ZONE 2

FOREST MANAGEMENT PLAN

2022 – 2026





Department of Fisheries, Forestry, and Agriculture Forestry & Wildlife Branch Regional Services

TABLE OF CONTENTS

INTRODUCTION 1	
SECTION 1 DESCRIPTION OF THE LAND BASE 2)
1.1 General)
1.1.1 Location)
1.1.2 History	3
1.1.3 Ownership	3
1.2 Physical	;
1.2.1 Topography and Hydrology3	3
1.2.2 Geology	3
1.2.3 Soils	ŀ
1.2.4 Climate	ļ
1.3 Ecosystems	ŀ
1.3.1 Forest Ecosystems	ŀ
1.3.2 Ecoregions and Subregions5	5
1.3.2.1 Central Newfoundland Forest Ecoregion5	;
1.3.2.1.1 North Central Subregion6	5
1.3.2.1.2 North Shore Ecoregion6	5
1.3.2.1.3 Eastern Hyper-Oceanic Barrens Forest Region6	5
1.3.2.2 Maritime Barrens Forest Ecoregion7	,
1.3.2.2.1 Northeastern Barrens Subregion7	,
1.3.2.2.2 Central Barrens Subregion7	,
1.4 Ecosystem Dynamics	3
1.4.1 Ecosystem Condition and Productivity 8	3
1.4.1.1 Productivity	3
1.4.1.2 Resilience)
1.4.1.3 Stability9)
1.4.1.4 Disturbance Regimes and Successional Patterns9)
1.4.1.4.1 Harvesting)
1.4.1.4.2 Fire)
1.4.1.4.3 Insect)
1.4.2 Biodiversity11	
1.4.2.1 Species Diversity11	-

1.4.2.2 Genetic Diversity	11
1.4.2.3 Ecosystem Diversity	12
1.5 Forest Characterization	12
1.5.1 Land Classification	12
1.5.2 Age Class	13
1.5.3 Site Class	14
1.5.4 Species and Working Group	15
1.5.5 Forest Disturbances	16
SECTION 2 PAST ACTIVITIES	17
2.1 Harvesting	17
2.1.1 Commercial	17
2.1.2. Domestic	18
2.2 Silviculture	19
2.3 Road Construction	19
2.4 Natural Disturbance	20
2.4.1 Fire	20
2.4.2 Insect	20
SECTION 3 TIMBER SUPPLY ANALYSIS	20
SECTION 3 TIMBER SUPPLY ANALYSIS	
	20
3.1 Introduction	20 20
3.1 Introduction 3.2 Guiding Principles and Policy Direction	20 20 21
3.1 Introduction3.2 Guiding Principles and Policy Direction3.3 Factors Affecting Timber Supply	20 20 21 21
 3.1 Introduction 3.2 Guiding Principles and Policy Direction 3.3 Factors Affecting Timber Supply 3.4 Timber Supply Analysis 	20 20 21 21 21 22
 3.1 Introduction 3.2 Guiding Principles and Policy Direction 3.3 Factors Affecting Timber Supply 3.4 Timber Supply Analysis	20 20 21 21 21 22 22
 3.1 Introduction 3.2 Guiding Principles and Policy Direction 3.3 Factors Affecting Timber Supply 3.4 Timber Supply Analysis	20 20 21 21 21 22 22 22
 3.1 Introduction 3.2 Guiding Principles and Policy Direction 3.3 Factors Affecting Timber Supply 3.4 Timber Supply Analysis 3.4.1 Forest Characterization 3.4.2.1 Non-Timber Related 3.4.2.1.1 No-Cut Buffer Zones 	20 20 21 21 22 22 22 22 22 23
 3.1 Introduction 3.2 Guiding Principles and Policy Direction 3.3 Factors Affecting Timber Supply 3.4 Timber Supply Analysis 3.4.1 Forest Characterization 3.4.2.1 Non-Timber Related 3.4.2.1.1 No-Cut Buffer Zones 3.4.2.1.2 Pine Marten and Caribou Habitat 	20 20 21 21 22 22 22 22 22 23 23
 3.1 Introduction	20 20 21 21 22 22 22 22 23 23 23
 3.1 Introduction 3.2 Guiding Principles and Policy Direction 3.3 Factors Affecting Timber Supply 3.4 Timber Supply Analysis 3.4.1 Forest Characterization 3.4.2.1 Non-Timber Related 3.4.2.1.1 No-Cut Buffer Zones 3.4.2.1.2 Pine Marten and Caribou Habitat 3.4.2.1.3 Protected Areas 3.4.2.2 Timber Related 	20 20 21 21 22 22 22 22 23 23 23 23 23
 3.1 Introduction 3.2 Guiding Principles and Policy Direction 3.3 Factors Affecting Timber Supply 3.4 Timber Supply Analysis 3.4.1 Forest Characterization 3.4.2.1 Non-Timber Related 3.4.2.1.1 No-Cut Buffer Zones 3.4.2.1.2 Pine Marten and Caribou Habitat 3.4.2.1.3 Protected Areas 3.4.2.2 Timber Related 3.4.2.2 Timber Related 	20 20 21 21 22 22 22 22 23 23 23 23 23
 3.1 Introduction 3.2 Guiding Principles and Policy Direction 3.3 Factors Affecting Timber Supply 3.4 Timber Supply Analysis 3.4.1 Forest Characterization 3.4.2.1 Non-Timber Related 3.4.2.1.1 No-Cut Buffer Zones 3.4.2.1.2 Pine Marten and Caribou Habitat 3.4.2.1.3 Protected Areas 3.4.2.2 Timber Related 3.4.2.2 Linsect/Fire/Disease Losses 3.4.2.2 Logging Losses 	20 20 21 21 21 22 22 22 23 23 23 23 23 23 23 23 23
 3.1 Introduction 3.2 Guiding Principles and Policy Direction 3.3 Factors Affecting Timber Supply 3.4 Timber Supply Analysis 3.4.1 Forest Characterization 3.4.2.1 Non-Timber Related 3.4.2.1.1 No-Cut Buffer Zones 3.4.2.1.2 Pine Marten and Caribou Habitat 3.4.2.1.3 Protected Areas 3.4.2.2 Timber Related 3.4.2.2 Timber Related 3.4.2.2 Logging Losses 3.4.2.2.3 Operational Constraints 	20 20 21 21 21 22 22 23 23 23 23 23 23 23 23 23 23 23

3.4.4.2 Spatial Analysis	24
3.4.4.3 Planning Horizons	25
3.4.4.4 Operable Growing Stock Buffer	25
3.4.4.5 Old Forest Targets	26
3.4.4.6 Operability Limits	26
3.4.4.7 Silviculture	26
3.5 Inventory Adjustments	27
3.5.1 Fire	27
3.5.2 Insects	27
3.5.3 Timber Utilization	27
3.5.4 Stand Remnants	27
3.6 Results	27
SECTION 4 VALUES	28
4.0 Guiding Principles of Sustainability	28
4.1 Value Structure	28
4.1.1 Biotic Values	30
4.1.1.1 Big Game	
4.1.1.1.1 Moose	
4.1.1.1.2 Caribou	
4.1.1.1.3 Black Bear	32
4.1.1.2 Furbearers	
4.1.1.3 Species of Interest	
4.1.1.3.1 American Marten	34
4.1.1.3.2 Rare Plants	35
4.1.1.3.4 Other Species	37
4.1.1.4 Water Resources	
4.1.2 Human Values	40
4.1.2.1 Timber Resource	40
4.1.2.2 Agriculture	41
4.1.2.3 Mining, Mineral Exploration and Quarrying	42
4.1.2.4 Historic Resources	45
4.1.2.5 Newfoundland T'Railway	46
4.1.2.6 Parks and Protected Areas	47
4.1.2.7 Outfitting	48

4.1.2.8 Tourism / Recreation	50
SECTION 5 MITIGATIONS	
5.1 General	
	54
SECTION 6 PUBLIC CONSULTATION	54
6.1 Planning Framework	54
6.2 Stakeholder Involvement	55
SECTION 7 MANAGEMENT OBJECTIVES AND STRATEGIES	57
7.1 Harvesting	57
7.1.1 Commercial	57
7.1.2 Domestic	58
7.2 Silviculture	58
7.3 Access Roads	60
7.4 Forest Protection	61
7.4.1 Insects and Disease	61
7.4.2 Fire	62
7.4.3 Windthrow	62
7.5 Environmental Protection & Climate Change	62
7.5.1 General Environment	62
7.5.2 Climate Change	63
7.5.3 Surveys	64
7.6 Information and Education	64
SECTION 8 PROPOSED ACTIVITIES	65
8.1 overview	65
8.1.1 Allocation of Timber Supply	65
8.1.1.1 Commercial	65
8.1.2.2 Domestic	68
8.1.2.3 Hardwoods	73
8.1.3 Silviculture	73
8.1.4 Primary Access Roads and Bridges	75
8.1.5 Activities in Protected Water Supply Areas	77
SECTION 9 PLAN ADMINISTRATION	77
9.1 Monitoring	77
9.2 Amendments	77

LIST OF FIGURES

Figure 1: Location of Planning Zone 2	2
Figure 2: Ecoregions and subregions of Planning Zone 2	5
Figure 3: Age class distribution for District 02	13
Figure 4: Age class distribution for District 03	14
Figure 5: Site class breakdown for District 02	14
Figure 6: Site class breakdown for District 03	15
Figure 7: Working group breakdown for District 02	15
Figure 8: Working group breakdown for District 03	16

LIST OF TABLES

Table 1: Land classification by district and area for Planning Zone 2.	12
Table 2: 2017- 2021 Commercial Harvest District 02	17
Table 3: 2017-2021 Domestic Harvest District 02.	18
Table 4: 2017-2021 Domestic Harvest District 03.	18
Table 5: 2017-2021 Silviculture treatments District 02	19
Table 6: 2017-2021 Road Construction District 02	19
Table 7: Annual Allowable Cut Zone 2 2016-2020	27
Table 8: Proposed forest harvest in Zone 2 from 2022-2026	65
Table 9: FMD 02 Proposed Commercial Harvesting from 2022 to 2026	66
Table 10: FMD 02 Proposed Domestic Harvesting From 2022 to 2026	68
Table 11: FMD 03 Proposed Domestic Harvesting From 2022 to 2026	72
Table 12: FMD 02 Proposed Silviculture treatment 2022-2026.	73
Table 13: FMD 02 Proposed Road Construction for 2022 to 2026.	75

INTRODUCTION

This new five-year plan is scheduled for the period January 1, 2022 to December 31, 2026 and represents proposed forestry activity upon crown timber lands within Forest Management Districts 02 and 03. The management of this land is consistent with strategies and philosophies implemented by Fisheries and Land Resources on all other crown land managed districts within the Province. This five-year operating plan incorporates established provincial planning requirements, environmental protection guidelines, and standard operating procedures developed under a stringent Environmental Management System (EMS) which is registered under the ISO 14001 standard. Topics that are provincial in scope such as carbon and global warming are included in the provincial sustainable forest management strategy, while sections that are more descriptive or depict local conditions such as values, forest characterization and ecosystem description are included into this five year operating plan. Forest Management Districts 02 and 03 are adjacent and share common ecoregion characteristics and collectively form Planning Zone Two (2). Within a planning zone, there is a requirement for each tenure to develop a five year operating plan. These plans have to be submitted to Forestry and Wildlife Branch and Department of Environment (for an environmental assessment review). As a result, there will be only one (1) five-year plan submissions for this zone. Throughout this five-year plan, references will be made to Districts 02 and 03 individually but when combined they will collectively be referred to as Planning Zone Two or the zone.

This document will attempt to fully integrate the presentation of information and discussions for crown land in the zone. Discussion and information will be presented separately for each district where warranted based on unique and distinct differences in scope and content. The more descriptive sections of this plan will be generic in nature and give information for the entire zone as well as some broad comparative statistics. Finally, this document will attempt to build on the positive results of previous five-year plan documents. Information will be updated as required or new sections will be added as any new information is available.

SECTION 1 DESCRIPTION OF THE LAND BASE

1.1 General

1.1.1 Location

Planning Zone two encompasses Forest Management Districts 02 and 03 (Figure 1). It is located in eastern Newfoundland and extends from Long Harbour River, Northwest River, and Terra Nova National Park in the west to the Come By Chance in the east and includes all of the Bonavista Peninsula, Burin Peninsula, and all the islands in Placentia Bay. Districts 02 and 03 are administered from Clarenville District office with satellite offices in Southern Bay and Winterland.




Figure 1: Location of Planning Zone 2

1.1.2 History

Since the earliest settlement, the forest and fish resources were the mainstay of the economy within this zone. Initially, the forest was used as a source of fuelwood and construction materials for houses and fishery related items (stages, lobster pots, boats etc.). The Bonavista Peninsula has one of the more active and successful agricultural communities in the Province. Tourism has been a growth industry in the region as well.

1.1.3 Ownership

The entire land area located in Planning Zone Two is currently managed by the crown.

1.2 Physical

1.2.1 Topography and Hydrology

The predominant physical feature throughout the District is rolling topography. Coastal hills dominate the landscape from Bull Arm north to Trinity and then strike north across the peninsula to Keels. The elevations of the hills on the peninsula are within a range of 150-250 metres above sea level, with the highest peak reaching 260 metres. The highest hills are to the west of Clarenville and strike southwest to Swift Current. These hills reach elevations of up to 400 metres. A relatively narrow plain is located in a transverse across the peninsula between Elliot's Cove and Goose Bay. This plain is underlain by Cambro Ordovician shales and contains most of the land within the District which is best suited for agricultural development.

A narrow swath of rolling topography runs parallel and to the south of the Trans Canada Highway between Port Blandford and Clarenville. West of this land form, the District spreads out to its western boundary in an expansive plateau of barren land and bog, with very intermittent forests occurring in narrow riparian areas and on valley slopes. Many small ponds are interspersed throughout this plateau and the eastern half of the peninsula.

1.2.2 Geology

Carbon aging indicates that it has been approximately 7400 years since the last glaciation occurred on the Bonavista Peninsula. The direction of ice flow was from west to east in the northern part of the District and from west/northwest to east/southeasterly in the southern portion. Glaciation was responsible for the rolling topography prevalent in much of the District and for the glacial deposits in some of the regions valleys. Climatic change caused a relatively rapid retreat of ice flows and the unstratified deposition of glacial till. Retreat of a later glacial advance during this period caused some out wash deposits to form. Out wash material (in the form of sand and gravel deposits) is present in various locations throughout the District. These deposits have been identified in the South West and North West Arm areas; west of Georges Brook; west of Lethbridge; in the Plate Cove area; south of the Northwest River and south of the Southwest River.

The type of rock, or parent material, underlying our forest soils is fundamental in the development of those soils. Some rock types, such as shales and slates, have many mineral elements which are essential to plant growth, and through geological time, become readily available. Generally, soils which form on parent material deriving from these type of rock group are fertile and can support favourable forestry/agricultural development. Geologic groups forming from igneous processes, such as granite, have far less inherent trace elements and generally lead to the development of soils in which fertility is a more limiting factor to plant growth. The geologic groups located on the Bonavista Peninsula are described in the soils report for that region by Heringa and Woodrow (1991). The peninsula contains the following geologic groups: Musgravetown, Adeytown, Connecting Point, Love Cove, Harcourt, and Granite Rocks.

1.2.3 Soils

Under the Canadian System of Soil Classification, Management District 2 has representation of three Orders of soil: podzolic, gleysolic and organic. The presence of shale parent material under many of the soils on the Bonavista Peninsula has helped build nutrient rich forest soils. As a result, many sites in this area are highly suitable for the development of productive forest or for agricultural development.

1.2.4 Climate

District 2 is located in the southeastern region of the province. Bordering the Atlantic Ocean on the eastern extremity, its climate is largely influenced by maritime conditions. Seasonal temperatures are moderated by this influence, yet conditions during certain seasons, especially the beginning of the growing season, can be harsh due to the cooling effect from the ice laden Labrador Current. The average mean daily February temperature in the District ranges from -5 0C to -8 0C with the minimum being -140C. The average mean daily July temperature ranges from +13 0C to +160C with the maximum being + 22 0C. The growing season in the District ranges between 140 and 160 days. Average precipitation ranges from 900mm - 1300mm and average snowfall ranges from 2.5 to 3.5m.

1.3 Ecosystems

1.3.1 Forest Ecosystems

The tree species which characterize the Canadian boreal forest include black spruce, white spruce, balsam fir, eastern larch, trembling aspen, white birch and jack pine. All of these, with the exception of jack pine, commonly occur on the Island. However, by far the dominant species are black spruce and balsam fir; together they represent more than 90 percent of the growing stock on the island.

1.3.2 Ecoregions and Subregions

Damman 1979, defined ecoregions as areas where comparable vegetation and soil can be found on sites occupying similar topographic positions on the same parent material, provided that these sites have experienced a similar history of disturbance. Thus, an ecoregion cannot be defined in isolation from the physical landscape, but vegetation toposequence, vegetation structure, floristic composition, and floristic distributions can provide the primary criteria. According to Damman, nine ecoregions are represented in Newfoundland. Each of these is further divided into subregions (also known as ecodistricts) All of the Newfoundland ecoregions and subregions contain many of the same ecosystem variables. It is the dominance and variance of these variables (e.g., vegetation and climate) that determine their classification.

Figure 2 depicts Planning Zone 2 relative to Damman's ecoregion classification system. FMD2 is located in a transition zone between central ecoregions, which have been influenced by interior continental-like climatic conditions, and eastern ecoregions, which have a greater coastal climate influence. There are portions of two subregions of the Maritime Barrens Ecoregion located in FMD2, including the Northeast Barrens Subregion and the Central Barrens Subregion. In addition, the Eastern Hyper - Oceanic Barrens Ecoregion crosses the northeastern tip of the peninsula.




Figure 2: Ecoregions and subregions of Planning Zone 2

1.3.2.1 Central Newfoundland Forest Ecoregion

This ecoregion is located in the north-central part of the island with a small outlet near Bay d'Espoir. The topography is gently rolling to hilly with most elevations between 150 and 450 meters. It has the most continental climate in insular Newfoundland with the warmest summers and coldest winters. It has the least wind and fog of any ecoregion and a growing season of 140-160 days and average precipitation of 900-1300mm. This ecoregion is heavily forested and is the most distinctly boreal part of the island. Balsam fir, black spruce, and to a lesser extent white birch

are the dominant tree species. There is an extensive fire history thus fire origin stands of black spruce and white birch cover extensive areas in the northern and eastern portions. Trembling aspen forms local stands after fire but is restricted to the central and northern portion. *Hylocomium*-balsam fir is the zonal forest type and is dominant in areas not disturbed by fire. *Kalmia*-black spruce and *Pleurosium*-balsam fir forests are also common. The *Kalmia*-black spruce-lichen forests, which occur on outwash sands and gravels, are unique to this ecoregion. Red pine also occurs but is restricted to extremely dry sites.

1.3.2.1.1 North Central Subregion

The North Central Subregion has the highest maximum temperatures, lowest rainfall, and highest forest fire frequency on the island. The subregion extends from Clarenville to Deer Lake with a mostly rolling topography of less than 200 meters. The history of fire is evident by the pure black spruce forest with white birch and aspen stands that dominate the subregion.

1.3.2.1.2 North Shore Ecoregion

The North Shore Forest Ecoregion does not have any subregions. It forms a coastal band approximately 20-25 kilometres wide extending from the Bonavista Peninsula in the east to the Baie Verte Peninsula in the west. The North Shore Forest Ecoregion occupies the northern half of the Bonavista Peninsula, bounded by the North Eastern Barrens Subregion in the south and the North Central Forest Subregion in the west. Black Spruce and Balsam Fir forest form a continuous forest except where barrens dominate on coastal headlands. The vegetation growing season is shorter and cooler than in the central part of the island, but it also has a frost free season which is several weeks longer. The summers are relatively dry and warm and soil moisture deficiencies may occur. Similar to the Central Newfoundland Forest Ecoregion, encroachment of ericaceous shrubs on drier nutrient poor sites after fire or cutting disturbance is common and presents a serious silvicultural problem. This problem is more prevalent after cutting.

1.3.2.1.3 Eastern Hyper-Oceanic Barrens Forest Region

The Eastern Hyper Oceanic Barrens Ecoregion occurs on the extreme south coast of the Avalon and Burin Peninsulas and in the Bay de Verde, Cape Freels and Bonavista areas. In District 2, the Eastern Hyper Oceanic Barrens Ecoregion is located on the very tip of the Bonavista Peninsula. Although at low elevation, this ecoregion has very cool summers due to the oceanic influence. The landscape is dominated by exposed coastal barren and blanket bogs and is completely without forest cover, except for Balsam Fir krummholz (ie. tuckamoor or low windswept scrub) (Figure 2.20). Arctic-alpine species occur even at sea level and are mixed with species common to southern coastal plain. This is a very unique ecological aspect of the oceanic barrens.

1.3.2.2 Maritime Barrens Forest Ecoregion

The Maritime Barrens Ecoregion extends from the eastern Peninsulas westward through southcentral Newfoundland to Port-aux-Basques. It is divided into four subregions, two of which are partially located in FMD2. The Maritime Barrens Ecoregion is characterized by cool, foggy and windy summers and relatively mild winters. Intermittent snow cover is common within the coastal portions of this ecoregion but increases in accumulation and duration in the interior barrens. The landscape pattern is one of almost pure Balsam Fir interspersed throughout extensive open heath-land. Productivity of the forest is better on long slopes which occur in infrequent valleys. A natural history of frequent wildfire occurrence in much of this ecoregion has had a significant influence in the development of this largely heath-covered landscape.

1.3.2.2.1 Northeastern Barrens Subregion

The North Eastern Barrens Subregion occupy the eastern third of the District. It is, generally speaking, that area east/southeast of a line running from Knights Cove southwest to Harcourt; and then south along the western end of Random Island and the bottoms of Northwest Arm and Southwest Arm to Queen=s Cove, and then southwest to Swift Current. This subregion has a lower frequency of fog and warmer summers than the other subregions of the Maritime Barrens Ecoregion, except for the Central Barrens Subregion. Portions of the landscape are, in fact, heavily forested - but these portions are interspersed throughout large expanses of soil/rock barrens, softwood scrub forest and local heath vegetation, usually along the coast. Balsam fir is the predominant tree species that occurs in the North Eastern Barrens Subregion. Natural regeneration of forests usually occurs after natural or cutting disturbance in this subregion. Fire is less prevalent than in other parts of the District. Natural forest succession usually occurs as a result of stand replacement insect or wind disturbances.

1.3.2.2.2 Central Barrens Subregion

This area occurs south of the Central Newfoundland Forest Ecoregion and north of the South Coast Barrens Subregion. In FMD2, the Central Barrens Subregion extends from the Northwest River south to Black River Pond in a strip of land bounded by the district boundary on the west and the Central Newfoundland Forest Ecoregion on the east. This subregion has warmer summers, less fog and more persistent snow cover then the other subregions in the Maritime Barrens Ecoregion. Forest patches, predominately composed of Balsam Fir Forest, are common throughout the barrens. Fire has played a significant role in the development of the ecology of the Central Barrens Subregion. Within FMD2, this subregion is unique in that it provides the eastern range for the Middle Ridge caribou herd.

1.4 Ecosystem Dynamics

1.4.1 Ecosystem Condition and Productivity

Landscape patterns determine the variety, integrity, and interconnectedness of habitats within a region. These landscape patterns are a direct result of the relationship amongst physical landforms and soils, disturbance history, and relationships among various species that make up the ecosystem communities. These factors, while listed separately for clarity, are unavoidably interrelated. Landscape patterns play a pivotal role in determining the current conditions and health of forest ecosystems. These variables are evaluated in terms of productivity, stability and resilience.

Another important role determining the condition of a forest is change. Forests are an ever evolving entity, resisting stagnation, and constantly moving through their cycles of life, death, and renewal. The process of change over time is the essence of nature itself. It has been nature's underlying storyline since time began, and will continue to be until time ends. The main forces of change in our natural forest ecosystems are disturbance and succession. A definition of disturbance would indicate that it initiates a change in a community structure which often ends up in the replacement of one set of species by another. However, replacement is not always the end result (e.g., a species like black spruce is aided in germination by disturbances like forest fire).

Disturbances range from the fall of a single tree, to the destruction of thousands of hectares by forest fires. While disturbances may be very destructive, they can often rejuvenate ecosystems and diversify landscapes.

Succession involves changes in both community composition and in the ecosystem structure and process. Succession is the orderly change whereby the dominant species is replaced by another species, then another etc. until a new dominant species establishes a relatively stable community.

1.4.1.1 Productivity

Productivity is the accrual of matter and energy in biomass. In simple terms, primary productivity is the sum total of all biomass produced through photosynthesis. Secondary productivity occurs when this "primary" biomass is ingested and is added to that organism's biomass. Since secondary productivity is directly dependent on primary productivity, it is this primary productivity component that drives the system. The level of primary production is dependent on the ability to produce biomass. This in turn is dependent on landscape features, soil, climate etc. In general terms, the more productive (ability to grow trees) a site is, the higher level of primary productivity. For example a forested stand would have a higher primary productivity than a bog and a good site would have a higher potential than a poor site.

Overall, the landscape in FMD2 has approximately 37 percent productive forest. As well, the relative proportion of site types is 2 percent good, 73 percent medium and 25 percent poor with a mean annual increment (MAI) of 2.6, 1.7, and, 0.8 m3/ha/yr respectively. This distribution of productive sites across the landscape and range of productivity within these sites is largely dependent on landscape patterns, climate, and soils. The more productive areas of FMD2 occurs in the lowlands of the river valleys. These areas have deeper soils and less exposed bedrock.

The landscape patterns are more consistent and the growing season is longer. In contrast, the eastern parts of FMD2 along the coast have soils are shallower with bedrock at or near the surface. The terrain in eastern parts is much rougher and the growing season is shorter.

1.4.1.2 Resilience

Ecosystem resilience reflects the ability of the ecosystem to absorb change and disturbance while maintaining the same productive capacity and the same relationships among populations. Healthy forest ecosystems maintain their resilience and adapt to periodic disturbances. The renewal of boreal forest ecosystems often depend on these disturbances. Resilience is characterized by the forest's ability to stabilize vital soil processes and maintain succession whereby the system is returned to a community composition and the productivity level is consistent with the ecosystems physical constraints following a disturbance. To a large degree, a forest ecosystem's resilience is controlled by properties such as climate, parent soil, topography and flora.

1.4.1.3 Stability

Nature is constantly changing and going through the unending processes of disturbance, growth, senescence, and decay. Therefore, stability of a forest ecosystem does not refer to one fixed position without variation. Ecosystem stability is more accurately defined as the maintenance of ecosystem changes within certain boundaries and the functional continuation of important potentials and processes such as energy capture.

There are three levels of stability; species stability, structural stability, and process stability. Species stability is the maintenance of viable populations or meta-populations of individual species. Structural stability is the stability of various aspects of ecosystem structure such as food web organization or species numbers. Process stability is the stability of processes such as primary productivity and nutrient cycling. To put stability in perspective, it must ensure that the system does not cross some threshold from which recovery to a former state is either impossible, (extinction) or occurs only after long time periods or with outside inputs (eg. loss of topsoil).

Some indicators of stability which can be monitored are: area of forest converted to non-forest use, area, percentage and representation of forest types in protected areas, percentage and extent of area by forest type and age class, and change in distribution and abundance of various fauna. These indicators can be measured and monitored to ensure stability is maintained and to evaluate the impact, if any, of forest activities on ecosystem stability.

1.4.1.4 Disturbance Regimes and Successional Patterns

There are four main driving forces that cause disturbance in the boreal forest. Forest Harvesting can be considered a major disturbance in the zone, occurring on a regular and consistent basis. Fire and insect damage are the other two major disturbances and occur on a more irregular or cyclic basis. With the exception of a major windstorm, wind throw usually occurs after a stand is weakened by some other agent like insects. For this reason successional patterns after insect damage and wind throw will be discussed together. The following is a brief synopsis of successional patterns after each major disturbance type by forest type and site type.

1.4.1.4.1 Harvesting

Regeneration patterns in the black spruce type after harvesting is generally back to the black spruce type with a minor component of balsam fir and some white birch on the better sites. There are two main white birch site types in the zone. The basic difference between them is terrain which impacts site quality. These sites are prone to revert to alder dominated NSR sites in the absence of very hot ground fire as the disturbance mechanism. Consequently the management prescription to ensure productivity on these valuable sites is to plant fast growing softwood species. The medium white birch sites are typically on more level terrain and will revert to white birch /balsam fir or white birch/black spruce after disturbance. Regeneration failure on these sites is low (10 percent). The management prescription to regenerate these site to white birch is to remove the overmature birch in a seed tree cut to provide a seed source for the next rotation of birch.

Harvesting of white birch in this zone has traditionally been for firewood purposes Recently, however, some of the harvest occurring has been directed to sawmilling with the development of a value added hardwood industry, which will place added pressure on the white birch resource in the zone. Evidence from domestic cutting in these types indicates that they will regenerate to mixed wood types dominated by balsam fir and white birch.

1.4.1.4.2 Fire

Since black spruce is a fire adapted species, it is not surprising that it is the most prolific regeneration species after fire across all forest types, site types and ecoregions within the zone. It regenerates as pure stands or in combination with white birch. Balsam fir is conspicuously absent after fir because most advanced regeneration in the under story is killed by the fire. Black spruce regeneration is somewhat correlated with the amount present in the pre fire stand. Generally, the higher the component of black spruce in the original stand, the higher the percentage of regeneration to black spruce. In mixed wood stands a higher component of white birch and sometimes trembling aspen is present after fire. Fire in pure hardwood stands can sometimes regenerate to trembling aspen in certain areas. Regeneration failure after fire is on average 20 percent across all forest types and is higher as sites get poorer.

1.4.1.4.3 Insect

Balsam fir is highly susceptible to insect attack from the hemlock looper, balsam woolly adelgid, balsam fir sawfly, and spruce budworm whereby black spruce is hardly impacted by these insects. For this reason, stands with a high component of balsam fir are more susceptible to insect attack and subsequent wind thrown. Mature balsam fir types usually regenerate to balsam fir or to balsam fir hardwood mixtures. Disturbance by insect kill in young balsam fir stands can cause succession to white spruce. Regeneration patterns in mixed wood types usually depend on the type of mixture. If black spruce is a component then it will persist and form part of the new stand. Otherwise balsam fir and balsam fir/hardwood mixtures regenerate after insect attack.

Regeneration failure occurs approximately 20 percent of the time particularly if pure stands of immature balsam fir are killed.

1.4.2 Biodiversity

Biodiversity is a term used to describe the variety of life on earth. A basic definition of biodiversity includes the variety of animals, plants and microorganisms that exist on our planet, the genetic variety within these species and the variety of ecosystems they inhabit.

While the boreal forest may not have the extent of biodiversity that some of the equatorial regions possess, Canada does have just over 70 000 species of plants, animals, and micro organisms in its boreal and other forest regions. While the boreal forest has less diversity of large plants than many other forest regions, it has greater biological diversity in some micro organisms. For example, the boreal forest has fewer tree species than the tropical rainforest but potentially up to 500 times as many mycorrhizal fungi. Despite the large number of organisms contained within the boreal forest, only a small amount are actually plants and vertebrates. The larger portion remains largely unrecorded and unstudied. As a result, we need to manage with caution so that species are not inadvertently extirpated.

Biodiversity provides such essential services for humans as: climate control, oxygen production, purification of freshwater supplies, carbon dioxide removal from the atmosphere, soil generation, and nutrient cycling.

The three components of biodiversity are species diversity, genetic diversity, and ecosystem diversity.

1.4.2.1 Species Diversity

Species diversity describes the overall range of species in a given area or ecosystem. Species are groups of animals, plants, and micro organisms capable of producing fertile offspring. Species extinction is the most dramatic and recognizable form of reduced biodiversity; habitat loss the most drastic in terms of far reaching effect. The prevention of species extinction is a key factor in the conservation of biodiversity. Changes in species population levels indicate the potential for serious changes in ecosystem integrity.

1.4.2.2 Genetic Diversity

Genetic diversity describes the range of possible genetic characteristics found within and among different species. Hair and eye colour, weight and height, are examples of genetic diversity found in humans. Genetic diversity within species is the foundation of all biodiversity. Assessing genetic diversity does not mean tracking every gene in the zone's forest. Responsible planning should design and implement measures which maintain or enhance viable populations of all forest vegetation species and which use the genetic diversity of commercially important species to a maximum benefit. The genetic diversity of commercially important species can also be managed to increase economic benefit from some portions of the landscape while allowing other portions to provide greater social and ecological values. Genetic diversity is the basis by which populations (flora and fauna) can adapt to changing environmental conditions.

1.4.2.3 Ecosystem Diversity

Ecosystem diversity describes the range of natural systems found throughout a region, a country, a continent, or the planet. Wetlands and grasslands are examples of ecosystems in Canada. A complex and intricate mix of plants, animals, micro organisms and the soil, water, and air they occupy create virtually limitless ecosystems around the world.

A forest interspersed with barrens, marshes, lakes and ponds provides for diversity across the landscape. Each ecoregion in the province should have representative areas protected which displays the diversity where such exists. These areas can serve as a benchmark from which to measure and guide management decisions. These representative areas protect the integrity of the ecoregion and are vital for guiding management actions. As benchmark areas, they will illustrate the multi-species mosaic that planning actions must maintain.

1.5 Forest Characterization

1.5.1 Land Classification

Table 1 displays the land classification broken down by ownership and district for Planning Zone 02. The total mapped land area in the zone is approximately 776,495 hectares. There are four basic categories that currently represent how the land is classified; productive, non productive, non-forest and fresh water.

Table 1: Land classification by district and area for Planning					
	Dist	Total			
Land Class	2	3			
disturbed	7921	2469	10390		
0-20 years	29168	8065	37233		
21-40 years	23489	14766	38255		
41-60 years	7315	13952	21267		
61-80 years	5427	13868	19295		
81-100 years	16804	8506	25310		
101-120 years	6699	3411	10110		
120+ years	4470	6893	11363		
Total Productive	101293	71930	173223		
softwood scrub	95082	61588	156670		
hardwood scrub	3722	5458	9180		
Total Non-Productive	98804	67046	165850		
non-vegetated	39370	102147	141517		
vegetated non-forested	61338	108468	169806		
clearedland	614	319	932		
agriculture land	236	3 60	596		
residential	2634	2212	4846		
right of ways	809	679	1487		
resource roads	378	13	391		
Total Non Forested	105378	214197	319576		
Fresh Water	39370	78477	117847		
Total All Classes	344845	431650	776495		

Table 1: Land classification by district and area for Planning Zone 2.

1.5.2 Age Class

Individual tree ages within any given stand have the potential to be the same after fire or planting. However; in most cases the ages vary. Foresters describe forest stand age in terms of age classes which generally encompass 20 years. The age classes present in the zone are described as regenerating (age class 1, 0-20 years), immature (age class 2, 21-40 years), semi-mature (age class 3, 41-60 years), mature (age class 4, 61-80 years), and over mature (age class 5, 81-100 years), (age class 6, 100-120 years), (age class 7, 120+ years). The age class distribution in each district for the entire productive forest is shown in Figures 3 and 4. In general terms, a continuous timber supply is limited by the lower age class. This means a more balanced age class distribution within a district would yield a greater opportunity for an even flow sustained yield of timber.




Figure 3: Age class distribution for District 02.




Figure 4: Age class distribution for District 03.

1.5.3 Site Class

The productive forest in the zone is further sub-divided along a gradient of productivity ranging from poor to good site class. The site class is determined through air photo interpretation supplemented with field checks, and is based primarily on the sites ability to produce timber. Site capability is determined on a number of factors including: soil fertility, moisture regime and geographic (slope) position. The distribution of area by site class for each district is shown in Figures 5 and 6. On average, good sites are capable of producing 2.6 m3/ha/yr, medium sites 1.7 m3/ha/yr, and poor sites 0.8 m3/ha/yr.




Figure 5: Site class breakdown for District 02.




Figure 6: Site class breakdown for District 03.

1.5.4 Species and Working Group

A Working group is a term used to describe the dominant tree species present in a forest stand. This species may occupy 100 percent of crown closure of a stand or may be present in association with other species. The working group designation describes the stand in general terms based on the prevalent species as opposed to species composition which specifically describes the relative proportion of each individual tree species that make up a stand.




Figure 7: Working group breakdown for District 02.




Figure 8: Working group breakdown for District 03.

1.5.5 Forest Disturbances

Forest harvesting, fire, insects and windthrow (blowdown) are considered forest disturbance types within the zone. The Bonavista Peninsula is very representative of these disturbance patterns. Historical records reveal that virtually all of the forest area has been swept by wildfires. Most of the FMD2 productive timberlands in existence at the turn of the century were destroyed by large forest Fires which occurred during the late 1880's and early 1900's (1907).

Wind damage has been witnessed to occur infrequently in naturally developing stands that have not had other disturbance impacts. Insect damage during the late 1970's and early 1980's and the widespread practice of partial stand cutting has contributed to subsequent wide-spread wind throw throughout much of the District. Recent weather events, including ice storms in 2008 and 2009 and hurricane IGOR in 2010, have exacerbated the situation and have resulted in extensive damage throughout FMD2.

SECTION 2 PAST ACTIVITIES

2.1 Harvesting

2.1.1 Commercial

Tables 2, summarizes the commercial and some domestic harvest in the zone for the period 2017-2021.

Dist	rict 02		Core Land	oase		Ор	erational La	andbas	e	Non-AA(C Wood
	Crown	AAC	Commercial		Total	AAC	Commercial		Total	Operational	Regulatory
	2017	60,840	44,888			8,112				0	209
	2018	60,840	53,501			8,112					731
Q	2019	60,840	56,144			8,112					0
SWD	2020	60,840	49,532			8,112					2,589
	2021	60,840				8,112					0
	Sub-Total	304,200	204,065			40,560	0			0	3529
		Core Landbase			Operational Landbase			Non-AAC Wood			
	Crown	AAC	Commercial		Total	AAC	Commercial		Total	Operational	Regulatory
	2017	2,161	3,985			234				0	0
	2018	2,161	3,908			234				0	3
HWD	2019	2,161	3,874			234					0
	2020	2,161	3,792			234					174
	2021	2,161				234					0
	Sub-Total	10,805	15,559			1170	0			0	177
Distr	ict Total									0	0

Table 2: 2017- 2021 Commercial Harvest District 02

2.1.2. Domestic

Tables 3 & 4 summarizes the Domestic harvest in the zone on Domestic Landbase for the period 2017-2021

District 02 Domestic Landbas		base		
Crow	'n	AAC	Domestic	
	2017	32,292	30,036	
	2018	32,292	32,967	
SWD	2019	32,292	42,167	
SV	2020	32,292	30,641	
	2021	32,292	N/A	
	Sub-Total	161,460	135,811	
		Domestic Landbase		
	Crown	AAC	Domestic	
	2017	663	1,655	
	2018	663	1,465	
DWH	2019	663	1,698	
	2020	663	N/A	
	2021	663	N/A	
	Sub-Total	3315	4,818	
District Tota	I			

Table 3: 2017-2021 Domestic Harvest District 02.

Table 4: 2017-2021 Domestic Harvest District 03.

District 03		Domestic Landbase		
Crown	Crown AAC Dom			
	2017	13,391	15,984	
	2018	13,391	13,400	
Q	2019	13,391	16,151	
SWD	2020	13,391	16,370	
	2021	13,391	N/A	
	Sub-Total	66,955	61,905	
		Domestic Landbase		
	Crown	AAC	Domestic	
	2017	N/A	130	
<u> </u>	2018	N/A	516	

	2019	N/A	781
	2020	N/A	728
	2021	N/A	N/A
	Sub-Total		2155
District Total			

2.2 Silviculture

Table 5 summarizes the completed silviculture treatments in the zone for the period 2017-2021

Table 5: 2017-2021 Silviculture treatments District 02.

Tractment Type	Area (ha)			
Treatment Type	Proposed	Treated		
Pre Commercial Thinning	0	0		
Site Preparation (raking)	370.2	342.8		
Planting	773.87	722.85		
Commercial Thinning	0	0		
Prescribed Burning	0	0		

2.3 Road Construction

Tables 10 & 11 summarizes forest access road construction in the zone for the period 2017-2021

Table 6: 2017-2021 Road Construction District 02

Roads				
	Proposed (km)	Constructed (km)		
New Construction	33.00	17.29		
Re-Construction	7.00	4.27		
Total	40.00	21.56		
Bridges	0.00	0.00		
Operator Proposals	33.85	22.37		

2.4 Natural Disturbance

2.4.1 Fire

This zone has had a very infrequent fire history due to its relatively long winters and abundant precipitation. There were no significant fires during the last planning period.

2.4.2 Insect

There was no defoliation and treatment for either the hemlock looper or balsam woolly adelgid in the last 5 years.

SECTION 3 TIMBER SUPPLY ANALYSIS

3.1 Introduction

The province conducts a review of timber supply every five years to reflect any changes in forest land base, growth rates, and management strategies. This schedule is consistent with the Forestry Act, with oversight by forest management districts, and mandates a wood supply analysis to be completed every five years. The result of this analysis is the establishment of annual allowable cuts (AAC's) for each forest management district. These AAC's are defined as the maximum annual rate at which timber can be harvested at a sustainable level into the future (applicable for a period of 160 years). Annual allowable cuts must be calculated on a district basis, and the cumulative sum would provide the total island annual allowable harvest level. The current Wood Supply is for the period January 1, 2016 to December 31, 2020. A new woodsupply analysis is currently being conducted for the period January 1, 2022 to December 31, 2026. More information on the Timber Supply Analysis Program can be found on Governments Forestry website using the following address:

https://www.faa.gov.nl.ca/forestry/timber/index.html

3.2 Guiding Principles and Policy Direction

The key underlying principles guiding the provincial woodsupply analysis are:

- (i) the AAC must be sustainable;
- (ii) the level of uncertainty (risk) associated with the AAC must be minimized by using empirical information wherever possible;
- (iii) there must be conformity between information and assumptions used in the analysis and actions and decisions taken on the ground;
- (iv) the analysis must be consistent with other forest values and objectives; and
- (v) the timber supply calculation must consider economic factors, not solely the physical supply of timber.

In addition to the establishment of sustainable timber harvest levels, the legislation also requires that forest harvesting not exceed the established AAC's. Likewise, government's policy is to optimize forest industry opportunities from the sustainable fiber supply. Government also requires consultation be conducted during the timber analysis. In the current analysis, the forest industry was consulted directly. As well, there was a 30 day consultation process whereby a draft of the gross AAC's and methodology was published on the government web site for public review and comment.

3.3 Factors Affecting Timber Supply

The forests of insular Newfoundland are very variable in terms of age distribution. Typically, there are significant amounts of mature/over-mature forest and regenerating forest, and limited intermediate aged forests. This imbalance is not unusual in a boreal forest where cyclic catastrophic disturbances are common.

This imbalanced age class structure of intermediate age forest within insular Newfoundland is one of the most important factors influencing AAC's and is therefore the basis for many of the department's forest management strategies. Essentially, the department utilizes a matrix of management techniques designed to marginalize the imbalance in age structure. These techniques range from an aggressive forest protection program (insect control and fire suppression), forest harvesting programs that attempt to exclusively target harvesting the oldest stands first, and pre-commercial thinning of the regenerating forest so that it becomes merchantable and ready for harvest at an earlier age.

Another important aspect of the province's forest that poses a challenge to forest managers is the natural fragmentation of the resource. The province's landscape is carved by many ponds, bogs, rivers, streams, and rock outcrops resulting in relatively small pockets of timber scattered across the landscape. These adverse conditions is very challenging when determining the economic availability of timber supply.

Arguably, the most important factor affecting present and future AAC's is the land base. The land base available for forest activity is constantly being reduced as a result of other users' requirements. There is an approximate correlation between AAC and land base in that a one percent loss of land base represents a one percent drop in AAC. Therefore, it is very important that we continue to determine methods to minimize the loss of productive landbase and expand on efforts to grow more volume on the existing land base.

3.4 Timber Supply Analysis

The timber supply analysis is structured to determine sustainable timber availability, while respecting social, economic and environmental objectives. Timber supply, in this context, refers to the rate at which timber is made available for harvesting on a sustainable basis.

The determination of supply (represented as AAC's) involves the use of computer models to forecast the sustainability of possible AAC levels. These models require three basic inputs as described below:

(1) a description of the current state of the forest (forest characterization and availability),

- (2) growth rates associated with the current forest, and
- (3) management strategies applied to the forest.

These basic inputs requires careful and detailed consideration of a broad range of both timber and non-timber values. The following topics in this chapter are considered when determining the sustainable timber supply.

3.4.1 Forest Characterization

To realize the current description of the forest resource (referred to as forest stock), the province has invested significant resources into creating and maintaining a Provincial Forest Inventory. This program is designed to ensure the estimate of forest stock is current and accurate, while other regular programs employed by the department also evaluates:

- 1. Natural and man-made disturbances (fire, insects and harvesting)
- 2. Enhancement activities (tree planting and pre-commercial thinning)
- 3. In addition, the actual stands within the forest inventory is updated to reflect any yield changes

3.4.2 Land Availability

Through a regular timber supply analysis, the Forest Inventory is updated and classified at the stand level on the basis of harvest potential. This classification system consists of three broad classes;

- i. Class 1 available for harvest under normal operating conditions
- ii. Class 3 adverse conditions for forest harvesting, making that landbase more expensive and less available under current economic conditions. However, an AAC is still calculated in the event of improved economic conditions making the area more feasible for commercial operations.
- iii. Class 5 unavailable for forest harvesting. No AAC is calculated on this landbase, which incorporates a broad range of timber and non-timber values as indicated in the following sections.

3.4.2.1 Non-Timber Related

Implementation of non-timber values has a direct impact on provincial AAC's. As the amount of productive forested landbase available for timber management declines, so will the AAC. With the current non-timber related considerations, the net landbase (area where harvesting operations can occur) is only 17% of the total landmass on the island or 66% of the total productive forest land base. Typically, in any given year, less than 1% of the productive forest land base is influenced by harvesting operations.

3.4.2.1.1 No-Cut Buffer Zones

The Department has implemented guidelines requiring all water bodies (visible on a 1:50,000 map sheet) be given a minimum 20 meter (from the edge of water) unharvested buffer. In addition to these legislated water buffers, District Ecosystem Managers, in consultation with interested stakeholders may have increased buffer zone widths to protect special values such as; salmon spawning areas, cabin development areas, aesthetic areas, wildlife habitat, outfitting camps, etc.

3.4.2.1.2 Pine Marten and Caribou Habitat

Wildlife Habitat specialists are working in consultation with industry to ensure future adequate habitat remains available for wildlife species such as pine marten and caribou. Analysis of the landbase continues with examining the quantity and quality of habitat, as well as, the required connectivity. Wildlife specialists also examine how this arrangement of habitat would change over time. Forest management strategies take into consideration the results and recommendations of the Wildlife Habitat Specialists.

3.4.2.1.3 Protected Areas

All established and proposed protected areas approved within the Natural Areas Systems Plan (NASP) are removed from potential harvest considerations and the AAC calculations.

3.4.2.2 Timber Related

The potential AAC within a Forest Management District is also further impacted by taking into account other potential losses of landbase or timber as indicated below:

3.4.2.2.1 Insect/Fire/Disease Losses

The department reduces AAC's to account for anticipated future losses resulting from insects, disease and fire using historical information.

3.4.2.2.2 Logging Losses

Surveys of recent harvested areas are conducted each summer throughout the province to determine the quantity and quality of fiber remaining. The estimates of loss from these surveys are used to reduce the AAC.

3.4.2.2.3 Operational Constraints

Areas that are inaccessible (surrounded by bogs or hills), timber on steep slopes, and low volume stands are removed from the class 1 AAC calculation. Also, significant adjustments are applied to the provincial forest inventory for stands deemed operable in the timber analysis but left unharvested within operating areas. The reasons for this are linked to the character of Newfoundland's forests; low volume, steep slopes, rough terrain, and excessively wet ground conditions etc. All these timber and non-timber related issues are applied directly in the AAC calculation to ensure harvest levels do not exceed the sustainable level. With the introduction of new values and the broader application of current values, negative pressure on future AAC's will continue to increase.

3.4.3 Growth Forecasting

A key requirement for forecasting future wood supply is an understanding of how forest stands grow and develop through time. That is, as a forest stand develops, how much merchantable (i.e. harvestable) volume does it carry at any given point? These yield forecasts (referred to as yield curves) are required for each type of forest stand (called a stratum) comprising the forest. In Newfoundland, there are dozens of distinct forest strattum for which separate yield curves are required. These are defined by the tree species in question (e.g., balsam fir, black spruce), the

site quality (e.g., good, medium, poor), the geographic region (e.g., the Northern Peninsula, Western Newfoundland) and other factors likely to affect yield.

Yield curves are a key element in a wood supply analysis. In fact, the validity, or "usefulness", of the wood supply analysis is determined by the truth, or "correctness", of the yield forecasts. While there is no way of predicting with one hundred percent certainty how stands will actually grow in the future, care must be taken to ensure that the yield projections are realistic and reasonable. Respecting the sensitivity and importance of these forecasts, the department has directed a large portion of its resources and time into developing realistic yield curves. Two growth models were used, one for projecting stand development under natural conditions and the other for projecting growth under managed (i.e., silviculturally enhanced) conditions. Tree and stand development data generated from the departments forest inventory program were used to make stand growth predictions. These projections were then evaluated against empirical data from thousands of temporary plots established throughout the island. If the projections varied from the real life evidence, the curves were adjusted to make them more accurate.

In this analysis, yield curves were developed on an ecoregion basis. As well, special yield curve sets were developed for defined geographic areas with demonstrated uniqueness. These included areas where chronic insect activity is ongoing and areas that have unique growth characteristics.

3.4.4 Management Strategies

With the current state of the forest described and the yield forecasts developed, the next step was to design a management strategy for each sector of the forest. The key objective was to maximize long term AAC while at the same time taking into account other forest values. This involved developing strategies that minimize fiber losses, and enhance forest sustainability.

3.4.4.1 Harvest Flow Constraints

An even-flow harvest constraint strategy is utilized in the wood supply analysis. This strategy produces the maximum even flow harvest but results in less than optimum economic use of the forest resource. Conversely, if this strategy was not applied, then harvest levels are permitted to fluctuate which may result in increased commercial potential of the forest at specific intervals. However; applying the even-flow constraint provides more stability within the forest industry.

3.4.4.2 Spatial Analysis

The provincial wood supply analysis implements a technique of manual harvest scheduling. In 2001, the harvest scheduling was an automated process where the software allocated the stands to be harvested over the upcoming 25 years, based on user supplied criteria. The 2001 approach of scheduling harvest stands was an improvement over previous wood supply analysis. However, the software used cannot realistically know all the operational restrictions within a forest management district. By utilizing the manual process, District Staff are able to identify specific ground conditions that restrict commercial harvesting, which are then incorporated into a spatial harvest schedule. The proposed harvest schedule is then vetted back through the modeling software to ensure sustainable and non-timber objectives are met. In most case, this process has to go through several cycles before an acceptable harvest schedule can be implemented.

The spatial arrangement of areas for timber harvesting is especially challenging in this province because of the natural fragmentation of our forests. This model provided forest planners with the ability to mimic realistic timber harvest schedules based on current practices and to identify other forest stands that are not as accessible for harvesting.

Manual harvest scheduling has several major benefits. First, it fosters the long term sustainability of our AAC's by mimicking current harvest practices and accounting for actual on the ground conditions that delay or restrict the harvesting of stands. These restrictions, which were previously unaccounted for, have made our past AAC's higher than was realistically sustainable. Secondly, the mapped harvest schedules build credibility into the forest management process. A common misconception is that the province is running out of wood and soon will not be able to support existing forest industries. Every stand that will be harvested over the spatial plan must already be in the second (20-40 years old) or third (41-60) age class and can be easily identified and highlighted.

Having the ability to visualize the timber that will be harvested in the future helps reassure the resource is being used in a responsible manner. Next, harvest scheduling helps integrate the management of other forest resource values into timber management planning. Specific forest values can be directly related to forest areas, which can be mapped and potential issues can be addressed. Finally, the harvest schedule maps developed for the wood supply analysis can be a starting point for a 5 year operational planning process. Worthy to note is that harvest scheduling is completed for class 1 landbase only. The class 3 AAC, for the most part, is considered opportunistic if economic conditions become favorable.

3.4.4.3 Planning Horizons

Given the province's commitment to long term sustainability of our forest resource, timber supplies are projected 160 years (equivalent to two forest rotations) into the future to ensure actions and strategies applied today will result in a sustainable forest in the future. Long term planning is fundamental in timber supply forecasting.

3.4.4.4 Operable Growing Stock Buffer

The province imposed an operable growing stock constraint in the analysis to ensure the sustainability of calculated timber supplies. This constraint imposes a condition that in any period there must be a minimum operable growing stock of two times the harvest level on the landscape. In other words, for every hectare that is harvested another harvestable hectare must exist on the landscape. The requirement for a growing stock buffer is based on a number of factors including:

- 1. Some of the non-timber objectives are not explicitly accounted for in the planning process and therefore will require a growing stock buffer to achieve them.
- 2. The ability to completely incorporate the optimum harvest schedule due to operational restrictions on commercial harvesting.
- 3. Lowers the overall risk associated with the sustainability of the timber supply.

For these reasons a growing stock constraint of two times is utilized. This constraint is used in concert with harvest scheduling to help map out a reasonable harvest for the upcoming 20 years.

3.4.4.5 Old Forest Targets

Within the woodsupply analysis, the department considers a target for Old Growth forest, where at least 15 percent of forests at any given time must be be older than 80 years. While this is a minimum target, actual results are usually higher. This initiative was designed to provide a coarse filter approach to maintaining representative forest structure. It ensures the presence of certain amounts of old forest across the landscape into the future and can be tracked across a district.

3.4.4.6 Operability Limits

Operability limits are considered the timeframe in which forest harvesting activity can be undertaken within specific forest stands. Stand growth development (merchantable timber volume) and individual piece size of trees are factors which determine a stands readiness for commercial harvest activity. Some younger stands may have acceptable harvest volumes, but still contain trees that are too small (diameter and height) to commercially harvest. In the 2006 wood supply analysis both stand volume and tree size were used to determine the earliest age when a stand could be initially harvested. In addition to determining the absolute earliest age a stand can be harvested, it was recognized that not all stands on the same site develop exactly the at the same rate. A small portion of a stand will develop faster; a small portion will lag behind; with the bulk of the stand type representing the average condition. Therefore, the first operability limit was staggered by 5 year intervals with the 10 percent, 30 percent, and 60 percent assigned to each availability class listed above respectively. The ending operability limits or the last age in which a stand can be harvested before it becomes too old to harvest is solely determined on a minimum stand volume of 60 m3/ha, after which that stand is not considered to have enough volume to make it economical feasible to commercial harvest operations. It should be noted that while the operability limits define the extreme end points of when stands can be harvested, very few stands are ever harvested at these extreme points. In order to meet other non-timber objectives and maximize the total volume of wood harvested, the model schedules stands to harvest somewhere inside the operability limit window.

3.4.4.7 Silviculture

Silviculture is one of the main forest management tools available to forest managers when analyzing the many different future forests that are generated using the wood supply modelling software.

The silvicultural actions used in the current woodsupply analysis include:

- 1. Pre-commercial thinning of balsam fir, black spruce, and softwood hardwood stands,
 - 2. Full planting of any areas that do not regenerate naturally with either white spruce, black spruce, or Norway spruce, and
 - 3. Gap planting of stands with either black spruce or balsam fir seedlings. Gap plant is the filling in of "holes" within stands that have inadequate natural regeneration of either balsam fir or black spruce.

The thinning levels (ha) for districts 09 and 16 used in the analysis were 25, and 0 ha respectively. The planting levels (ha) for districts 09 and 16 used in the analysis were 100 and 75 ha respectively.

3.5 Inventory Adjustments

One of the limitations of the current wood supply model is its inability to account for volume depletions outside of what is reported for harvesting operations. The model produces a gross merchantable volume (GMV) figure which needs to be adjusted to account for volume losses as: fire, insects/disease, timber utilization practices and the presence of stand remnants. It was recognized that a need existed to study each component more intensely and the staff from the Forest Engineering and Industry Services Division, over a seven year period, completed an analysis of the individual components. The results of these (and other) analysis are incorporated into the current woodsupply analysis.

3.5.1 Fire

An estimate of productive area loss as a result of fire was based on an analysis of the historical fire statistics maintained by FFA

3.5.2 Insects

An aerial mortality survey was completed on areas with historically high insect infestations. This information along with a GIS analysis of areas salvaged enabled FFA to determine the amount of productive area lost to insect mortality each year.

3.5.3 Timber Utilization

Information for this adjustment was derived from a series of intensive on-the-ground surveys which measured the amount of wood remaining on cutovers following harvesting. This wood was comprised of solid merchantable wood (logging losses) and wood with inherent cull (butt/heart rot). Surveys were conducted province wide and on all tenures where the information is analyzed by harvesting system and season.

3.5.4 Stand Remnants

Following harvesting operations, small fragments of stands often are left for a variety of reasons (operational constraints, low volume stands, terrain conditions). These often result in the inability of the operator to achieve volumes predicted by the computer models. Surveys were conducted across the province and the results analyzed to determine the amount of productive area attributed to remnants. The total inventory adjustment for Forest Management Districts 09 and 16 was 18 percent.

3.6 Results

The results of the timber supply analysis for Forest Management District 02 for the period 2016-2020 is shown in Table 16. A new woodsupply is being developed for the same districts for the period 2022-2026.

	Annual Allowable Cut Volume (m ³)		
	Core Softwood	Operational Softwood	Domestic Softwood
DISTRICT 02	60,840	8,112	32,292
DISTRICT 03	0	0	13,391

Table 7: Annual Allowable Cut Zone 2 2016-2020

SECTION 4 VALUES

4.0 Guiding Principles of Sustainability

Environmental, Economic, Political, Social, and Cultural are considered the five guiding principles of sustainability.

Environmental sustainability evaluates current and future ecosystem health. It ensures the needs of the present are obtained without compromising the ability of future generation's needs. Ecosystem health is determined by such factors as ecosystem integrity, biodiversity, productive capacity, and resiliency. The five year operating plan strives to ensure these factors are maintained.

Economic sustainability requires forest resources to be managed efficiently and equitably among stakeholders. Economic development remains high priority for many of the residents within the Province. However; economic development should only proceed with the incorporation of the other principles of sustainability.

Political sustainability refers to goals and management objectives being applicable, administrable, and practical. With the aid of public input and support, these goals and objectives must maintain these qualities into the future.

Social sustainability means fairness and equity to all interested stakeholders. The forest management strategy should not jeopardize the basic requirements of the public. As a result, public involvement/awareness, participation, and decision-making are considered necessary to development of proper forest management plans.

Cultural sustainability is attained by applying Newfoundland and Labrador's culture to the planning process. A forest management strategy cannot be successful without allowances within the strategy for traditional access and use of the land. For generations, many of Newfoundland and Labrador's public has had free range in our pristine wilderness, a fact that cannot be ignored when planning for the zone. All are key interlocking components and each must be maintained if sustainable development is to be properly achieved.

4.1 Value Structure

The forest ecosystems of the zone provide a wide range of values to different individuals and groups, which include:

- Consumptive values such as: timber products, hunting, trapping, sport fishing, and berry picking.
- Non-Consumptive values such as: skiing, snowmobiling, hiking, and bird watching.
- Intrinsic and intangible values such as a feeling of wilderness and peace which some people describe as spiritual. Although difficult to spatially describe or quantitatively measure, spiritual values are considered to be a product or an accumulation of all values.

Other values such as water quality, parks and protected areas provide the protection of forest ecosystems, which can enhance the above identified values. Many of the values in the zone are identified by many years of forest management planning and engagement with interested stakeholders. The following represents a framework for characterizing values in a clear and consistent manner. This approach consists of three components:

VALUE STRUCTURE		
Characterization	Description: Why the value is important, types of activities, intensity, spatial extent, employment, etc. Data in support: Statistical references	
Critical Elements	Forest Features: Elements at risk from harvesting or enhanced by harvesting (viewscapes, adjacency to water, mountains, habitat, wilderness ambiance, road Access, etc.)	
Guiding Principles	 A guiding principle can be defined as a fixed or predetermined policy or mode of action. These 'modes of action' would be implemented in the five year plan in the form of: Policies that should be in place to protect or enhance the resource value; Methods for negotiation or inclusion of other interested stakeholders in resolving potential conflicts; Special management provisions/strategies such as: riparian buffer zone consideration, temporal operating periods, modified harvesting, or best management practices, and/or Models and/or forecasting strategies to determine economic contribution, biodiversity impact, or community sustainability 	

In many instances, the Environmental Protection Guidelines (EPG's) developed by the department help form the guiding principles for a value. Quite often the spatial extent or location of all values is not known (eg., raptor nests). Specific guidelines are still listed in order to provide a direction or course of action when and if these values are encountered.

4.1.1 Biotic Values

4.1.1.1 Big Game

4.1.1.1.1 Moose

MOOSE				
Characterization	• Moose were introduced in 1878 (Gander Bay) and 1904 (Howley) and are now distributed throughout the entire Island.			
	• The 2020 population estimate is 118,000 animals.			
	 The Province is divided into Moose Management Areas (MMA); boundaries, season dates and license quotas are reviewed annually for each MMA 			
	• Approx. 100,000 residents actively participate in the annual moose license application process. If successful, it provides an opportunity to hunt, to take part in an outdoor fall activity, to be part of an important social event and to provide local food.			
	• Non-resident harvest is conducted through outfitting establishments, employing guides, cooks, maintenance people, pilots, etc. Other businesses benefiting from the non-resident outfitting industry include local stores, accommodations and restaurants, outdoor stores, aviation companies, etc.			
	• Moose are also important for non-consumptive activities. Intrinsic value is added to hiking, cycling, camping, paddling, sightseeing, etc., when there is a chance to view megafauna such as moose. The tourism industry relies on this intrinsic value to promote many of the activities offered.			
Critical	Habitat			
Elements	 Moose select different habitats at different times of the year and therefore require a mixed forest landscape year-round: a mixture of closed, mature stands for cover, shelter and winter moose yards; early- seral stages (due to disturbances such as insects, fires and logging) with a high amount of young regenerating areas for browsing; and aquatic foraging areas in the summer. 			
	Forest harvesting			
	 Recently harvested large scale areas can be low quality moose habitat as forage and cover are removed. 5+ year old cutovers tend to have enough regeneration to support a slow return of moose to the area. Regenerating cutovers provide excellent foraging opportunity until trees become too tall for moose to reach. 			
	Forest roads			
	 Provide hunters access into areas previously free from vehicle traffic. Removal of access, in particular in areas of new forest regeneration, may allow moose densities to increase beyond sustainable levels. 			
Guiding Principles	 Proposed forestry activity is reviewed by the staff at the Wildlife Division and recommendations are incorporated into this five-year plan. 			

4.1.1.1.2 Caribou

	CARIBOU
Characterization	 Caribou in Canada occur in two primary ecotypes: Barren-ground caribou and Woodland caribou, often referred to as boreal caribou. The latter ecotype inhabits mosaics of barren, wetlands, and mature forests. The herds on the island of Newfoundland belong to this ecotype. They are cyclic by nature mainly due to being density-dependent; meaning restricted by food and/or space, as well as regulated by predators such as wolves and managed hunters.
	 Caribou is the only native ungulate species on the island. Numbers are anecdotal at best for the 1800's (ranged from 40,000 to 200,000). By 1925 hunting was closed as the island population was down to a few thousand caribou. From the 1960's through the 1990's the caribou population recovered, peaking at over 90,000 in the 1990's. In late 1990's caribou started another downward trend (2014 assessment: just over 32,000 animals). The 2019 population estimates suggests 30,600 animals island-wide.
	• The Province is divided into Caribou Management Areas (CMA): boundaries, season dates and license quotas are reviewed annually for each CMA.
	• Approx. 100,000 residents actively participate in the annual big game license draw process. If successful, it provides an opportunity to hunt, to take part in an outdoor fall activity, to be part of an important social event and to provide local food.
	 Non-resident harvest is conducted through outfitting establishments, employing guides, cooks, maintenance people, pilots, etc. Other businesses benefiting from the non-resident outfitting industry include local stores, accommodations and restaurants, outdoor stores, aviation companies, etc.
	 Caribou are also important for none-consumptive activities. Intrinsic value is added to hiking, cycling, camping, paddling, sightseeing, etc., when there is a chance to view megafauna such as caribou. The tourism industry relies on this intrinsic value to promote many of the activities offered. Caribou is also iconic to the NL Regiment.
	 In 2014 the Committee on the Status of Endangered Wildlife in Canada (COSEWIC) assessed Newfoundland caribou and recommended a listing of "special concern". COSEWIC assessments are usually done on a 10 year cycle.
Critical	• Caribou display a highly mobile nature and some herds occupy large home ranges.
Elements	• Caribou show a high selection for bogs and mature coniferous forests, as well as shrubs and water bodies. These habitats provide refuge from predation and support an abundance of forage.
	• Direct impacts from large cut overs and linear features include habitat destruction and fragmentation, increase in mortality and reduced recruitment rates likely due to increased and improved access for humans as well as predators, and avoidance of disturbed areas

Guiding Principles	 Caribou populations are monitored and assessed on a regular basis via Annual fall classification surveys: provide regular updates to sex ratios and recruitment within each herd. This work feeds into herd level assessments and quota evaluation/development.
	 Collar deployment: analysis of caribou movements, mortality rates and range occupation
	• This supports the development and assessment of Caribou Management Guidelines for forest management planning. New iterations of these guidelines may incorporate maximum targets for disturbance and linear features (e.g., forest roads) as well as no net losses within certain caribou core ranges.
	• The guiding principles for forestry operations and road construction will apply to all crown operations within the province. Wildlife Division will provide herd specific guidance and recommendations.

4.1.1.1.3 Black Bear

BLACK BEAR	
Characterization	• The black bear is native to the island and is found in forested areas. Black bears are solitary creatures and do not truly hibernate but enter a state of lethargic sleep. A den is chosen in mid fall under tree stumps or logs which are lined with grass, twigs and leaves. If reproducing females gained enough body fat they will give birth between January and February before emerging from the den with the cub(s) in April or May.
	• Currently, the number of black bears occurring on the island has been estimated at approximately 6,000 - 10,000 animals.
	• The Province is divided into Black Bear Management Areas (BMA), which correspond to MMAs (except for sub-areas). Currently, only one license (with a bag limit at two black bears) is required by hunters on the Island to hunt black bear in both the spring and fall.
Critical Elements	 Heavily wooded areas with dense bush intercepted with smaller clearings or early successional seral stages represent suitable habitat for black bears.
Guiding Principles	• A 50-metre undisturbed buffer must be maintained around known bear den sites (winter) or those encountered during harvesting.

4.1.1.2 Furbearers

FURBEARERS	
Characterization	• There are a variety of furbearers within the Province: lynx, red fox, beaver, otter, muskrat, short-tailed weasel, red squirrel, mink, coyote, wolf (Labrador) and

Critical	 Newfoundland marten (currently listed as a Species at Risk). Of these, mink and red squirrel are not native to the Island and coyotes are considered recent colonizers as they expanded their range naturally. Furbearer populations fluctuate from year to year and are influenced by factors such as predator/prey relationships, food supply and disease. Trapping pressure (i.e. due to fur market prices) can also influence population growth or decline. Trapping furbearers for their fur, meat and other natural products has a long tradition in this Province. Today's trapping is a choice of lifestyle, most times deeply rooted in the family and local culture, provides food, clothing and money as well as an enduring connection and experience with nature. Approx. 2,500 people in the province trap and snare furbearers. Management of furbearer species within the province is highly regulated and the responsibility of the Wildlife Division. There are 13 furbearer management zones throughout Newfoundland and Labrador. Within these zones, harvest activities for each species occurs during optimum periods for fur quality, which is generally during the fall and winter period. However, furbearer seasons may vary for each different species.
Critical Elements	• Embedded in conservation of furbearers is the conservation of biodiversity and the ecological processes that support that biodiversity. Furbearers depend on other species; lynx on snowshoe hares, marten on mice and beaver on deciduous trees. Each species belong to a diverse community that depends on specific habitat conditions. Habitats change from one forest successional stage to another and are therefore inextricably linked to forest management. (Fur Institute of Canada, 2019)
	 Species respond differently to logging operations or Silviculture treatments (i.e. thinning, prescribed burns). Hare and grouse increase with logging, which is advantageous to lynx, coyote and fox. The wolf depends on caribou and moose, and beaver.
	 Snags, coarse woody debris, and sufficient understory provide sites for shelter, denning, nesting, traveling, etc.
	 Maintaining sufficient undisturbed riparian buffer zones along aquatic areas such as wetlands and waterbodies ensures water quality maintenance as well as shelter from upland disturbances
	• A mix of various forest conditions and successional stages that provides diversity of habitats supports a diversity of prey and thus a diversity of furbearers.
Guiding Principles	• The Wildlife Division develops and implements an annual <i>Furbearer Management Strategy</i> . Similar to the big game management plan, a furbearer management plan reviews the status of each furbearer species, addresses the season dates and lengths, and if necessary closure of areas (or no open season as currently in place for marten).

4.1.1.3 Species of Interest

4.1.1.3.1 American Marten

	AMERICAN MARTEN
Characterization	In 1986, the Committee on the Status of Endangered Wildlife in Canada (COSEWIC) assessed the Newfoundland population of the american marten and the species was listed as threatened. Revisions in 1996 and 2000 resulted in an uplisting to "endangered" due to further declines. Habitat loss, trapping and incidental snaring are possible reasons for the marten population decline. Through the work of the Marten Recovery Team, the status of marten has been upgraded from "endangered" to "threatened" in 2007 because new population estimates were stable and distribution of marten was increasing. The American marten (island population) is currently (2010) listed as "threatened" under both the federal <i>Species at Risk Act</i> and provincial <i>Endangered Species Act</i> .
	To identify factors affecting marten survival, stakeholders from the Canadian Forest Service, Wildlife Division, Corner Brook Pulp and Paper and the Forestry Branch became members of the Newfoundland Marten Recovery Team. The primary function of the Recovery Team was to prepare and periodically revise the recovery plan for American marten in Newfoundland and to provide advice on species recovery. The initiation of the live-trapping program, revealed that Main River, Little Grand Lake and Red-Indian Lake are high-density marten areas on the island. Based on this information, it is important that marten habitat be protected in these areas. Furthermore, it is important that some remnant stands of old growth (80+) forests remain on harvested areas throughout the province and provision made to have connectivity (i.e., unbroken corridors of forest).
Critical Elements	 Older studies of habitat associations of American marten in Newfoundland have found that they are dependent on old-growth or overmature (>80-years-old) forests due to a depauperate prey base which restricts marten to older forests where prey are accessible. More recent research suggests that marten will use a wide variety of habitat types, including immature regenerating forests, precommercially thinned forests, areas disturbed by forest insects, and areas of mature and overmature forests sufficient habitat to support a viable population of marten; Overall experts agreed that forest management promotes marten populations when some remnant stands of over mature/ old-growth (80+) forests remain unharvested across the island and a provision made to have connectivity between these over mature or old growth stand with the habitats that are listed above in the first bullet (this does not include recently harvested areas). Under Brian Hearn's work with the Canadian Forest Service, previous proposed
	harvest schedules within various forest management districts have been analyzed and indicate suitable habitat remained after harvest.
Guiding Principles	These guiding principles are put in place to further recovery of the Newfoundland Marten and allow for forest harvesting.
	Sustainably manage and conserve core marten areas:

In the early 1990's-2000's there were three high-density marten areas on the island: Main River (southern Northern Peninsula), Little Grand Lake- Red-Indian Lake (Western) and Terra Nova (Eastern). Marten have and continue to expand their range from these core areas. Based on this information, it is important that marten habitat be managed with a focus on the sustainable use and conservation of these core areas and the management of forests to promote and continue range expansions beyond the three core areas.
Ensure habitat connectivity:
Forest planning ensure sufficient connectivity between habitat patches to allow continued dispersal and expansion of the species back to its historical range.
Protect denning female and kits:
Activities previously identified by the Newfoundland Marten Recovery Team, as having the potential to disrupt female marten and her kits during the critical denning period, where possible should be limited during the period April 1st to June 30th within the areas identified as marten critical habitat in the 2010 Provincial Recovery Plan. This includes any activity that results in the removal of trees, significant noise, or compression or disturbance to brush or soil. Wildlife Division must be consulted on activities proposed in critical habitat during this period.
• The guiding principles for Habitat requirements during forestry operations will apply to all crown operations within the province.

4.1.1.3.2 Rare Plants

RARE PLANTS	
Characterization	
	 The distribution of specific species as well as plant communities depend on regional and landscape-level factors, such as climate, geology, altitude and terrain as well as more local factors, such as the depth, moisture regime and fertility of the soil, microclimate and other biota, such as presence of herbivores, pathogens or pollinators.
	 Information on the distribution of plant species in Newfoundland and Labrador is obtained from various surveys initiated by government, research institutions, individual naturalists or local organizations, as well as those by consulting firms required through Environmental Assessment processes.
	• Currently, 27 plant and 3lichen species are listed under the <i>Endangered Species Act</i> , with several hundred other plant and lichen species considered to be of conservation concern.

	 Four of the Endangered plant species are not found anywhere in the world outside of the Gulf of St. Lawrence Region, and three of them, the Fernald's Braya, Long's Braya and Barrens willow, are only found on the Island of Newfoundland. There are many species of plants, lichen and fungi throughout the Province that are not rare or endangered but still have an important social or economic value. Many of these species also play important roles in the functioning of the province's ecosystems. The vast majority of the rare plant species throughout Newfoundland are inhabitants of open habitats, such as river gravels, salt marshes, wetlands, aquatic habitats, alpine areas and and coastal barrens, which are not targeted by commercial forest operations and in some cases receive special protection
Critical Elements	 A small subset of the plant and lichen species of conservation concern, such as the Endangered Graceful Felt Lichen, depends on forests, and some of them are specialists requiring late-successional forests. Without proper protection measures, negative effects can occur from: quarrying and road construction logging and extraction using heavy equipment mechanical site preparation all terrain vehicle traffic prescribed burning stand conversion to different tree species and/or earlier seral stages
Guiding Principles	 Work with the Wildlife Division to educate forestry Personnel and develop mitigative measures in areas where rare plants occur. Identify and update all rare plant sites on GIS forestry data base Encourage domestic harvesting in the winter The guiding principles for Protection during forestry operations will apply to all crown operations within the province.

4.1.1.3.3 Waterfowl

WATERFOWL	
Characterization	Maintaining waterfowl populations depends on effective conservation efforts
	throughout North America. Newfoundland and Labrador most productive waterfowl
	breeding and staging areas are either owned by or are under management control of

	major corporations, private or communities. To help manage for waterfowl areas, stewardship agreements are signed at either the municipal, corporate or private levels.
Critical Elements	 Habitat maintenance of habitat Breading disturbance of waterfowl during the brood rearing, breeding, and staging period
Guiding Principles	 Habitat 50-metre treed buffer will be established around designated sensitive waterfowl areas. Breeding A minimum 30-metre, treed buffer must be maintained from the high water mark in other waterfowl breeding, molting, and staging areas. no forestry activities are recommended during the brood rearing, breeding, and staging period The guiding principles for Habitat and Breeding during forestry operations will apply to all crown operations within the province.

4.1.1.3.4 Other Species

Other species, particularly the red crossbill, are currently listed as endangered. The Forestry and Wildlife Division has a representatives on the recovery team for this species. Any recommendations on modified forestry activities, if any, for this species will be developed with input from all members and implemented accordingly.

4.1.1.4 Water Resources

WATER RESOURCES	
Characterization	The protection of water resources is an important topic both nationally and provincially. Human impacts both locally and globally have the potential to impair or alter water quality and water quantity.
Critical Elements	 Road construction / maintenance, Timber harvesting, and Silviculture Potential to alter the quantity and quality of water draining from watersheds. Negative Impacts could include: Negative impacts could include: stream hydrology (e.g. water quantity/flood risk),sediment loadings, and stream characteristics. Such activity would also potentially increase access by the public to land inside a Protected Public Water Supply Area (PPWSA) that was not previously accessible.

	 Storage and handling of fuels Public Protected Watersupply Areas (PPWSA) Human Activity (e.g. domestic and commercial forestry has the potential to negatively affect the area of land and water designated as a Protected Public Water Supply Area, for a municipal authority or local service district (LSD) operating a waterworks or using or intending to use a water sources, under Section 39 of the Water Resources Act. 3 Working in and within 15 metres of a waterbody or wetland The management of water resources helps to protect, enhance, conserve, develop and effectively utilize NL water resources and is accomplished through the issuance of Permits to Alter a Body of Water and Water Use Licences under Section 48 and Part 1 of the Water Resources Act respectively.
Guiding Principles	 Road Construction Any alteration to a body of water (e.g. culvert or bridge installation) within 15 metres of a natural waterbody (i.e. any water identified on the latest 1:50,000 NTS map) or development within a protected public water supply area, will require prior approval by the Water Resources Management Division of the Department of Environment and Climate Change (ECC). Approvals from Transport Canada are required for culverts, bridges and abutments on navigable waters (i.e. any waterbody capable of being navigated by floating vessels of any description for the purpose of transportation, commerce or recreation. Transport Canada's Navigability Self-Assessment Tree) must be utilized for each project to determine if a stream is Navigable or Not Navigable. When extraction trails and winter roads are to be constructed, soil disturbance and impacts on water bodies are to be minimized.
	 Timber Harvesting Heavy equipment and machinery are not permitted in any waterbody, on a wetland or a bog, unless frozen, without a permit from Water Resources Management Division Woody material of any kind (i.e. trees, slash, sawdust, slabs, etc.) is not permitted to enter a waterbody. Depositing woody material on ice within the high water floodplain of any waterbody is also prohibited. Buffers inside PPWSAs vary and are applied using Policy 95-01 (Policy for Land and Water Related Developments in Protected Public Water Supply Areas) Extraction trails and landings shall not be established within 30 metres of a waterbody A minimum 30 metre, no harvesting activity buffer zone shall be established around all water bodies that are identified on the latest 1:50,000 national topographic system (NTS) maps. All buffers may be increased inside PPWSAs. Where possible, a closure plan will be identified at the end of Forestry activity to minimize access.

 Storage and handling of fuels No heavy equipment or machinery is to be refueled, serviced, or washed within 30 metres of a waterbody. Gasoline or lubricant depots must be placed at least 100 metres from the nearest waterbody All fuel-storage tanks must be registered with Digital Government and Service NL Used oil storage, handling and disposal is to comply with the Used Oil Control Regulations, NLR, 82/02 under the Environmental Protection Act. Contaminated soil or snow must be disposed of at an approved treatment facility
 Protected Public Watersupply Areas (PPWSA) In Newfoundland and Labrador forestry operations may be permitted in Protected Public Water Supply Areas on a limited and controlled basis provided the potential negative impacts of the proposed operations can be mitigated. Approvals required include:
 Working in and within 15 metres of a waterbody All waterbodies, including wetlands will be identified within the project area as per the most up to date 1:50,000 NTS topographic maps. Work within mapped wetland areas will follow the WRMD's Policy for Development in Wetlands to ensure that adverse effects to water quantity, quality, hydrological functions and terrestrial and aquatic habitats are restricted. All necessary erosion/siltation control measures will be taken during infilling, dredging and debris removal activities near waterbodies.

 Impacts to the natural drainage pattern of the work area shall be identified including the redirection and discharge of water. Flood risk areas will be identified and work activities categorized as per the WRMD Policy for Flood Plain Management to ensure structural, public, and environmental health and safety. Required permits and licenses will be attained prior to commencing work include; Issuance of a Permit to Alter a Body of Water under Section 48 of the Water Resources Act Issuance of a Part 1 – Water Use Licence for all non-domestic water use.
 The guiding principles for forestry operations, road construction, fuel handling and operating with PPWSA's will apply to all crown operations within the province.

4.1.2 Human Values

4.1.2.1 Timber Resource

TIMBER RESOURCE	
Characterization	Commercial logging contractors are allocated the majority of the annual allowable cut level in the zone. Commercial harvesting and sawmilling activity provides many jobs in harvesting, sawmilling, trucking, pulp and paper manufacturing and related spin off industries for local residents. Commercial uses have arisen for timber, which includes: lumber, pulp and paper products, and value added products. Domestic harvesting provides fuelwood to heat many homes and sawlog material for residential house construction in the zone. Domestic harvesting is conducted in specific domestic cutting areas via a crown domestic cutting permit that is required and issued within each forest management district. Unless otherwise specified; domestic cutting is limited to these designated cutting areas. A Domestic Permit specifies the volume and species that can be harvested, utilization standards, and other relevant conditions. While some domestic cutting areas are designated for hardwood only, the majority of areas will allow the harvest of all hardwood and softwood species.
	Silviculture treatments are important to the forest resource of the zone as it ensures that a vigorous and healthy forest is maintained. Forest renewal activities facilitate renewal of productive landbase by manual planting areas that are not sufficiently restocked. Forest improvement activities help improve and enhance the growing stock which can reduce harvest cost, enhance forest product options and increase sustainable timber supply. There will be a significant investment on silviculture in the zone each year creating seasonal employment. Timely access to timber is critical to planning any forestry operations. Primary, secondary and tertiary roads form an integral part of operating areas for commercial harvesting

	 activity. Upon completion, these roads are often used for silviculture and recreational purposes. A significant amount of money will be spent to construct forest access roads each year in the zone. Protection of the forest from various disturbances is a major characteristic of resource management, which includes: integrated pest management and forest fire prevention/suppression techniques. Other resource values are protected through modification of activities and enforcement.
Critical Elements	 maintenance or enhancement of productive land base planting of non-regenerating areas minimizing loss of land base to other users minimize losses to fire, insect and disease timely access road construction enhancement of younger age classes through thinning to correct age class imbalance
Guiding Principles	 enforcement of forestry act, regulations, guidelines and policies minimize loss of productive land base through spatial and temporal compromises and continuous dialogue with other resource users education (staff, public, operators) aggressively conduct silviculture, access road, and protection activities implement best management practices. The <i>EPG's</i> outline courses of action and mitigative measures for forest activities. These EPG's are outlined in their entirety in Appendix, with some highlighted subject areas listed below. garbage disposal fuel storage mineral soil exposure buffer requirements road and bridge construction silviculture and harvesting activities

4.1.2.2 Agriculture

AGRICULTURE	
Characterization	The agricultural industry provides both direct and indirect employment to residents of
	the province.
Critical	Landbase
Elements	 Land resources are at the foundation of all agricultural enterprises and provide the requirements for crop production. It is not possible to identify and plan all sites for future agriculture use and often there is a conflict with other land uses particularly forestry because these sites are of high growing capability.

	 Although a suitable landbase is the first critical element necessary for a successful agriculture operation, markets and the interest of individuals are also prime factors in the development and location of future farms. Operational Requirements When approval is granted for an agricultural lease, the land needs to be cleared and prepared for production
Guiding Principles	 Landbase Both the Forestry and Agrifoods sections have identified Agricultural Areas of Interest (AOI's) across the province, representing approximately 155,600 hectares. The area in AOI's have been removed from woodsupply calculations and is available for agricultural development. Agricultural Development is still considered for areas falling outside established AOI's Operational Requirements Forestry Act has been amended to allow clearing of land to occur without a commercial cutting permit if applicant is utilizing the fibre for their own private use. Where possible, existing commercial forest operators should be encouraged to work with farmers to clear new land for development. Where possible, existing commercial forest operators are encouraged to clear identified land within the departments Agriculutre areas of Interest (AOI's). Five Year Plans includes AOI areas which are not part of woodsupply calculations Home gardening leases should be confined to areas already developed for this activity. The guiding principles for Landbase and operational requirements will apply to all crown operations within the province.

4.1.2.3 Mining, Mineral Exploration and Quarrying

MINING, Mineral Exploration, and Quarrying	
Characterization	Mineral exploration, mining, and quarrying are recognized as separate and distinct activities, each of which is approved and regulated under a separate piece of legislation.
	Mineral exploration activities may consist of prospecting, geological mapping, grid line- cutting, geochemical surveys, ground-based and airborne geophysical surveys, the preparation and use of access trails, mechanized trenching, diamond drilling, and – in remote areas – the preparation and use of campsites.
	Mineral exploration takes place province-wide and is a significant contributor to the provincial economy, particularly in rural areas.

	There are a large number of active quarries throughout the province which generate significant royalties and which provide the raw material for the development and maintenance of infrastructure such as highways, building lots, and concrete.
	There are a number of active mines in the province at any given time and mining represents a major component of the provincial economy.
Critical Elements	 Forest Ecosystem Mining, quarrying, and mineral exploration activities can have potential negative impacts to Forest Ecosystems and future Woodsupply calculations. Mining and quarrying represent permanent (but sequenced – not all at once) alterations to the landscape whereas mineral exploration activities at most involve temporary disturbance. Each activity is subject to rehabilitation requirements. Utilization of Timber Resource
	O When exploration activity occurs, merchantable trees may need to be harvested to gain access to work sites. Under forestry act, all merchantable trees can only be removed with a cutting permit and that holders of a cutting permit must utilize all portions of the tree to a top diameter of 8cm (outside bark). Section 18 of cutting of timber regulations state that all timber cut shall be removed from the cutting area to a roadway while harvesting operations are in progress unless otherwise specified in the cutting permit.
Guiding Principles	 Forest Ecosystem Regulations and permitting conditions require that all mine sites, quarry sites, and mineral exploration sites be rehabilitated. While each activity is regulated separately, rehabilitation requirements for each include the requirement that the organic overburden (e.g., topsoil, ground vegetation) be stockpiled and stored in a manner so that it can be used to rehabilitate the site. Rehabilitation requirements can be further specified in special terms and conditions and could include, if warranted, the requirement that the site be left in a condition conducive to forest regrowth. Utilization of Timber Resource As part of exploration and development activities, individuals must attempt to extract timber harvested. If timber cannot be feasibly extracted using conventional means, then timber shall be piled so that it may be extracted during winter months by snowmobiles. Mineral Exploration companies are permitted to use some harvested timber within access routes for the purposes of corduroy or brushmatting to prevent rutting and minimize ground disturbance on sensitive sites.
	 Additional: The Forestry Branch will consult with the Mining and Mineral Development Branch in determining appropriate silviculture buffer distances from the boundaries of sites covered by a quarry permit or quarry lease. In many cases, 100 meters is an appropriate buffer distance that will accommodate

 discrepancies in plotting. However, in other cases, there may be valuable aggregate resources present and expectations that the area will see proposals for new quarry developments in the foreseeable future, in which case a buffer zone should be specially planned in consultation with Mines. In other cases, where quarry development is expected to proceed slowly, silviculture buffer distances may be much reduced. For mine sites (including all associated infrastructure), an appropriate silviculture buffer distance should be chosen in consultation with the site operator and the Mining and Mineral Development Branch. Mineral exploration activity that proposes to explore or develop within a silviculturally treated area must be undertaken with minimal disturbance and under approval of Forestry Branch. A standard condition has been developed by the Mineral Development Branch and Forestry Services Branch to include as a condition in mineral exploration approvals document where silviculture treatments may be impacted. Mineral exploration and/or development on mineral licenses will not be impeded and will follow government policy. Specific proposed forest management activities are identified in annual operating plans for each upcoming calendar year. Should future quarry or mineral resource developments or exploration programs (i.e., new quarry development, existing quarry expansion, new mine development, exploration for quarry materials, or mineral exploration) be considered by the Forestry Services Branch as having the potential to cause a significant impact on the forest resource and forest mutual impacts are minimized. For the purpose of road construction, quarry permits or quarry leases are required only for aggregate material taken outside of the road right-of-way. Non-compliance with exploration permits identified by Forestry Sarncies arant mutual impacts are minimized. For the purpose of road construction, quarry permits or decommission roads or

4.1.2.4 Historic Resources

HISTORIC RESOURCES	
Characterization	The provincial archeology office (PAO) is the agency responsible for the management and protection of archaeological sites and artifacts in Newfoundland and Labrador. This program is carried out under the Historic Resources Act which ensures that developments with potential to have adverse impacts on historic resources are investigated as and monitored by a qualified archaeologist through archaeological impact assessments. Archaeological sites are non-renewable resources and play a vital role in understanding our heritage. Most often, archaeological sites are small in size, so it is important to protect these sites and professionally record as much information possible to fully understand its history. To do this properly, the site must not be disturbed.
Critical Elements	 Protection Activities which disturb soil layers and/or provide unintended public access to an archaeological site can have a negative impact on that historic resource. Without applying best management practices, forestry activities such as: construction of access roads and bridges, harvesting, and mechanical site preparation have the potential to destroy historic resources. While forestry activities can have adverse impacts on historic resources, beneficial effects can be realized. Where impact assessments are carried out and new sites found, it adds to our understanding of Newfoundland and Labrador's heritage. When archaeological sites are discovered through impact assessments, these resources are protected from damage or destruction and preserved.
Guiding Principles	 Protection the provincial archaeology office will review forestry's five year plan proposals through the Environmental Assessment process. Applicable protection measures will be identified by the archaeology department and incorporated into the forestry plans. Buffer Zones will be implemented to protect known archaeological sites and potential unknown sites. If deemed necessary, archaeological assessments may be required to fully assess the site. Buffer zones are required along all rivers and ponds, as well as along the coastline, where there is potential for archaeological resources to be found. Occasionally there are accidental discoveries made of historic resources. In the event that this does happen, activities should cease in this area and contact be made immediately with the provincial archaeologists. The guiding principles for Protection during forest Harvesting will apply to all crown operations within the province.

4.1.2.5 Newfoundland T'Railway

NÉWFOUNDLAND T'RAILWAY	
Characterization	The T'Railway is a linear park approximately 850 km in length, comprising the former CNR right of way, which varies from 25 to 100 feet each side of the center line. It provides for an all season, multi-use recreation corridor and is developed and managed with Parks and Natural Areas Division in conjunction with the T'Railway Council. The <i>Provincial Parks Act</i> provides the legislative framework for the administration and management of the T'Railway and is protected for present and future enjoyment of the public. The T'Railway constitutes the province's contribution to the Trans Canada Trail System and is used primarily by snowmobile and all-terrain vehicles. Industrial or traditional uses such as: commercial and domestic harvesting, quarry and mining access and cabin access are also granted with a special permit.
Critical Elements	 Protection protection of the historical landscape integrity of the T'Railway corridor preservation of the scenic quality along the corridor Operational Regulating land usage adjacent to the T'Railway
Guiding Principles	 Protection co-ordination of activities with various other agencies responsible for land management outside the T'Railway corridor to ensure that the integrity of the park is maintained build partnerships with other stakeholders and user groups such as communities, industry and recreational organizations for the long term maintenance and development of the T'Railway Establishment of a 100 meter buffer along the right-of-way corridor to preserve the natural value of the T'Railway. Also, consider viewscapes in forestry management plans. where access is required, any landings or turnaround areas shall be 100 meters or more along the resource roads from the T'Railway. Operational where feasible and possible, Forestry Activity utilizing the T'Railway will avoid peak snowmobile and ATV seasons. Applicable permits will be obtained for use of vehicular and heavy equipment on the T'railway The guiding principles for Protection and operational requirements will apply to all crown operations within the province.

4.1.2.6 Parks and Protected Areas

	PARKS AND PROTECTED AREAS
Characterization	 PARKS AND PROTECTED AREAS There are several different types of conservation areas in the province contributing to the provincial system of protected areas. Wilderness Reserves are designed to protect complete ecological systems, while Ecological Reserves may be established to protect representative samples of each of the province's natural regions. Both Wilderness and Ecological Reserves are established via the Wilderness and Ecological Reserves are established via the Wilderness and Ecological Reserves Act. Provincial Parks established under the Provincial Parks Act, do play a conservation role, but are primarily established as sites for outdoor recreation and nature-based education. Wildlife Reserves may be established under the Wildlife Act for the protection of specific species or habitats. Public or Crown Reserves may be established for conservation reasons under the Lands Act. National Parks such as Terra Nova, Gros Morne and Torngat Mountains are established under the federal National Parks Act.
Critical Elements	 research, opportunities for environmental education, provide standards against which the effects of development can be measured, and provide natural venues for enjoyment of nature. Protection preservation of biodiversity maintenance of protected area integrity maintain natural processes and features Operational Forestry activity occurring around Parks and Protected Areas.
Guiding Principles	 Protection Some protected areas prohibit new development such as mining activity, hydroelectric projects, forestry activity, agriculture activity, roads/trails, cabins and new structures; the type of activities encouraged or permitted within various protected areas in the province depends entirely on the type of protected area and the rational for its establishment Operational a 500 m no roads buffer is to be maintained around all existing and proposed protected areas to reduce access and minimize damage from motorized vehicles

\circ where forestry operations are scheduled within one kilometre of
provisional and ecological reserves, wilderness reserves or provincial parks, modified operations may be necessary
• The guiding principles for Protection and operational requirements will apply to all crown operations within the province.

4.1.2.7 Outfitting

	OUTFITTING			
Characterization	utfitting is an iconic, high yield tourism demand generator and one of Newfoundland and Labrador's most developed tourism products. Outfitters annually attract high end unting and angling enthusiasts that benefit local communities, other private operators, ansportation providers and guides. Ince the early 1900's, the outfitting industry has been an integral component of the purism industry in Newfoundland and Labrador. The province has been a popular unting and fishing destination because of the pristine environment and abundance of sh and wildlife species. There are many outfitters operating within the boundaries of is forest management zone, which provide seasonal employment for many local dividuals. Wer the past decade or so, a significant number of traditional hunting and fishing cilities have diversified into the non-consumptive areas of the tourism industry. Such strivities include: snowmobiling, dog sledding, kayaking, canoeing, nature viewing, king, and wildlife photography. The ability to diversify has positively impacted the ability of outfitting operations and as a result, increasing numbers of operators are posidering this opportunities. Pristine wilderness settings are necessary for many of			
	viability of outfitting operations and as a result, increasing numbers of operators are considering this opportunities. Pristine wilderness settings are necessary for many of these types of diversification.			
Critical Elements	 Resource Roads Some outfitting camps are considered remote and construction of forest access roads too closely to a main camp could have negative impacts to this remote appeal. Increasing accessibility through increased access roads has the potential for increased hunting and fishing pressures within in a given area. Increased pressure on the wildlife resource within a given area could potentially lead to decreased success rates of outfitter guests. With improved road access, there is also a potential for increased cottage development, which can also impact both remoteness and wildlife availability. 			
	 Visual While clients of hunting and fishing outfitters are primarily interested in the actual hunting or fishing experiences, they also show a great respect and admiration for pristine conditions and a healthy looking landscape. Activities such as forest harvesting can potentially detract from the overall visitor experience. 			

	 The landscape view experienced by clients plays a large role in leaving a lasting impression of the province. The viewscape experience may also have a direct impact on repeat client bookings and recommending the destination to others. Viewscapes become even more important as outfitters begin diversification into non-consumptive tourism activities. Wildlife Without proper application of best management practices, forest
	 harvesting has the ability to negatively impact wildlife travel corridors, bear denning areas, and moose and caribou feeding and calving areas. Removal of large areas of forest can simulate the same effect of reducing wildlife habitat, particularly winter staging areas. Forest harvesting has the ability to lead to disturbances of wildlife, including shifts in where they live, disturbances for hunters accessing hunting areas
	and finding animals in these traditional hunting areas, and other disturbances for outfitters. This has the potential to significantly impact their business.
Guiding Principles	 Resource Roads Where possible, new access roads will be strategically located away from an outfitters existing main outfitting Lodge. Consideration will be given to decommissioning roads and bridges (where possible) after harvesting is completed, which will help reduce possibilities of increased hunting pressure. When roads are in use actively for harvesting purposes, access to hunters could be restricted or limited. Where possible and feasible, conduct harvest activity in the winter and construct winter roads that are less passable in summer/fall and will help to reduce vehicular traffic around an outfitters main lodge. Visual In consultation with the outfitter, efforts will be made to minimize negative effects regarding the view from the outfitter's main lodge and their hunting areas when conducting nearby forest operations. Environmental Management System (EMS), developed procedures for: Appropriate stream and wildlife buffers are implemented and management and wildlife buffers are implemented and management
	 maintained Wildlife All forestry Plans are reviewed by the Wildlife Branch. Travel Corridors and Buffer Requirements are designed and implemented upon direction of Wildlife Branch using scientific data. Modified or deferred harvest based on Wildlife Habitat requirements is under the direction of the Wildlife Branch
	 Forest Operations shall be undertaken in compliance with existing regulations In consultation with the outfitter, the timing forest harvesting can be modified around the main lodge during the season of operation.

 Unless located on registered crown land issued to the specific outfitting business, Bear Bait Stands are considered temporary hunting tools, which do not require protection mechanisms and are not considered permanent fixtures on the landscape. However, through the five year plan development and consultation process, outfitters can identify the location of temporary hunting tools such as Bear Bait Stands in an effort to mitigate any issues prior to implementation.
• The guiding Principles for Resource Roads, Visual, Wildlife and Forest Operations above are applicable for all licensed outfitting businesses within the province. In addition, direct consultations with specific outfitters and the Newfoundland and Labrador Outfitters Association may occur to mitigate specific forestry proposals near the Main Lodge.

4.1.2.8 Tourism / Recreation

TOURISM / RECREATION			
Characterization	Non-timber values such as: hiking, skiing, canoeing/kayaking, ATV/UTV and snowmobiling constitute an important role within the Province for tourism and recreational purposes. The Province has outstanding scenery, varying topography and opportunities for viewing wildlife and flora in a natural setting. The tourism industry in Newfoundland and Labrador has experienced significant growth over the years and is now a significant economic driver, generating \$1.14 billion in revenue in 2019. Newfoundland and Labrador has the resources to compete nationally and internationally with tourist destinations. As such, protection of these resources is		
Critical	vital for continued growth and prosperity.Wilderness		
Elements	 Adventure tourism activities require the existence of wilderness areas. If best management techniques are not applied, forest harvesting may result in the alteration of this feeling of pristine wilderness, which could have short and long term adverse effects. Accessibility 		
	 Accessionity Construction and maintenance of Forest Access roads has both a positive and negative effect. On a positive side, it provides the ability to increase vehicular and ATV/UTV traffic allowing more opportunity for this activity. However, on a negative side, it also has the potential to decrease the value of the experience for those individuals seeking a "remote" type setting. 		
	Viewscapes		
	 Either walking a trail, snowmobiling on a groomed trail or canoeing down a river, the visual experience of the surrounding landbase plays an important role in the overall pleasure of the activity. Over the past number of years, viewscapes have become an integral portion of forest management planning. 		

	 The Trans Canada Highway is a strategically important touring route for the tourism industry. Any development that is near this touring route and would impact scenic settings, must be managed to minimize the negative visual viewscapes.
Guiding Principles	 Wilderness Forest operations will avoid established ecological reserve areas and will make every attempt to consult with local stakeholders in areas of high concentrations of tourism and recreational activities. In such areas, stakeholder meetings could prevent conflicts through temporal scheduling. Limiting Accessibility Where possible in sensitive areas, forest harvesting will be scheduled during winter months and take advantage of winter road construction techniques. Winter roads usually restrict vehicular traffic and decommission naturally. In addition, decommissioning of regular forest access roads near sensitive areas is a possible option when forest operations are completed.
	 Viewscape Where possible within areas where high concentrations of tourism and recreational activities occur, negative impacts to viewscapes could be managed using landscape design techniques. This could mean that forest harvesting operations employ: treed buffers, tree retention methods or implement reforestation activity immediately to return the site to a forested condition. Viewscape analysis using computer modelling has been completed in areas along the Exploits River and the Trans Canada Highway to minimize the initial visual effects of Forest Harvesting. The guiding principles for Wilderness, Accessibility, and Viewscapes will apply to all crown operations within the province.

SECTION 5 MITIGATIONS

5.1 General

Best Management Practices adopted from previous planning processes to be incorporated into this plan

- A 30 m buffer will be maintained on both sides of any other rivers, brook, ponds or other water bodies that are shown on 1:50,000 topographic maps.
- There will be no cutting buffer within 100 meters of the Newfoundland T'Railway.
- There will be no cutting buffer within 100 meters of a cabin development area and 30 meters of an approved cabin.
- Scheduled salmon rivers will be evaluated on a site-by-site basis and buffers will vary in width from 30 -100 meters.
- Within protected water supplies, there will be no cutting within 150 meters of the intake pond or stream and no cutting within 75 meters of the main river channel. There will be no cutting within 50 meters of all ponds and streams flowing into the intake pond or stream.

Local known stakeholders were notified as per section 6. Some further information was required as indicated below

DATE	Community	Reason for Contact	Action for Forestry	Results
05/27/21	Town of Fortune	-CC03501: Unable to open files. They recognize Adobe files. Solution: save kmz attachements directly to harddrive then open google earth and add the files through google earth's file tab	-Reached out to municipality and walked them through the process. Sucessfully opened files at the Municipality	no further action
05/28/21	Garnish Pond Road Committee	-CC03508: Irene Hodder emailed regarding information from Town of Marystown regarding five year plan and they would like to be consultated. They would like a No Cut 30m buffer along the Garnish Pond Forest Access Road.		
05/26/21	Port Blandford	-CC02014: May 26/21 Email from the Town of Port Blandford, have ongoing concerns with harvesting in the Southwest River Valley. Specifically with viewscape concerns that might affect Tourism in the area.	alley. town representatives. Town agreed to provide vantage	
06/16/21	Canning's Cove	-CC02531: June 16/21 Mel Simmonds responded by email with LSD having no concerns as long as the propper buffer zone is	-June 18/21 Emailed response to Mel Simmonds. 'No Cutting Within 100m of Clode Sound' marked directly on	no further action
05/25/21	Clarenville	-CC02505, CC02507: 2020 Anonymous requested to extend Block CC02505 to the boundary of Block CC02507.	-May 25/21 Decision made in last planning period to separate Domestic Block CC02505 and CC02507, the District Forest Manager recommends to not extend these Domestic	no further action
06/16/21	Clarenville	-CC02507: Council Requested meeting with Forestry for June 29/21.	-June 18/21 Emailed Rick Well and Public Works Clerk that Tim and Inga to attend Council meeting on June 29/21. Discussion map was provided to the Town.	ongoing discussions

DATE	Community	Reason for Contact	Action for Forestry	Results
06/04/21	Come By Chance	-CC02502: June 4/21 Colin Holloway with the Town of Come By Chance emailed regarding a CL referral that was located in the Domestic Block. As well, wondering if we thought of expanding Block 02.	-June 18/21 Emailed response to Colin Holloway that the Crown Lands Application was revised and it didn't include the portion located in the Domestic Block with the revision. The Provincial LandUse Atlas shows this change as well. The District Forest Manager recommends to not expand the Domestic Block as the transmission line provides a good boundary for permit holders to follow. As well, the number of permits issued for this block and our inventory of volume of wood available indicates a sufficient supply of firewood to accommodate the average number of 30 permits issued per year."	no further action
06/11/21	Dunfield	-CC02559:June 11/21 letter returned, no such addresss	-No other contact info found for LSD	no further action
12/08/21	Hickman's Harbour	-CC02504, CC02562: Dec. 8/20 David Vardy requested the Memorial Trail around Long Pond be put on the Domestic Maps so a No-Cut buffer will be implemented.	-Dec. 22/20 Letter sent indicating the the District Forest Manager recommends to show the Memorial Trail on Domestic Block CC02504 and CC02562 Domestic maps.	no further action
05/25/21	Hillview	-CC02524: 2020 Anonymous requested Block CC02524 be extended to the northwest.	-May 25/21 The the District Forest Manager recommends to not extend Domestic Block CC02524 as the powerline provides a definitive boundary for permit holders to follow.	no further action
05/25/21	Trinity	-CC02559: 2020 Anonymous requested area around old Trinity Loop Park be added into Block CC02559.	-May 25/21 The District Forest Manager recommends to not extend the Domestic Block around the old Trinity Loop Park as the area was taken out during the last planning period and proposals are being made to make use of the existing park.	no further action
05/31/21	Trinity Bay North	-CC02557: May 31/21 - Emailed with concerns of no cut buffers in PPWSA - THMs are high due to organic matter getting into water.	-May 31/21 - Emailed a copy of the PPWSA Development Permit for the Domestic Block to Darryl Johnson. The Town will be referred again for the PPWSA Development Permit in the Fall of 2021. Advised that if they have additional conditions, to contact us.	Apply for PPWSA Development Permit in Fall 2021
04/20/21	Charleston	-CC02548: April 20/20 Mr. Hayward Pike sent a letter wanting a section north of Block CC02548 reserved for No Cutting.	-April 28/20 District called Mr. Pike, explained the FYOP and consultation process. This area was approved for commercial harvest and was scheduled to be harvested in 2020.	no further action
05/25/21	Thorburn Lake	-CC202571: 2020 Eric Stephenson requested Block CC02571 be extended to the plantation.	-May 25/21 The District Forest Manager recommends not to extend the Block CC02571 as young wood in the area will be merchantable when the plantation is available for Commercial Harvest. As well, many cabin owners want the size of the block decreased. Domestic Block CC02571 inventory shows there is enough wood to accomodate the number of permit holders for this block. -June 30/21 emailed Fric Stephenson the District Forest Manager recommends to not extend the Domestic Block.	no further action
09/20/19	Thorburn Lake	-CC02571: Sept. 20/19 Email from Lloyd Avery representing the Thorburn Lake North Cabin Owners requesting removal of a section of Domestic Block CC02571 on the north side of Thorburn Lake. Nov. 20/19 Email from Boyce Mews/Mike Penney representing another group of cabin owners around Thorburn Lake and suggesting that cabin owners need access to the entire block and no changes be made.	-Nov 29/19 Email to Lloyd Avery from Tim Andrews informing that any decisions would be deferred to the FYOP planning process. June 30/21 Email to Lloyd Avery that after review the District Forest Manager recommends to not revise the Domestic Block boundaries. -Call made to Mike Penney to inform the decision would be deferred to the FYOP planning process . June 30/21 Email response to Boyce Mews and Mike Penney that after review the District Forest Manager recommends to not revise the Domestic Block boundaries.	no further action

DATE	Community	Reason for Contact	Action for Forestry	Results
05/31/21	Bonavista	-CC02528, CC02568: Emailed that they couldn't open the KMZ files.	-May 31/21 Saved the blocks as pdf's and emailed them back to the Town.	no further action
06/01/21	Fortune	-CC03501: The Fortune Town Council emailed, they do not have any concerns as per discussion at regular meeting of Council held on June 1 st , 2021 with respect to the domestic block relevant to their jurisdiction.	No action required.	no further action
04/19/21	Marystown	-CC03509: The Town of Marystown requested 'The Tolt' be added back in, but Johnny's Mt. to remain out due to visualization issues.	-Jan. 8/21 Email to the Town indicated the District Forest Manager recommends the Tolt be put back into Domestic Block CC03509 and for Johnny's Mt. to remain out.	no further action
06/03/21	Parker's Cove	they have in their Town that is in Domestic Block CC03511. They are concerned the trail will be cut out by Domestic Permit holders.	-June 3/21 Returned Ambrose Abbotts call, the District Forest Manager recommends to put the trail on the Domestic Map and it will have a 30m no cut buffer as stated in the conditions of the domestic permit.	no further action
10/16/21	St. Lawrence	east of Water Street East and north of Route 220 as well as no	-Jan. 01/21 The District Forest Manager recommends to have no cutting 2 km east of Water Street East and north of Route 220, as well as no cutting within 2 km of Salmonier Pond on Route 220.	no further action
05/31/21	Lethbridge		-April 12/21 Forestry and Wildlife Branch responded to Crown Lands Referral for the Lethbridge Recreational Trail that this was a commercial harvest areaMay 31/21 Through phone call, the District Forest Manager recommends to work with LSD to protect the trail as part of the harvesting plan.	ongoing discussions
06/22/21	Lower Lance Cove	-CC02504, CC02512: June 22/21 Leslie Ivany, LSD chair, emailed with concerns of protecting area in proposed PPWSA extension.	-June 30/21 Emailed Leslie Ivany that the District Forest Manager recommends to revise the Domestic Block boundaries of Blocks CC02512 and CC02504 to exclude the drainage area surrounding Stickley's Pond, Figure Eight Pond and Long Narry Pond.	Adjustments to be made to Domestic Block CC02512 and CC02504. No further action
01/13/21	Hodges Cove		-Jan 13/21 Responded to Crown Lands Referral that the District Forest Manager recommends to show the hiking trail on the Domestic map and that there would be a 30m No Cut buffer around Traditional Paths hiking trail in Hodges Cove.	Hiking Trail to be shown on Domestic Block ma No further action

DATE	Outfitter	Reason for Contact	Action for Forestry	Results
6/2/2021	Deep Country Lodge	Email received indicating owner	Sent more detailed pdf maps	There were no proposed activity located within the
		wanted to be fully consultated on	indicating location of the 4 lodges	general vicinity of 3 lodges. No further action required.
		plans	under this business.	With respect to the one lodge near the proposed

SECTION 6 PUBLIC CONSULTATION

6.1 Planning Framework

Forest Resource managers in Canada are striving for a society that successfully integrates economic, environmental and social considerations into all resource-related decision making. Since the early 1990's, there has been a country-wide shift from single resource management to a more comprehensive approach of forest ecosystem management. Sustainable Forest Management (SFM) must be balanced in light of social, economic, and environmental issues. In the context of SFM, this shift has resulted in a move from the traditional narrow focus of timber management, to incorporate non-timber values into the management planning framework. Another term that has become closely associated with SFM is "sustainable development" or in this case "sustainable forests", which not only takes into account the social, cultural, economic, and environmental benefits of the present, but those of future generations as well. Involvement of

Interested Stakeholders into the five-year planning process is recognized by the Forestry Services Branch as a key component to achieving sustainable development.

As a result of the 1995 Environmental Preview Report, the Forestry Services Branch adopted an adaptive management planning process, which has three objectives:

- Establish a productive planning framework to include all interested stakeholders. An effective planning framework must have information and defined spatial issues.
- Learn more about forest ecosystems while they are being actively managed (i.e. adaptive management). Adaptive management incorporates strategies which help us learn about the forest ecosystem and to deal with uncertainties.
- Establish an ecosystem approach to forest management which integrates the scientific knowledge of ecological relations and limits of growth with social values. This will help to attain the goal of sustaining natural ecosystem integrity and health over the long term.

Adaptive management makes decisions based on input from interested stakeholders and establishes a continuous learning program. The adaptive approach allows us to communicate, share information and learn about forests being managed. This sharing of information, both old and new, then provides the flexibility necessary to adjust to changes and to set new goals. Such interaction is an absolute necessity for a subject as complex as an ecosystem.

More information on the Forest Management Planning Process can be found on Governments Forestry website using the following addresses

(https://www.faa.gov.nl.ca/forestry/managing/district.html) and

(<u>https://www.faa.gov.nl.ca/forestry/managing/public_consultation.html</u>), as well as Governments Engage NL website.

6.2 Stakeholder Involvement

Since the mid 1990's, for each five-year plan, the Forestry Services Branch embarked upon a rigorous public consultation process involving a series of meetings spanning a number of months at an established venue, where interested stakeholders could discuss a range of forest management issues at an operational level.

With respect to the strategic level, in 2014, the Forestry Services Branch released a 10- year Provincial Sustainable Forest Management Strategy (PSFMS) Document (2014-2024), which emerged through wide consultation with citizens of the Province. The 2014-2024 PSFMS builds on the strengths of the previous strategy plans and uses a landscape-scale planning approach to implement the progressive and innovative ecological policies required for Sustainable Forest Management (SFM). The strategy builds on the strengths of the many modern and high-quality forest management programs that are currently being implemented in this province to ensure a vibrant and competitive forest industry.

Taking into account the many five-year plans successfully implemented within the province since the mid 1990's through public consultation processes and the recent PSFMS developed through public consultations, The Forestry Services Branch strives to improve its methods to garner advice from the public while also mitigating land-use conflicts. To this effect, as new five-year plans are being developed and implemented provincially, relevant issues raised from previous planning processes are considered the foundation the new plans.

A new stakeholder involvement process was initiated in 2020 utilizing the department of Engage NI. Through this process, interested individuals could either contact forestry directly or complete an online form that specifically indicates a particular issue or concern. For Zone 2, an Engage NI consultation process was targeted and directed stakeholder involvement began on May 25, through emails to targeted groups. Information regarding the proposed forest management plan, and associated google earth files were provided

Clarenville	Lewin's Cove	
King's Cove	Lord's Cove	
Port Blandford	Marystown	
Baine Harbour	Musgravetown	
Bay L'Argent	Parker's Cove	
Bonavista	Point au Gaul	
Burin	Point May	
Come By Chance	Port Rexton	
Elliston	Red Harbour Rushoon	
English Harbour East		
Fortune	St. Bernard's	
Burin		
Frenchman's Cove	St. Lawrence	
Garnish	Sunnyside	
Grand Bank	Terrenceville	
Grand Le Pierre	Trinity	
Keels	Port Union	
Lamaline	Winterland	
Lawn	George's Brook-Milton	

1) Town Councils - email sent on May 25, 2021

2) Local MHA's – email sent on May 25, 2021

3) Outfitters / Tourism - email sent on May 25, 2021

Deep Country Lodge	Nousfoundland and Johnsdon Outfitters Accessible a
Efford's Hunting Adventures (Thornburn Lake Lodge)	Newfoundland and Labrador Outfitters Association
Efford's Hunting Adventures (Triton Brook Camp)	Newfoundland and Labrador Snowmobile Federation
Beaulieus Caribou Hunts (2005) Limited (Caribou Pond Camp)	Hospitality NL
Beaulieus Caribou Hunts (2005) Limited (Long Harbour River Lodge)	Legendary Coasts NL DMO
Canadian Northern Outfitters	Adventure Central DMO
Efford's Hunting Adventures (Island Pond West Lodge)	Go Western DMO

- 4) Indigenous Leaders email sent on May 25, 2021
 - Qalipu First Nation
 - Miawpukek First nation

SECTION 7 MANAGEMENT OBJECTIVES AND STRATEGIES

7.1 Harvesting

The boreal forest is characterized by natural stand replacement following a natural disturbance, which results in the formation of relatively even aged stands. A forest management technique called the Clearcut Silvicultural system is utilized as it closely emulates this natural disturbance pattern. The size, shape, arrangement and juxtaposition of harvest areas vary across the landscape depending on localized topography and terrain conditions.

7.1.1 Commercial

The older unalienated timber considered in the worst condition is targeted as a high harvest priority, followed by stands that have been damaged by insects and disease. In managed stands, this priority changes to allow for a faster rotation on sites that have been silviculturally treated.

Specific commercial strategies include:

- design irregular cut blocks that follow contours and natural boundaries
- vary buffer widths to protect other values (ie. larger buffers on salmon rivers)
- utilize winter harvest on wet and sensitive sites
- maintain current size and distribution of clear cuts
- where possible, maintain unharvested strips between harvest blocks as wildlife utilization corridors
- use landscape design techniques to mitigate viewscapes
- minimize timber utilization loss (< 6 m3/ha)

7.1.2 Domestic

The harvest of domestic fuelwood and sawlogs occurs from designated areas, recent commercial harvest areas and sporadically throughout the zone with the removal of blowdown trees. Utilization of cutover residue, dead timber and scrub areas do not form part of the woodsupply analysis. Generally, domestic cutting areas are generally established near communities. However, within areas of the zone not covered by any operating areas, domestic permits may be issued to remote cabin owners for firewood as requested. The number of permits and volume associated with these permits will be extremely low and insignificant.

Specific domestic strategies include:

- target low volume stands having poor commercial opportunity
- encourage use of poor quality hardwood (birch, larch and aspen). In areas where there
 are future softwood commercial operations, domestic harvesting is limited to noncommercial hardwoods
- target dead, burnt and insect damaged stands that are beyond commercial salvage throughout the zone and specifically within 100 meter buffer of highways.
- target alienation class 3 lands that have low commercial opportunity
- in areas of high domestic demand, limit volume allocation in designated cutting areas and encourage alternate sources (cutovers, landings, scrub etc)

7.2 Silviculture

As a general rule, approximately 80% of the Boreal Forest regenerates naturally following a disturbance. Forest renewal management programs are applied by forest managers within the 20% that do not successfully natural regenerate. Forest renewal silvicultural treatments are designed to help facilitate a new forest after disturbances caused by harvesting, insect, wind or fire. These prescriptions can involve either Site Preparation (scarification), Planting or Pre-Commercial Thinning.

Site Preparation

When a site does not regenerate at all, a full planting program is required. In some cases, the site may need to be manually prepared to aid in the establishment and growth of the planted seedlings (generally black or white spruce and to a lesser extent, norway spruce). Site preparation techniques can include:

- Mechanical site preparation (scarification) involves using heavy equipment (skidder) equipped with special attachments to reduce the thickness of the duff layer, and remove or disturb any kalmia that is present, which would restrict seedling growth.
- Prescribed burning is used to sanitize some sites where adelgid is present. This treatment reduces the slash loading and duff thickness to prepare the site for planting and kills any balsam fir which could potentially perpetuate the adelgid problem.
- Treatment to prepare sites that have been overgrown with hardwoods and other herbaceous species has been done with herbicides to reduce this competition, making the site more accessible and suitable for planting. Release herbicide treatment reduces the competition for a few years to allow planted seedlings to get established. In other

instances, herbicides are used to control Kalmia either before or after planting. Herbicides, while used sparingly, are sometimes a necessary tool to help establishment of a new forest, particularly on the better sites.

Planting

A full planting technique is required when no regeneration occurs to ensure regeneration of selected tree species is at acceptable levels. Gap planting is normally achieved with spruce seedlings, coupled with the natural regeneration already present on site to increase seedling density to acceptable levels.

On adelgid sites partially regenerated to balsam fir, planting is done through the existing regeneration to obtain a sufficient stocking level of an adelgid resistance species. However, where adelgid has been a problem, balsam fir regeneration is sometimes ignored and the site is planted with spruce seedlings.

Where possible, seedlings used in the silviculture program are grown with seed from local sources. Seed orchards have been established at Pynns Brook and Wooddale to produce seed from plus trees collected throughout the province. Plus trees are normally selected because they have superior growth and physiological characteristics. It is hoped that once the orchard is in full production, the majority of the planting stock will be grown from this source. The ultimate goal is to plant seedlings that have superior growth characteristics and thus increase yield and maintain genetic diversity.

Exotic species have been planted in trials at some locations in the zone. However; it is not anticipated to form any substantive proportion of the planting program in the foreseeable future.

Thinning

In an attempt to enhance development, silviculture thinning programs are designed to treat established forest stands.

Pre-Commercial Thinning (PCT) usually involves partial removal of overstocked balsam fir stands at a young age 10 -15 years. In areas which have high moose browsing potential, the age is increased to 20 – 25 years, so that crop trees are tall enough to be out of reach of moose. PCT reduces density levels which facilitates maximizing volume increment and operability (piece size). Trees removed are not of merchantable size and remain on site, returning the nutrients back into the soil. In the zone, balsam fir is usually thinned to favour any spruce present within the stand. This prescription results in a mixed softwood stand (depending on the original density of spruce) which is more diverse and less susceptible to insect infestation. As well, any hardwood species that are not in direct competition with spruce or fir are left to increase the biodiversity of the stand.

Commercial and Diameter Limit thinning would occur in the intermediate age 25 - 35 years and is undertaken in older balsam fir stands (either natural or previously thinned). It is designed to capture any mortality that would normally occur in the stand through self thinning. The trees removed from commercial thinning operations are extracted and utilized. The remaining trees are left to grow, free from competition and are harvested when mature. As with PCT, spruce and hardwoods are left where possible to increase the stand diversity.

Thinning programs aim to shorten the rotation period of a stand and produce large diameter stems. This program should increase the percentage of merchantable volume considered suitable for sawlogs. Commercial thinning has not been completed in the zone and diameter limit thinning has been done sparingly. In recent years the precommercial thinning program has dropped significantly. This trend is expected to continue.

More information on the Silviculture Program can be found on Governments Forestry website using the following address

(https://www.faa.gov.nl.ca/forestry/managing/silviculture/index.html)

Specific silviculture strategies include:

- ensure regeneration of areas disturbed by harvest, insect, wind and fire to prevent loss of productive land base
- use thinning techniques in young stands to promote enhanced stand development, reduce rotation age, and increase the percentage of sawlogs
- leave hardwoods, where possible, in pre-commercially thinned areas to increase stand diversity
- where possible, promote species mixes particularly with spruce and hardwoods to reduce susceptibility to insect attack and increase biological diversity
- where possible, use seedlings grown from local seed sources to protect genetic diversity
- ensure levels of planting and thinning used in the wood supply analysis are achieved
- work towards pre harvest planning to identify areas with potential silviculture problems so that optimal prescriptions can be promptly employed

7.3 Access Roads

Forestry roads are required to gain access to scheduled commercial harvest areas. Access roads also provide opportunities for other recreational and commercial values such as: hunting, fishing, skiing, berrypicking, hiking, outfitting, cabin development and mineral exploration.

As a general rule of thumb, only the minimal amount of access roads are constructed to effectively and efficiently conduct commercial harvest operations. Access roads are constructed to specifications minimizing right-of-way and running surface width. Forwarding distances will be maximized to curtail the overall amount of road constructed. In sensitive and wet areas, winter harvesting and road construction are encouraged. Following these principles helps to ensure the minimum amount of road will be constructed, reducing the loss of productive forest landbase and minimizing environmental disturbance. Road and bridge maintenance and/or decommissioning are considered depending on cost, and mitigation of conflicting uses for a particular road. More information on the Roads Program can be found on Governments Forestry website using the following address (<u>https://www.faa.gov.nl.ca/forestry/roads/index.html</u>)

Specific roads strategies include:

- construct winter roads in sensitive and wet areas
- minimize amount of road built by maximizing forwarding distances
- use minimum road standard to safely and effectively match the logging chance
- consider road decommissioning on roads near remote outfitting lodges and other areas of concern where requested and where feasibly possible
- determine impacts and explore alternatives (cost sharing) in areas where road and bridge decommissioning impacts other stakeholders
- explore all avenues to secure funding for road construction and encourage operators to build their own roads in exchange for royalty reductions

7.4 Forest Protection

7.4.1 Insects and Disease

As indicated in section 1.5.5, insects have been considered a major natural disturbance within the zone. Balsam fir is susceptible to most of the major insects including spruce budworm, hemlock looper, and balsam woolly adelgid. In the event of a major insect infestation, salvage efforts may change harvest priorities. However, deviations from harvest schedules will be closely monitored to ensure that the validity of the AAC is not compromised.

Monitoring and protection programs for insects and disease are coordinated by the forest protection division in Corner Brook. Local district staff provide assistance in detection, monitoring, and protection surveys against insects and disease. More information on the Forest Insect Control Program can be found on Governments Forestry website using the following address (https://www.faa.gov.nl.ca/forestry/idc/index.html)

Specific insect and disease strategies include:

- use silvicultural techniques at the stand level to alter species mix and increase stand vigor to make stands less susceptible to insect attack
- where possible, use harvest scheduling techniques to alter species mix across the landscape to minimize potential for severe insect infestation
- in conjunction with provincial and federal initiatives, use pertinent and approved insecticides

7.4.2 Fire

There has been a cyclic fire history in the zone. A fire in an unusually dry year can have devastating effects on the forest and can exacerbate an established wood supply. The risk of a serious forest fire can minimized by maintaining a highly trained, efficient and effective fire control program and by minimizing the risk in forest stands through maintenance of forest health and vigour. Within the zone, there have been major forest fires in the past. However, in recent years, wildfire has not been a major issue. There have been some minor wildfire's but all have been quickly contained and not much timber has been lost. The Department of Fisheries and Land Resources is committed to protection of the resource and continues to invest in a fire suppression program to ensure any future losses are minimized. There are fire crews and equipment stationed at local forestry depots within the zone during the forest fire season, whose direct responsibility is forest fire protection. In addition, support, equipment and manpower at both the regional and provincial level is available should the need arise. There are air tankers stationed at Deer Lake and Gander and helicopters in Gander that are available for initial attack. More information on the Forest Fire Program can be found on Governments Forestry website using the following address (https://www.faa.gov.nl.ca/forestry/forest_fires_index.html)

Specific fire strategies include:

- use silvicultural treatments and protection from insects to increase health and vigour of stands
- maintain fire control capabilities
- promote species mixes in stands to minimize risk

7.4.3 Windthrow

Wind throw usually occurs in older stands that have been predisposed by some other disturbance such as insects and disease. To minimize the effects of Windthrow (blowdown), stands will be managed to promote forest health and vigour mainly through silvicultural treatments and protection from insects.

Specific windthrow strategies include:

- avoid thinning in areas with high wind damage potential (hilltops on high elevations etc.)
- maintain forest in healthy vigorous condition through silvicultural treatments and protection from insects
- design cut blocks to follow contours and natural boundaries to minimize risk of windthrow to residual forest

7.5 Environmental Protection & Climate Change

7.5.1 General Environment

The Department of Fisheries and Land Resources have developed has developed an Environmental Management System (EMS) that is registered with the International Standards Organization (ISO). As part of this process, an EMS Policy was developed and proper operating procedures developed for various forest management activity. Initial registration was on December 17, 2015 and through regular monitoring and audits (internal and external), the EMS

remains registered. Under the EMS, the department has developed stringent operating procedures for fuel handling, working around waterbodies, and overall pollution prevention. In addition, inspection programs are implemented are implemented to evaluate forest operations and rectify any deviations from established protocols. More information on the EMS can be found on Governments Forestry website using the following address

(https://www.faa.gov.nl.ca/forestry/ems/index.html)

To ensure forestry activity is conducted to minimize any potential negative impacts to the environment, operating procedures and best management practices called Environmental Protection Guidelines (EPG's) have been developed and implemented across the province. Highlights of measures to avoid these impacts include no activity buffer zones, modification of harvesting design and equipment, avoidance of sensitive site during critical periods, consultation with other regulatory agencies, and monitoring. More information on EPG's can be found on Governments Forestry website using the following address

(<u>https://www.faa.gov.nl.ca/forestry/managing/pdf/Environmental-Protection-</u>Guidelines.pdf)

Through implementation of the EMS and the EPG's, the department strives to be responsible stewards of the landbase. As well, the programs illustrated in this document relating to forest protection from Insects and Fires, help to maintain a forested landbase. As indicated in previous sections, harvested sites are evaluated for regeneration potential and proper reforestation techniques are implemented to facilitate tree growth. Maintaining and achieving a stocked forest at the earliest timeframe help provide for carbon storage.

7.5.2 Climate Change

From an initial review, it appears there are two options for incorporating carbon accounting into our standard wood supply modelling process. First, there is the option of using the National carbon accounting model CBM3. CBM3 accepts outputs from Woodstock, the wood supply model used by the Department, and from these Woodstock outputs generated the expected carbon flows. The advantage of this approach is that CBM3 is an internationally recognized model that is used nationally to project carbon flows for Canada. The disadvantage of the CMB3 that the model is that can only process Woodstock models with 10 or less landscape themes. A landscape theme is a feature used to describe or characterized the area being modeled. Our present Woodstock model has 13 themes and reducing the themes down to 10 will not meet some of our planning commitments. There have been scripts developed by the Federal scientists that can convent models with more than 10 themes down to 10 themes models, however using these scripts requires additional data processing and adds complexity to the model structure.

The second options involves developing similar carbon curves as the ones used in CBM3 and incorporating them into Woodstock itself. With these carbon curve in place it will be possible to tracking carbon outputs similar to the CBM3. The advantage of the carbon curve approach is it reduces over all modelling complexity; however, this approach does not have the "brand recognition" of the CBM3 modelling approach.

The plan going forward is to hire a wood supply analyst who will spend 50% of his time evaluating the two approaches highlighted above, or any other approach that becomes apparent, and

recommend the approach that will works the best for the Department. The remaining 50% of the analyst time will be spending supporting the ongoing Province wood suppy review.

7.5.3 Surveys

Utilization surveys will be conducted on both commercial and domestic cutovers to ensure loss of merchantable timber is minimized. Results of these surveys will be used to evaluate the expected volume in an operating areas to those actually attained. The results of this survey will help refine inventory deductions in future woodsupply analysis.

Reconnaissance and intensive regeneration surveys will be conducted on commercial cutovers in this upcoming five year period, and as well as those created in the past five years to determine the requirement for silvicultural activity. Reconnaissance surveys will be completed on regenerating stands to determine the suitability for pre-commercial thinning.

7.6 Information and Education

Information and education is one of the key elements to providing active and effective participation in the planning process at all levels. Through interaction with various user groups and the general public, a better understanding of ones values and positions is gained. The more we know about other values and their location, the better the ability to mitigate any potential negative impacts. For example, learning where a cabin is located can help planners when selecting areas for harvest and provide a contact to discuss impacts and mitigations. Districts within the zone will continue to educate the general public and engage in meaningful consultations with interested stakeholders where applicable. Annual National Forest Week activities provides a great opportunity for interested individuals to gain a greater understanding of the Provinces' Forest.

Sources of information can include:

- government website
- field trips
- school visits
- information meetings
- general day to day contact

SECTION 8 PROPOSED ACTIVITIES

8.1 overview

This section will outline all forest activities that will occur on crown land in the zone from 2022-2026, including: proposed commercial and domestic harvesting, silviculture, access road construction, and activities proposed within protected water supply areas.

To present a more comprehensive overview of proposed activities on the entire district an overview map is presented in Appendix 3. This map shows all proposed operating areas so that operations can be viewed from a landscape perspective. Maps of individual operating areas and summary sheets are also presented in Appendix 3. The summary sheets give a brief description of each area, the type of activities that will occur and any issues raised and mitigative measures employed.

Digital copies of the Zone 2 plan can be found on the governments forestry website at the following address

(https://www.faa.gov.nl.ca/forestry/managing/public consultation.html)

8.1.1 Allocation of Timber Supply

Table 8 below indicates the scheduled proposed forest harvest for the upcoming five year period 2022-2026. Note: the total volume identified maybe over the final AAC for each forest management district, but when scheduling activity, the maximum sustainable harvest over the five year period will not be exceeded.

		PROPOSED HARVEST TOTAL VOLUME m ³								
HARVEST TYPE	Class 1 softwood	Class 3 Softwood	Hardwood	TOTAL						
COMMERCIAL	1,073,754	23,321	97,634	1,194,709						
DOMESTIC FMD02	210,880			210,880						
DOMESTIC FMD03	130,320			130,320						
TOTAL	1,414,954	23,321	97,634	1,535,909						

Table 8: Proposed forest harvest in Zone 2 from 2022-2026

*note: when final AAC's are developed for this zone, if the above schedule represents more than the established AAC, the harvest will be monitored to ensure established AAC's are not exceeded.

8.1.1.1 Commercial

The timber scheduled for commercial harvest in the district is overmature with some small pockets of mature dispersed throughout. This proposed harvest approximates the harvest schedule that was used to determine the AAC. The allocated operating area and associated harvest volumes represent as much as two times the actual proposed harvest. The purpose of including more

volume than is actually proposed is to allow for operational flexibility and inventory deviations within operating areas without having to constantly amend the plan. These operations occur manually or mechanically using conventional harvesting equipment such as chainsaws, shortwood harvesters, skidders and forwarders and are conducted year round. The more sensitive sites are usually harvested in winter and most operations are integrated utilizing sawlogs, pulpwood and fuelwood.

Opora	ting Area		FMD 02 (Crown Propos	ed Commer	cial Harvest	t (m ³) 2022-20	26	
Opera	ling Area		Softw	vood			Hardwo	bod	
Number	Name	Core	Operational	Total	Area (Ha)	Core	Operational	Total	Area (Ha)
CC02001	Goobies	3860		3860	37.56	176		176	37.56
CC02002	Hatchet Cove	10642	897	11539	141.53	818	37	855	141.53
CC02003	Burnt Point	18930		18930	209.82	2078		2078	209.82
CC02004	Snook's Harbour	4219		4219	43.44	451		451	43.44
CC02005	Foster's Marsh	23810		23810	244.12	2375		2375	244.12
CC02006	Hickman's Harbour Road	18756		18756	183.38	740		740	183.38
CC02008	Northwest Brook	5024		5024	63.27	2575		2575	63.27
CC02009	Beaver Pond	19343	3335	22678	231.91	1179	263	1442	231.91
CC02010	Clarenville	37452		37452	326.47	4178		4178	326.47
CC02011	White Hills	49824		49824	480.48	2283		2283	480.48
CC02012	Shoal Harbour Pond	91924	1972	93896	798.33	8576	716	9292	798.33
CC02013	Thorburn Lake South	10355	304	10659	101.52	433	15	448	101.52
CC02014	Southwest River Valley	115163		115163	902.72	5450		5450	902.72
CC02015	Port Blandford Access Road	11910		11910	108.23	978		978	108.23
CC02016	Thorburn Lake North	5420		5420	50.12	476		476	50.12
CC02017	George's Pond	4273		4273	39.42	211		211	39.42
CC02018	Salt Lake	6299	2580	8879	100.06	903	200	1103	100.06

Table 9: FMD 02 Proposed Commercial Harvesting from 2022 to 2026

CC02019	Shoal Harbour Valley	5326		5326	59.34	533		533	59.34
CC02020	Muddy Hole Brook	10988		10988	124.6	1461		1461	124.6
CC02021	George's Brook	6074		6074	79.83	630		630	79.83
CC02022	Barton	771		771	12.5	24		24	12.5
CC02023	Harcourt Access Road	18694		18694	287.99	1214		1214	287.99
CC02024	Jack's Pond	1237		1237	16.83	67		67	16.83
CC02025	Nut Cove	3290	2032	5322	59.08	373	81	454	59.08
CC02030	Island Pond	1322		1322	18.62	109		109	18.62
CC02031	Bloomfield Access Road	2167		2167	31.54	119		119	31.54
CC02032	North Brook	20282		20282	178.58	2931		2931	178.58
CC02033	Branches Pond Access Road	17997		17997	238.24	2974		2974	238.24
CC02036	New Country	9923		9923	117.05	869		869	117.05
CC02037	Dalton's Pond	12189		12189	161.99	1170		1170	161.99
CC02038	Rocky Pond	8992		8992	138.87	2149		2149	138.87
CC02040	Rattle Falls	14702		14702	153.21	526		526	153.21
CC02041	Princeton Pond	26472	252	26724	348	1637	10	1647	348
CC02042	Blue Gull Pond	2808		2808	38.68	52		52	38.68
CC02043	Portland	26106		26106	422.28	8289		8289	422.28
CC02044	Winterbook	13240		13240	217.32	3054		3054	217.32
CC02045	Bakeapple Bog	13158	3665	16823	231.99	3277	649	3926	231.39
CC02046	Chance Harbour	15216		15216	230.06	1907		1907	230.06
CC02047	Charleston	28252		28252	440.38	3971		3971	440.38
CC02050	Trinity Pond	3555		3555	45.81	822		822	45.81
CC02051	Plate Cove Access Road	198726	1013	199739	1793.64	9131	51	9182	1793.6 4

CC02052	Stock Cove Access Road	32480	730	33210	338.56	1836	37	1873	338.56
CC02054	World Pond	35849		35849	404.36	1906		1906	404.36
CC02056	Black Bay	15141	839	15980	249.5	1723	256	1979	249.5
CC02057	Terra Nova South	7744	5702	13446	106.61	284	179	463	106.61
CC02058	Deep Bight	43996		43996	457.05	4885		4885	457.05
CC02059	Andrew's Pond	35717		35717	269.16	3080		3080	269.16
CC02060	Camp Pond	4136		4136	51.43	257		257	51.43
	TOTAL	1,073,754.00	23,321.00	1,097,075.00	11,385.48	95,140.00	2,494.00	97,634.00	11,384. 88

8.1.2.2 Domestic

Harvesting will occur in designated domestic cutting areas and is generally conducted on a small patch cut system. However, within areas of the zone not covered by any operating areas, domestic permits may be issued to remote cabin owners for firewood as requested. The number of permits and volume associated with these permits will be extremely low and insignificant. All domestic cutting is done under permit which has conditions attached that outline the species, volume, location and utilization standards to be employed. Most cutting occurs in fall and winter with extraction by snowmobile or ATV. Domestic permit allocation is 16 m3.

TADIE TU. FI	Table TO. FIND 02 Proposed Domestic Harvesting From 2022 to 2020.										
	FME	Estimated 5 year volume (m ³)									
Number	Name	Tenure	Total Area (ha)	Number of Permits	Softwood	Hardwood					
CC02501	Sunnyside	Crown		59	4720						
CC02502	Com By Chance	Crown		26	2080						
CC02503	Goobies	Crown		50	4000						
CC02504	Hickman's Harbour	Crown		61	4880						
CC02505	Adeytown	Crown		40	3200						
CC02506	George's Brook	Crown		104	8320						
CC02507	Clarenville	Crown		251	20080						

Table 10: FMD 02 Proposed Domestic Harvesting From 2022 to 2026.

1	I	1	1	1	l	1
CC02508	White Hills	Crown		9	720	
CC02510	Elliott's Cove	Crown		36	2880	
CC02511	Aspey Brook	Crown		14	1120	
CC02512	Britannia	Crown		50	4000	
CC02513	Burgoyne's Cove	Crown		39	3120	
CC02514	Weybridge	Crown		40	3200	
CC02517	Branches Pond Access Road	Crown		10	800	
CC02518	Lethbridge Access Road	Crown		7	560	
CC02519	Harcourt	Crown		35	2800	
CC02520	Monroe	Crown		10	800	
CC02521	Ocean Pond Access Road	Crown		25	2000	
CC02522	Sweet Bay Road	Crown		5	400	
CC02523	Plate Cove Access Road	Crown		18	1440	
CC02524	Hillview	Crown		113	9040	
CC02525	Bunyan's Cove	Crown		77	6160	
CC02526	Hodge's Cove	Crown		89	7120	
CC02527	Trinity Pond	Crown		18	1440	
CC02528	Bonavista	Crown		36	2880	
CC02529	Catalina Road	Crown		19	1520	

1		I	1	1	
CC02530	Lockston Path South	Crown	20	1600	
CC02531	Musgravetown	Crown	102	8160	
CC02532	Port Blandford	Crown	89	7120	
CC02533	Morley's Siding	Crown	15	1200	
CC02534	Hatchet Cove	Crown	25	2000	
CC02535	Bloomfield	Crown	74	5920	
CC02536	Lethbridge	Crown	79	6320	
CC02537	Little Heart's Ease	Crown	46	3680	
CC02538	Open Hall	Crown	11	880	
CC02539	Long Beach	Crown	19	1520	
CC02540	Random Heights	Crown	21	1680	
CC02541	Swift Current	Crown	34	2720	
CC02542	Garden Cove	Crown	18	1440	
CC02543	Lady Pond Access Road	Crown	6	480	
CC02544	North Harbour	Crown	21	1680	
CC02545	Portland	Crown	53	4240	
CC02546	Winterbrook	Crown	37	2960	
CC02548	Charleston	Crown	23	1840	
CC02549	Princeton	Crown	40	3200	
CC02550	Sweet Bay	Crown	15	1200	
CC02551	Summerville	Crown	20	1600	

CC02552	Plate Cove	Crown	39	3120	
CC02553	Duntara	Crown	34	2720	
CC02554	King's Cove	Crown	55	4400	
CC02555	Amherst Cove Access Road	Crown	16	1280	
CC02556	Newman's Cove	Crown	52	4160	
CC02557	Trinity Bay North	Crown	111	8880	
CC02558	Port Rexton	Crown	54	4320	
CC02559	Trouty	Crown	61	4880	
CC02560	Trynor's Pit	Crown	4	320	
CC02561	Chain Pond	Crown	12	960	
CC02562	Robinson's Bight	Crown	18	1440	
CC02563	King's Cove Road	Crown	5	400	
CC02564	Stock Cove	Crown	17	1360	
CC02565	Lockston Path North	Crown	9	720	
CC02566	English Harbour	Crown	9	720	
CC02567	Little Catalina	Crown	14	1120	
CC02568	Elliston	Crown	 30	2400	
CC02569	Chute Brook	Crown	2	160	
CC02570	Amherst Cove	Crown	21	1680	
CC02571	Thorburn Lake	Crown	35	2800	

CC02572	Deer Harbour Pond	Crown	1	80	
CC02573	Deer Harbour	Crown	5	400	
CC02574	Hatchet Cove Access Road	Crown	5	400	
CC02576	Lockston	Crown	18	1440	
CC02577	St. Jones Without	Crown	0	0	
Total			2636	210880	

Table 11: FMD 03 Proposed Domestic Harvesting From 2022 to 2026.

	FM		Estimated 5 year volume (m³)			
Number	Name	Tenure	Total Area (ha)	Number of Permits	Softwood	Hardwood
CC03501	Fortune	Crown		61	4880	
CC03502	Molliers	Crown		112	8960	
CC03503	Lord's Cove	Crown		57	4560	
CC03504	Lawn	Crown		114	9120	
CC03505	Grand Beach	Crown		63	5040	
CC03506	Main Brook	Crown		215	17200	
CC03507	Garnish	Crown		74	5920	
CC03508	Marystown	Crown		2620	20800	
CC03509	Burin	Crown		214	17120	
CC03510	Red Harbour	Crown		74	5920	
CC03511	Boat Harbour	Crown		113	9040	
CC03512	Petite Forte	Crown		16	1280	
CC03513	St. Bernard's	Crown		45	3600	
CC03514	Bay L'Argent	Crown		67	5360	
CC03515	Baie De L'eau	Crown		53	4240	

CC03516	Terrenceville	Crown	23	1840	
CC03517	English Harbour East	Crown	11	880	
CC03518	Grand le Pierre	Crown	27	2160	
CC03519	Monkstown Road	Crown	11	880	
CC03520	Monkstown	Crown	18	1440	
CC03521	Sandy Harbour River	Crown	1	80	
Total			3,989	130,320	

8.1.2.3 Hardwoods

This domestic harvest of birch occurs as a mixture in softwood stands and is utilized as fuelwood. The commercial hardwood harvest is for sawlogs and fuelwood and occurs in some pure stands but mostly as residual in hardwood/softwood and softwood/hardwood stands.

8.1.3 Silviculture

Potential silvicultural treatment areas need to undergo reconnaissance and / or intensive surveys to determine the regeneration level and severity adelgid attack. Such surveys will be conducted during this five year period but until they are completed, specific locations and treatment amounts cannot be identified. However, silviculture prescriptions have been developed for implementation on specific site conditions. Areas that are scheduled for commercial harvest or have been harvested are identified on the operating area maps and are candidates for planting or gap planting to black, white or Norway spruce. These areas will undergo reconnaissance and or intensive regeneration surveys to determine the need for planting and the presence of adelgid.

Site preparation using either mechanical methods or prescribed burning will be employed on suitable sites having impediments to planting. On black spruce cutovers where kalmia is present, mechanical site preparation (row scarification) or prescribed burning will be used to disturb the kalmia and create suitable microsites to plant black spruce. In fir areas, burning is a preferred treatment to sanitize the site of any existing adelgid infested trees.

FMD	BLOCK ID	NAME	PROPOSED TREATMENT
2	CC02001	Goobies	Scarification/Planting
2	CC02002	Hatchet Cove	Scarification/Planting
2	CC02003	Burnt Point	Scarification/Planting

Table 12: FMD 02 Proposed Silviculture treatment 2022-2026.

2	CC02004	Snook's Harbour	Scarification/Planting	
2	CC02005	Foster's Marsh	Scarification/Planting	
2	CC02006	Hickman's Harbour Road	Scarification/Planting	
2	CC02008	Northwest Brook	Scarification/Planting	
2	CC02009	Beaver Pond	Scarification/Planting	
2	CC02010	Clarenville	Scarification/Planting	
2	CC02011	White Hills	Scarification/Planting	
2	CC02012	Shoal Harbour Pond	Scarification/Planting	
2	CC02013	Thorburn Lake South	Scarification/Planting	
2	CC02014	Southwest River Valley	Scarification/Planting	
2	CC02015	Port Blandford Access Road	Scarification/Planting	
2	CC02016	Thorburn Lake North	Scarification/Planting	
2	CC02017	George's Pond	Scarification/Planting	
2	CC02018	Salt Lake	Scarification/Planting	
2	CC02019	Shoal Harbour Valley	Scarification/Planting	
2	CC02020	Muddy Hole Brook	Scarification/Planting	
2	CC02021	George's Brook	Scarification/Planting	
2	CC02022	Barton	Scarification/Planting	
2	CC02023	Harcourt Access Road	Scarification/Planting	
2	CC02024	Jack's Pond	Scarification/Planting	
2	CC02025	Nut Cove	Scarification/Planting	
2	CC02030	Island Pond	Scarification/Planting	
2	CC02031	Bloomfield Access Road	Scarification/Planting	
2	CC02032	North Brook	Scarification/Planting	
2	CC02033	Branches Pond Access Road	Scarification/Planting	
2	CC02036	New Country	Scarification/Planting	
2	CC02037	Dalton's Pond	Scarification/Planting	
2	CC02038	Rocky Pond	Scarification/Planting	
2	CC02040	Rattle Falls	Scarification/Planting	
2	CC02041	Princeton Pond	Scarification/Planting	
2	CC02042	Blue Gull Pond	Scarification/Planting	
2	CC02043	Portland	Scarification/Planting	
2	CC02044	Winterbook	Scarification/Planting	
2	CC02045	Bakeapple Bog	Scarification/Planting	
2	CC02046	Chance Harbour	Scarification/Planting	
2	CC02047	Charleston	Scarification/Planting	

2	CC02050	Trinity Pond	Scarification/Planting	
2	CC02051	Plate Cove Access Road	Scarification/Planting	
2	CC02052	Stock Cove Access Road	Scarification/Planting	
2	CC02054	World Pond	Scarification/Planting	
2	CC02056	Black Bay	Scarification/Planting	
2	CC02057	Terra Nova South	Scarification/Planting	
2	CC02058	Deep Bight	Scarification/Planting	
2	CC02059	Andrew's Pond	Scarification/Planting	
2	CC02060	Camp Pond	Scarification/Planting	

8.1.4 Primary Access Roads and Bridges

There are 105 km of new forest access roads scheduled to be constructed within the zone for the next five years to access timber for commercial purposes. All roads will be built to the specifications of the Class C, C-2 standards and all pertinent EPG's will be followed. In addition, secondary, operational and winter access roads and upgrading of existing road will be required and will be submitted in the annual operating plan prior to the year that they are planned to be built. As well, referrals will be sent to all relevant agencies (including DFO and Water Resources Division) before any construction is initiated.

Operating Area		Construction/	Length	Water Crossings	
Name	Harvest Block	Reconstruction	(km)	Culverts	Bridges
Goobies	CC02001				
Hatchet Cove	CC02002	С	1.5	8	
Burnt Point	CC02003				
Snook's Harbour	CC02004				
Foster's Marsh	CC02005	С	3	5	1
Hickman's Harbour Road	CC02006	С	0.85	3	
Northwest Brook	CC02008				
Beaver Pond	CC02009				
Clarenville	CC02010				
White Hills	CC02011	С	2.2	2	
Shoal Harbour Pond	CC02012	С	10.5	15	2
Thorburn Lake South	CC02013				
Southwest River Valley	CC02014	С	4.5	9	
Port Blandford Access Road	CC02015	С	1.2	1	

Table 13: FMD 02 Proposed Road Construction for 2022 to 2026.

Thorburn Lake North	CC02016				
George's Pond	CC02017	С	1	1	
Salt Lake	CC02018	С	1.2	4	
Shoal Harbour Valley	CC02019	С	1.2	2	
Muddy Hole Brook	CC02020	С	2.9	2	
George's Brook	CC02021	С	1.8	6	1
Barton	CC02022				
Harcourt Access Road	CC02023	С	4.3	6	
Jack's Pond	CC02024				
Nut Cove	CC02025				
Island Pond	CC02030				
Bloomfield Access Road	CC02031				
North Brook	CC02032	С	2.3	4	1
Branches Pond Access Road	CC02033	С	3	5	
New Country	CC02036				
Dalton's Pond	CC02037	С	3.6	5	
Rocky Pond	CC02038	R	4.7	9	
Rattle Falls	CC02040	С	1	1	
Princeton Pond	CC02041	С	2.2	4	1
Blue Gull Pond	CC02042				
Portland	CC02043	С	3.4	4	
Winterbrook	CC02044				
Bakeapple Bog	CC02045	С	4.8	3	
Chance Harbour	CC02046	С	2.8	4	
Charleston	CC02047	С	3.9	2	
Trinity Pond	CC02050				
Plate Cove Access Road	CC02051	С	18.4	21	1
Stock Cove Access Road	CC02052	С	5	9	
World Pond	CC02054	С	3.8	6	
Black Bay	CC02056	С	2.6	5	
Terra Nova South	CC02057				
Deep Bight	CC02058	С	1	3	
Andrew's Pond	CC02059	С	6.3	6	1
Camp Pond	CC02060				

8.1.5 Activities in Protected Water Supply Areas

Larger buffers are established inside PWSA and the pertinent EPG's will be attached to any commercial or domestic permits issued for these areas. There will be continuous monitoring inside these areas and buffers will be flagged to ensure compliance with the guidelines. All activity within a PPWSA must be approved by the Department of Environment.

SECTION 9 PLAN ADMINISTRATION

9.1 Monitoring

Monitoring of planned activities is critical to ensure objectives and operations are carried out in a manner consistent with various guidelines and provincial and federal legislation. Monitoring occurs at the operational level and the planning level. Forest harvesting activity is regulated using a permitting system and all activities are inspected and monitored on the ground by conservation officers to ensure compliance with applicable legislation, cutting permit conditions, and Environmental Protection Guidelines. Permit holders and contractors are also subject to financial penalties if deviations occur.

9.2 Amendments

Changes to an approved Operating Plan maybe required occasionally resulting from operational challenges or unforeseen circumstances. These changes are submitted as amendments and must be approved by the Forest Ecosystem Managment Division prior to implementation. There are two types of possible amendments:

- 1) Internally within the Department of Fisheries and Land Resources, where approval is required by the Forest Ecosystem Management Division. Internal amendments are governed by the following conditions:
 - a. within one kilometer of an operating area described in the five year operating plan, an additional area for timber harvesting that is, in total, not more than 50 hectares in each year of the plan
 - b. within a forest management district, an additional areas for silviculture treatment of not more that 20 percent of the total operating area described in the five year operating plan over the five year term of the plan
 - c. within an operating area described in the five year operating plan, not more than one kilometer, in total, of new primary forest access road in addition to existing and proposed primary forest access road in each year of the plan
 - d. adjacent to an operating area described in the five year operating plan, not more that half a kilometer, in total, of new primary forest access road in each year of that plan.
- 2) Externally, through the Department of Environment. Any required revisions which are not covered by the above internal requirements must be submitted for Environmental Assessment (EA) in the form of an amendment to the five year operating plan.