

# Real-Time Water Quality Deployment Report

## Vale - Long Harbour Annual Report

2023



Government of Newfoundland & Labrador  
Department of Environment and Climate Change  
Water Resources Management Division  
St. John's, NL, A1B 4J6 Canada

## Table of Contents

Introduction.....	1
Methods and Procedures.....	2
Results and Discussion.....	2
Surface Water Network.....	3
Water Temperature.....	3
pH.....	5
Specific Conductivity.....	6
Dissolved Oxygen.....	7
Turbidity.....	9
Groundwater Network.....	10
Water Temperature.....	10
pH.....	12
Specific Conductivity.....	14
Oxidation-Reduction Potential (ORP).....	16
Water Elevation.....	18
Path Forward.....	20
Appendix.....	21

## Introduction

Real-time water quality (RTWQ) monitoring of surface and groundwater quality on the Vale Long Harbour Processing plant site is carried out by the Department of Environment and Climate Change (ECC), Water Resources Management Division (WRMD). This work is undertaken in circumstances where industrial development has the potential to impact water bodies. The RTWQ program consists of more than 30 stations across the province from Voisey's Bay to St. Lawrence and Stephenville to St. John's.

RTWQ work in Long Harbour has been ongoing for more than 15 years – beginning with the first station, Rattling Brook below Bridge, in late 2006. In 2009, two additional surface water stations were deployed in the headwaters of Rattling Brook (Big Pond station) and lower in the river system (Rattling Brook below Plant Discharge). These surface water stations were positioned to monitor for long-term changes and water quality events related to the construction and operation of Vale's nickel processing plant.

As the nickel processing plant began to move towards operation, Sandy Pond was chosen as a residue storage area (RSA) to contain solid waste material. A groundwater monitoring network of five stations was deployed around the RSA in late 2012.

Surface and groundwater monitoring stations are depicted in Figure 1 below (blue triangles are surface water monitoring stations on Rattling Brook and yellow arrows are groundwater monitoring stations around the RSA).




Figure 1: Real-time water quality monitoring stations in Long Harbour, Newfoundland

## Methods and Procedures

Work under the RTWQ program is conducted according to the Protocols Manual for Real-Time Water Quality Monitoring in NL<sup>1</sup>. This document outlines the procedures, methods, and QAQC regimen used by all staff involved in the RTWQ program at all stations, province wide. For surface water monitoring, water quality instrumentation – in this case the Hydrolab DS5X multi-parameter sonde – is deployed on six-week intervals with *in situ* data validation at the beginning and end of deployment using an equivalent and freshly calibrated multi-parameter sonde. A grab sample is collected at the start of a deployment as an independent indicator of data quality.

Groundwater monitoring utilizes YSI EXO1 multi-parameter sondes. Due to the narrow confines of a 2” monitoring well, insertion of additional instruments into the well for verification purposes results in considerable changes to the well chemistry. As a result, data validation is restricted to capturing a grab sample immediately prior to insertion of newly-calibrated monitoring equipment in the well. Protocol requires a volume equivalent to three well casings to be purged from the well prior to sampling. This process flushes stagnant water from the well and ensures that the water being observed is aquifer water.

In the next section, long-term data from both the surface and groundwater monitoring networks are presented as line and boxplots. Guidelines set by the Canadian Council of Ministers of the Environment (CCME) and site-specific guidelines are indicated by dashed lines. Grab sample data for pH, specific conductivity, and turbidity is presented as black dots in the same figures. Boxplots are presented to illustrate how water quality parameters change from year to year.

Summary statistics and weather data are presented for each surface and groundwater parameter in the next section. Each table lists the 2023 median, minimum, and maximum values. *Average median* values for each parameter are calculated from the median values of each previous year and are provided in the same tables and labelled as *average* for simplicity. Median values are preferred throughout this report as a more robust indicator of central tendency than average values, especially given the highly skewed characteristic of environmental data.

## Results and Discussion

In the next sections, data from both surface and groundwater networks are presented as a series of line and box plots for water quality visualization over time and between stations. Summary statistics are presented in the appendix by year and station.

---

<sup>1</sup> <https://www.gov.nl.ca/ecc/files/waterres-rti-rtwq-nl-rtwq-manual.pdf>




## Surface Water Network

### Water Temperature

Water temperatures for all stations were lower than 2022. (Table 1).

**Table 1: Water temperatures at Rattling Brook**

Station	Segment	Median	Min	Max
Big Pond	2022	8.235	-0.04	24.39
	2023	7.71	0.08	23.6
Bridge	2022	9.27	-0.52	25.4
	2023	7.855	-0.19	26.6
Discharge	2022	9.29	-0.56	26.24
	2023	8.4	-0.4	27.51



**Figure 2: Water Temperature at Rattling Brook from 2021 to 2023**

Boxplots in Figure 3 illustrate the spread of data at each of the Rattling Brook stations.




Figure 3: Boxplots of water temperature at Rattling Brook from 2021 to 2023


## pH

In 2023, pH levels were above 2022 values at Bridge and Big Pond stations, but relatively similar between years at Plant Discharge station, according to median values (Table 2).

In 2023, median pH values were within site-specific guidelines<sup>2</sup> (dashed lines) at each station as shown in Figure 4.

**Table 2: pH at Rattling Brook**

Station	Segment	Median	Min	Max
Big Pond	2022	6.43	4.4	6.89
	2023	6.52	5.13	6.79
Bridge	2022	6.48	5.56	7.18
	2023	6.52	5.85	6.94
Discharge	2022	6.47	6.02	6.8
	2023	6.47	5.65	7.06



**Figure 4: pH at Rattling Brook from 2021 to 2023**


### Specific Conductivity

In 2023, specific conductivity decreased at all three stations when compared to 2022 data, according to Figure 5 and Table 3.

Big Pond has been experiencing more fluctuations in conductivity since the water level rose in 2017. Conductivity may remain elevated until soil stabilizes.

**Table 3: Specific Conductivity at Rattling Brook**

Station	Segment	Median	Min	Max
Big Pond	2022	62.7	30.9	99.8
	2023	60	46.9	95.1
Bridge	2022	54.6	20.8	157.8
	2023	52	27.9	77.4
Discharge	2022	56.1	33.9	192.7
	2023	55.7	17	118.9



**Figure 5: Specific Conductivity at Rattling Brook from 2021 to 2023**




### Dissolved Oxygen

In 2023, dissolved oxygen levels were within normal ranges for all stations. All values at the stations were found to be above the minimum guideline for other life stages (min values in Table 4). Big Pond experiences lower dissolved oxygen due to warmer temperatures as the large waterbody holds heat longer.

Two CCME minimum guidelines shown as dashed lines give a conservative value of 9.5 mg/l for early life stage organisms and another value of 6.5 mg/l for other life stages.

**Table 4: Dissolved Oxygen at Rattling Brook**

Station	Segment	Median	Min	Max
Big Pond	2022	11.37	7.59	14.44
	2023	11.23	7.43	14.7
Bridge	2022	10.99	7.9	14.76
	2023	11.22	7.57	14.85
Discharge	2022	10.71	7.6	14.39
	2023	11.41	7.15	14.84



**Figure 6: Dissolved oxygen at Rattling Brook from 2021 to 2023**

Figure 7 shows the range of dissolved oxygen concentrations at each station for the past three years.

In 2023, the medians at all stations were very similar.




Figure 7: Boxplots of dissolved oxygen at Rattling Brook from 2021 to 2023


### Turbidity

In 2023, median turbidity level at Big Pond was 0.0 NTU – equal to the long-term average. Below Bridge and Plant Discharge saw a slight increase in Median values (Table 5).

Maximum turbidity level at Big Pond was higher in 2023 than 2022, while Below Bridge and Plant Discharge were lower.

**Table 5: Turbidity at Rattling Brook**

Station	Segment	Median	Min	Max
Big Pond	2022	0.6	0.0	253.1
	2023	0	0	273.9
Bridge	2022	0.0	0.0	313.7
	2023	1.8	0	269.2
Discharge	2022	0.0	0.0	220.7
	2023	0.3	0	193.5



**Figure 8: Turbidity at Rattling Brook from 2021 to 2023**

## Groundwater Network


### Water Temperature

As shown in Table 6, in 2023, median water temperature increased slightly from 2022 levels: within 0.08°C at well 1 Deep, 0.14°C at well 2 Deep, 0.05°C at well 2 Shallow, 0.01°C at well 3 Deep and 0.15°C at well 4 Deep (Table 6).

The narrow range and stability of water temperatures is reinforced by the scale of the y-axes in Figure 9.

**Table 6: Temperature at Residue Storage Area**

Station	Segment	Median	Min	Max
1 Deep	2022	6.95	6.83	7.51
	2023	7.03	6.94	7.16
2 Shallow	2022	6.69	3.45	11.39
	2023	6.74	3.53	11.14
2 Deep	2022	6.53	6.13	6.97
	2023	6.67	6.25	7.04
3 Deep	2022	7.02	6.9	7.35
	2023	7.1	6.98	7.46
4 Deep	2022	6.74	6.64	6.83
	2023	6.89	6.79	6.99



**Figure 9: Water temperature at the Residue Storage Area from 2021 to 2023**

Within each well, water temperature ranges were similar in 2023 compared to the previous year (Figure 10).




Figure 10: Boxplots of water temperature at the Residue Storage Area from 2021 to 2023



pH

2023 Median pH values were found to be higher than 2022 levels at all stations with exception of well 4 Deep (Table 7).

Long term trends are difficult to extrapolate from Figure 11, but the gray trend lines show pH plateaus at well 1 Deep, 2 Shallow, and 3 Deep for 2023. Well 4 deep had a decreasing trend, while 2 Deep showed no trend. Well station equipment was upgraded late 2020 - early 2021.

Table 7: pH at Residue Storage Area

Station	Segment	Median	Min	Max
1 Deep	2022	5.48	5.27	5.86
	2023	5.53	5.3	6.16
2 Shallow	2022	5.47	4.81	5.75
	2023	5.68	5.22	6.03
2 Deep	2022	6.44	5.7	7.35
	2023	6.87	5.57	7.69
3 Deep	2022	5.9	5.7	6.0
	2023	5.98	5.74	6.11
4 Deep	2022	7.07	6.81	7.82
	2023	6.81	6.04	7.6




Figure 11: pH at the Residue Storage Area from 2021 to 2023

Figure 12 shows the range of pH at each station for the past three years.




Figure 12: Boxplots of pH at the Residue Storage Area from 2021 to 2023

### Specific Conductivity


In 2023, median specific conductivity values were below 2022 values.

For 2023, figure 13 shows variable trends at all stations throughout the year, with the exception of 4 deep which shows a noticeable decreasing trend.

Figure 13 also shows grab samples taken during routine maintenance (black circles).

**Table 8: Specific conductivity at Residue Storage Area**

Station	Segment	Median	Min	Max
1 Deep	2022	176	150	207
	2023	162	116	213
2 Shallow	2022	335	162	481
	2023	266	170	372
2 Deep	2022	338	224	545
	2023	260	214	549
3 Deep	2022	129	110	146
	2023	128	118	134
4 Deep	2022	236	196	304
	2023	197	115	279



**Figure 13: Specific conductivity at the Residue Storage Area from 2021 to 2023**

From 2021-2023 (Figure 14), specific conductivity appears to be trending downward overall at well 1 deep, 3 deep and 4 deep. Well 2 shallow and 2 deep are variable and show no significant trends over the past 3 years.




Figure 14: Boxplots of specific conductivity at the Residue Storage Area from 2021 to 2023


### Oxidation-Reduction Potential (ORP)

Due to the high variability in ORP values following equipment maintenance, raw ORP values can be challenging to observe. As such, the gray trend lines in Figure 15 give a more intuitive indication of ORP tendency over time.

Table 9 shows that median ORP values from 2023 are greater than 2022 at well 2 Shallow while all other stations recorded lower median values in 2023.

**Table 9: ORP at Residue Storage Area**

Station	Segment	Median	Min	Max
1 Deep	2022	432.39	157.98	452.64
	2023	378.29	59.37	470.28
2 Shallow	2022	70.55	-258.71	412.54
	2023	104.76	-356.83	382.78
2 Deep	2022	-280.68	-633.11	111.5
	2023	-379.88	-633.36	115.39
3 Deep	2022	390.4	212.57	453.35
	2023	377.33	211.45	448.1
4 Deep	2022	486.45	-6.53	550.4
	2023	474.93	-86.45	553.68



**Figure 15: Oxidation-Reduction Potential at the Residue Storage Area from 2021 to 2023**



Figure 16 shows the range of ORP at each station for the past three years.

Year-over-year tendency in ORP values are most clearly shown in Figure 16. Wells 2 deep and 4 deep show the greatest degree of change over the previous three years.




Figure 16: Boxplots of Oxidation-Reduction Potential at the Residue Storage Area from 2021 to 2023


### Water Elevation

The local height of the aquifer surrounding each well is closely indicated by the water level in each well. These values are not expected to change a great deal over the long term, barring unforeseen circumstances or major changes to water level in nearby water bodies.

Periodic variation is commonplace as illustrated by Figure 17, but levels in 2023 were close to previous years as shown in Table 10. During dam construction in 2023 Well 1 experienced a longer recharge of elevation. An equipment upgrade in Well 2 Shallow led to a noticeable adjustment in elevation in early 2021.

**Table 10: Water level at Residue Storage Area**

Station	Segment	Median	Min	Max
1 Deep	2022	131.9716	131.4263	132.7686
	2023	131.5441	127.9078	132.6498
2 Shallow	2022	114.1629	113.661	115.0802
	2023	114.2904	113.6155	115.1724
2 Deep	2022	113.2821	112.6758	114.0687
	2023	113.1770	112.6446	114.1597
3 Deep	2022	133.3953	131.511	136.5113
	2023	133.6958	132.3863	135.78
4 Deep	2022	138.0313	137.0165	138.9141
	2023	137.974	136.0023	139.112



**Figure 17: Water elevation at the Residue Storage Area from 2021 to 2023**

Figure 18 shows the elevation of water levels in each well from 2021 to 2023. Values are largely stable at all stations with only slight variations over the previous three years.




Figure 18: Boxplots of water elevation at the Residue Storage Area from 2021 to 2023

## **Path Forward**

A 1.5 m water level increase at Rattling Brook Big Pond in November 2017 saw the inundation of shoreline vegetation and soils. As the organic matter decays, changes in water quality are expected over the short- to mid-term. In particular, dissolved oxygen concentrations may fall below guidelines set by the CCME during warm water conditions.

Station maintenance and calibration activities will continue to take place every six weeks at surface water stations while maintenance and calibration activities will occur four times per year at groundwater stations.

Reports on surface water stations will be generated at the end of each deployment period while a summary of the groundwater data will be included in the annual report, due to the relatively static nature of data from monitoring wells.

In partnership with ECC, a weather station was installed on site at Vale Long Harbour in December 2020. A wind event damaged the weather station in January of 2022. There is a plan to repair and have the station operational again by the end of 2023. The additional weather station in the Long Harbour community was completed in February 2023.

In 2021 the ground water equipment in the Residue Storage Area was upgraded.

Equipment upgrade for the surface water stations at Rattling Brook is scheduled for completion 2024-25.

Ongoing and cooperative efforts between the department of Environment and Climate Change and Vale to monitor water quality on a real-time basis have been successful in identifying areas worthy of attention.

## Appendix

**Table 1: Summary statistics of Water Temperature from 2008 to 2023**

Station	Year	Mean	Median	Min	Max
Big Pond	2009	5.13	5.15	0.2	12.26
	2010	10.08	9.65	0.04	22.4
	2011	9.58	9.875	-0.02	20.88
	2012	10.00	11.28	0	22.87
	2013	9.67	10.04	-0.02	22.17
	2014	10.58	11.37	0.01	23.1
	2015	10.11	11.68	-0.39	21.46
	2016	10.87	11.52	-0.44	22.24
	2017	11.24	11.96	-0.47	20.61
	2018	8.07	8.44	-0.43	20.87
	2019	8.24	6.76	0.23	22.26
	2020	11.90	12.72	0.25	21.77
	2021	8.92	8.16	0	20.96
	2022	9.22	8.24	-0.04	24.39
2023	9.19	7.71	0.08	23.6	
Below Bridge	2008	6.73	6.195	-0.42	21.93
	2009	9.14	8.025	-0.5	23.97
	2010	8.65	7.73	-0.5	22.84
	2011	7.70	6.43	-0.48	22.2
	2012	9.52	9.77	-0.51	23.82
	2013	9.03	9.16	-0.49	24.98
	2014	8.65	7.455	-0.5	24.93
	2015	7.91	6.69	-0.03	22.69
	2016	9.10	9.15	-0.54	24.69
	2017	8.30	7.25	-0.54	23.67
	2018	8.49	8.65	-0.41	23.98
2019	8.65	7.39	-0.05	24.42	
2020	8.85	6.83	-0.56	25.23	

Plant Discharge	2021	9.23	8.53	-0.5	24.29
	2022	9.52	9.27	-0.52	25.4
	2023	9.46	7.86	-0.19	26.6
	2009	4.02	4.2	0.02	11.37
	2010	9.04	8.12	0.02	23.67
	2011	8.43	7.49	-0.07	22.89
	2012	9.98	10.16	-0.03	24.33
	2013	10.05	10.9	-0.03	24.7
	2014	9.27	9.36	0	25.48
	2015	8.05	6.705	-0.51	23.25
	2016	9.10	9.22	-0.55	25
2017	8.49	7.74	-0.52	24.13	
2018	8.21	7.88	-0.53	24.77	
2019	8.66	7.705	-0.07	24.67	
2020	9.36	7.63	-0.04	25.8	
2021	9.23	8.37	-0.49	25.02	
2022	9.59	9.29	-0.56	26.24	
2023	9.49	8.4	-0.4	27.51	

**Table 2: Summary statistics of pH from 2008 to 2023**

Station	Year	Mean	Median	Min	Max
Big Pond	2009	6.24	6.24	5.86	6.41
	2010	6.22	6.25	5.34	6.8
	2011	6.29	6.32	5.45	6.74
	2012	6.48	6.51	5.37	7.14
	2013	6.41	6.42	5.02	7.51
	2014	6.43	6.46	5.65	6.78
	2015	6.58	6.59	5.57	7.07
	2016	6.49	6.54	5.23	8.74
	2017	6.43	6.41	5.54	7.16
	2018	5.84	5.95	4.93	6.36



Appendix – Weather

	2019	5.81	5.98	4.49	6.89
	2020	6.10	6.13	4.81	6.38
	2021	6.15	6.24	4.72	6.83
	2022	6.16	6.43	4.4	6.89
	2023	6.41	6.52	4.13	6.79
Below Bridge	2008	6.08	6.11	5.42	6.5
	2009	5.98	5.99	5.25	6.71
	2010	6.19	6.24	5.22	6.81
	2011	6.16	6.19	5.41	6.81
	2012	6.29	6.29	5.15	7
	2013	6.14	6.21	4.89	6.94
	2014	6.09	6.09	5.13	7.1
	2015	6.34	6.37	5.45	6.94
	2016	6.21	6.39	4.84	7
	2017	6.31	6.38	5.15	7.2
	2018	6.08	6.25	4.74	6.92
	2019	6.21	6.26	4.72	7.05
	2020	6.21	6.28	4.9	6.91
2021	6.39	6.44	5.34	6.79	
2022	6.44	6.48	5.56	7.18	
2023	6.48	6.52	5.85	6.94	
Plant Discharge	2009	6.30	6.29	5.82	6.78
	2010	6.45	6.44	5.12	6.95
	2011	6.61	6.57	6.07	7.67
	2012	6.58	6.58	5.92	7.48
	2013	6.54	6.6	5.45	7.12
	2014	6.62	6.63	4.83	7.17
	2015	6.66	6.66	6.37	6.96
	2016	6.46	6.57	5.17	7.03
	2017	6.68	6.6	5.62	7.53
	2018	6.08	6.22	4.59	6.84
	2019	6.27	6.39	5.19	6.97
2020	6.48	6.44	5.77	7.41	

2021	6.43	6.45	5.48	6.88
2022	6.46	6.47	6.02	6.8
2023	6.44	6.47	5.65	7.06

**Table 3: Summary statistics of Conductivity from 2008 to 2023**

Station	Year	Mean	Median	Min	Max
Big Pond	2009	33.05	33.2	29.6	35.4
	2010	35.18	35.6	27.4	55.7
	2011	43.38	44.6	33.1	57
	2012	52.99	52.8	28.2	73.8
	2013	54.82	56.5	32.5	77.4
	2014	58.35	58.8	30.6	68.1
	2015	60.56	60.8	39.1	70.3
	2016	62.13	62.4	37.6	76.3
	2017	67.23	68.3	45.8	110.9
	2018	76.12	76.6	54.2	135.8
	2019	73.73	73.8	56.8	118.3
	2020	65.25	64.3	46.2	125.3
	2021	63.12	63.3	41.9	108.5
2022	62.58	62.7	30.9	99.8	
2023	60.55	60	46.9	95.1	
Below Bridge	2008	32.17	31.8	21.6	44.4
	2009	36.91	36.5	27.5	51.6
	2010	38.06	38	27.4	83.6
	2011	40.80	40.6	21.2	87.1
	2012	52.87	50.1	20.2	81.1
	2013	55.11	53.9	29.3	116.6
	2014	56.08	57	20.3	120.7
	2015	59.01	58.3	50.6	82.6
	2016	59.15	58.7	47.3	119.1
	2017	60.79	61.9	38.8	94.1
	2018	66.42	66.6	37.3	96.2
2019	62.88	62.7	37.6	162.2	
2020	53.61	53.7	33.6	91.4	

Appendix – Weather

Plant Discharge	2021	54.88	55.5	33.1	140.1
	2022	54.16	54.6	20.8	157.8
	2023	51.85	52	27.9	77.4
	2009	36.24	35.5	30.6	60
	2010	46.48	44.9	35.5	99.8
	2011	53.36	51.9	36.5	147.9
	2012	69.13	64.7	45.5	202
	2013	75.81	72.5	51	158.7
	2014	72.45	70.4	43.9	161.4
	2015	74.03	73	52.3	121
	2016	72.40	71	54.3	203
	2017	66.89	65.9	45.4	275
	2018	71.50	70.9	3.7	144.7
	2019	69.55	68	57.5	222
	2020	58.17	56.7	45.9	188.1
	2021	58.25	57.9	47	168.3
	2022	56.61	56.1	33.9	192.7
	2023	55.9	55.7	17	118.9

**Table 4: Summary statistics of Total Dissolved Solids from 2008 to 2023**

Station	Year	Mean	Median	Min	Max
Big Pond	2009	0.02	0.0213	0.0189	0.0226
	2010	0.02	0.0228	0.0175	0.0357
	2011	0.03	0.0285	0.0212	0.0365
	2012	0.03	0.0338	0.018	0.0473
	2013	0.03	0.0355	0.0208	0.0495
	2014	0.04	0.0376	0.0196	0.0436
	2015	0.04	0.0389	0.025	0.045
	2016	0.04	0.0399	0.022	0.0489
	2017	0.04	0.0437	0.022	0.071
	2018	0.05	0.049	0.0347	0.0869
	2019	0.05	0.0472	0.0364	0.0757
	2020	0.04	0.0412	0.0296	0.0802
	2021	0.04	0.0405	0.0268	0.0694

Below Bridge	2022	0.04	0.0401	0.0198	0.0639	
	2023	0.04	0.04	0.03	0.0609	
	2008	0.02	0.0204	0.0138	0.0284	
	2009	0.02	0.0234	0.0176	0.033	
	2010	0.02	0.0243	0.0176	0.0535	
	2011	0.03	0.026	0.0136	0.0557	
	2012	0.03	0.0321	0.0129	0.0519	
	2013	0.04	0.0345	0.0187	0.0746	
	2014	0.04	0.0365	0.013	0.0773	
	2015	0.04	0.0373	0.0324	0.0528	
	2016	0.04	0.0375	0.0303	0.0762	
	2017	0.04	0.0396	0.0249	0.0602	
	2018	0.04	0.0426	0.0239	0.0616	
	2019	0.04	0.0401	0.024	0.1038	
	2020	0.03	0.0344	0.0215	0.0585	
	2021	0.04	0.0355	0.0212	0.0896	
	2022	0.03	0.035	0.0133	0.101	
	2023	0.3	0.0333	0.0179	0.0496	
	Plant Discharge	2009	0.02	0.0227	0.0196	0.0384
		2010	0.03	0.0287	0.0227	0.0639
2011		0.03	0.0332	0.0234	0.0946	
2012		0.04	0.0414	0.0291	0.129	
2013		0.05	0.0467	0.0329	0.1016	
2014		0.05	0.0451	0.0281	0.1033	
2015		0.05	0.0467	0.0335	0.0774	
2016		0.05	0.0455	0.0347	0.13	
2017		0.04	0.0422	0.0291	0.176	
2018		0.05	0.0454	0.0023	0.0926	
2019		0.04	0.0435	0.0368	0.142	
2020		0.04	0.0363	0.0294	0.1204	
2021		0.04	0.0371	0.0301	0.1077	
2022	0.04	0.0359	0.0217	0.1233		
2023	0.04	0.0357	0.0109	0.0761		

**Table 5: Summary statistics of Dissolved Oxygen from 2008 to 2023**

Station	Year	Mean	Median	Min	Max
Big Pond	2009	11.56	11.72	9.42	12.88
	2010	10.68	10.69	8.06	13.53
	2011	10.99	10.71	8.39	14.42
	2012	10.86	10.47	8.17	14.69
	2013	10.74	10.55	8.29	14.43
	2014	10.80	10.36	8.27	13.27
	2015	10.90	10.26	8.68	14.54
	2016	10.74	10.42	8.62	15.93
	2017	10.59	10.35	8.41	13.68
	2018	10.36	10.67	4.6	13.5
	2019	10.63	10.85	2.61	13.23
	2020	10.02	9.83	7.26	13.1
	2021	10.84	10.97	6.98	13.68
	2022	10.99	11.37	7.59	14.44
2023	10.98	11.23	7.43	14.7	
Below Bridge	2008	12.06	12.15	8.35	14.63
	2009	11.30	11.26	7.72	14.61
	2010	11.43	11.36	7.81	14.9
	2011	11.74	11.7	8.08	15.11
	2012	11.32	10.95	7.54	15.51
	2013	11.17	11.04	7.65	14.21
	2014	11.41	11.53	7.86	14.4
	2015	11.70	11.82	8.34	14.68
	2016	11.38	11.15	7.61	14.53
	2017	11.61	11.64	7.98	14.69
	2018	11.48	11.28	7.64	14.47
	2019	11.48	11.44	8	14.73
	2020	11.49	11.7	8.02	14.48
	2021	11.19	10.99	8.04	14.48
2022	11.32	10.99	7.9	14.76	
2023	11.17	11.22	7.57	14.85	

Plant Discharge	2009	12.25	12.28	10.29	14.1
	2010	10.94	10.95	7.02	14.48
	2011	11.24	10.99	7.12	14.76
	2012	10.91	10.66	6.46	14.45
	2013	10.96	10.52	7.28	14.2
	2014	11.09	10.95	7.39	14.3
	2015	11.55	11.79	7.59	14.68
	2016	11.23	11.1	7.18	14.57
	2017	11.53	11.5	7.5	14.91
	2018	11.45	11.35	7.03	14.73
	2019	11.30	11.185	6.88	14.44
	2020	11.18	11.37	7.56	14.13
	2021	11.10	11.09	7.68	14.51
	2022	11.07	10.71	7.6	14.39
2023	11.16	11.41	7.15	14.84	

**Table 6: Summary statistics of Percent Saturation from 2008 to 2023**

Station	Year	Mean	Median	Min	Max
Big Pond	2009	92.14	91.8	86.8	98.9
	2010	94.06	94	85.9	104.4
	2011	94.67	94.2	87.3	109.3
	2012	93.94	93.1	84.8	109.6
	2013	92.31	91.9	81.4	109.1
	2014	95.00	95	82.7	106.4
	2015	95.10	95.1	86.8	106.2
	2016	95.00	94.5	84.8	106.8
	2017	95.07	95.7	82.5	104.8
	2018	85.82	87.2	46.9	99.9
	2019	89.99	91.2	26.7	108.6
	2020	91.71	92	77.8	102.2
	2021	92.75	92.7	71.5	107.2
	2022	93.77	93.5	82	104.4
2023	94.40	94.6	82.2	107.0	
Below Bridge	2008	96.07	96.3	89.8	100.6

Appendix – Weather

	2009	96.09	96.3	88.3	102.6	
	2010	96.65	96.7	88.9	103.3	
	2011	95.93	96.05	88.2	104.4	
	2012	96.04	96.5	85.6	105.3	
	2013	94.36	94.3	88.7	101.6	
	2014	94.80	94.9	87.7	101.5	
	2015	96.91	97	89.5	103	
	2016	96.72	96.8	88.9	103.5	
	2017	96.38	96.7	89	102.5	
	2018	95.99	96.2	88.9	102.9	
	2019	97.17	96.9	91.2	104	
	2020	96.75	96.2	88.7	108.9	
	2021	95.92	95.8	89.4	102.5	
	2022	97.23	97.1	89.8	106.3	
	2023	96.15	96.4	88.1	103.9	
	Plant Discharge	2009	94.94	95.5	88.1	101.6
		2010	93.63	94.1	80.1	105.7
		2011	94.33	95.1	72.5	103
		2012	93.33	94.8	70.8	103.4
		2013	94.37	94.3	83.1	103.6
		2014	93.47	94.1	84	101
		2015	95.47	95.9	84.7	103.1
		2016	95.32	95.9	83.5	103.6
2017		95.48	95.6	85.9	102.8	
2018		94.75	95.6	80.9	103.7	
2019		95.37	95.1	70.7	108	
2020		95.07	95.2	85.2	102.3	
2021		94.91	95	86.7	103	
2022	95.33	95.2	88.3	105.1		
2023	96.02	96.3	87.1	104.7		

**Table 7: Summary statistics of Turbidity from 2008 to 2023**

Station	Year	Mean	Median	Min	Max
Big Pond	2009	3.37	1.7	0	22

	2010	2.36	0	0	116.6	
	2011	0.63	0	0	44.9	
	2012	0.20	0	0	22	
	2013	0.05	0	0	84.8	
	2014	0.03	0	0	81.1	
	2015	0.29	0	0	25.3	
	2016	0.48	0	0	15	
	2017	0.10	0	0	77	
	2018	1.19	0	0	672	
	2019	2.53	0	0	555	
	2020	0.96	0.2	0	114.5	
	2021	0.95	0	0	215.4	
	2022	2.59	0.6	0	253.1	
	2023	2.72	0	0	273.9	
	Below Bridge	2008	0.61	0	0	963
		2009	10.37	0	0	1486
		2010	10.24	2.5	0	445
		2011	6.00	0.4	0	2259
		2012	22.64	3.35	0	1437
		2013	6.42	2.4	0	998
		2014	2.31	0	0	886
		2015	2.91	0	0	396.9
		2016	5.42	0	0	781
2017		1.80	0	0	371.7	
2018		1.91	0	0	353.5	
2019		3.77	0.3	0	460	
2020		2.39	0	0	251.6	
2021	3.77	0	0	784		
2022	0.9	0	0	313.7		
2023	3.77	1.8	0	269.2		
Plant Discharge	2009	67.35	23.6	4.3	1094	
	2010	11.48	3.3	0	460	
	2011	6.69	1.7	0	734	

Appendix – Weather

2012	19.41	4.8	0	586
2013	11.10	4.5	0	580
2014	2.57	0	0	277.2
2015	2.51	0	0	282.5
2016	7.66	0.6	0	314.6
2017	1.79	0	0	430
2018	1.93	0	0	548
2019	2.42	0.5	0	485
2020	1.83	0.1	0	415
2021	1.82	0.40	0	243.8
2022	2.96	0	0	220.7
2023	2.12	0.3	0.0	193.5

**Table 8: Summary statistics of Water Temperature at the Residue Storage Area from 2013 to 2023**

Station	Year	Mean	Median	Min	Max
Well 1 Deep	2012	7.40	7.43	7.2	7.49
	2013	6.78	6.87	6.3	7.45
	2014	6.69	6.68	6.59	6.97
	2015	6.63	6.63	6.56	6.7
	2016	6.62	6.62	6.59	6.68
	2017	6.64	6.64	6.58	6.71
	2018	6.68	6.67	6.64	6.73
	2019	6.70	6.72	6.51	7.02
	2020	6.74	6.74	6.54	7.08
	2021	6.82	6.83	6.64	7.06
	2022	6.92	6.95	6.83	7.51
2023	7.03	7.03	6.94	7.16	
Well 2 Shallow	2012	6.17	6.27	4.23	7.6
	2013	6.83	6.71	2.22	11.81
	2014	6.77	6.53	1.03	11.81
	2015	6.15	5.8	2.91	10.5
	2016	6.21	6.1	1.68	11.07
	2017	5.96	5.61	2.44	9.95
2018	6.38	5.94	3.21	11.27	
2019	5.90	5.65	2.5	10.56	
2020	6.44	5.925	3.18	11.22	

Well 2 Deep	2021	7.54	7.66	3.79	11.62
	2022	7.13	6.69	3.45	11.39
	2023	7.04	6.74	3.53	11.14
	2013	6.53	6.53	6.17	6.91
	2014	6.35	6.32	5.77	6.88
	2015	6.35	6.35	5.95	6.81
	2016	6.20	6.19	5.78	6.59
	2017	6.15	6.13	5.74	6.58
	2018	6.14	6.16	5.79	6.46
	2019	6.03	6.02	5.45	6.51
	2020	6.02	6.04	5.64	6.47
2021	6.35	6.37	6.03	6.94	
2022	6.51	6.53	6.13	6.97	
2023	6.66	6.67	6.25	7.04	
Well 3 Deep	2012	6.44	6.44	6.41	6.47
	2013	6.62	6.63	6.46	6.77
	2014	6.75	6.76	6.5	7.01
	2015	6.69	6.715	6.52	6.89
	2016	6.62	6.63	6.52	6.87
	2017	6.67	6.68	6.53	6.9
	2018	6.70	6.69	6.61	6.93
	2019	6.69	6.75	6.4	7
	2020	6.59	6.57	6.44	6.82
	2021	6.84	6.83	6.67	7.17
	2022	7.02	7.02	6.9	7.35
2023	7.14	7.1	6.98	7.46	
Well 4 Deep	2012	6.24	6.24	6.18	6.29
	2013	6.40	6.39	6.28	6.53
	2014	6.46	6.46	6.32	6.6
	2015	6.46	6.48	6.32	6.64
	2016	6.39	6.39	6.3	6.49
	2017	6.39	6.4	6.28	6.52
	2018	6.40	6.42	6.27	6.51
	2019	6.44	6.47	6.29	6.56
	2020	6.22	6.41	5.76	6.5
	2021	6.57	6.57	6.43	6.96
	2022	6.75	6.74	6.64	6.83
2023	6.89	6.89	6.79	6.99	

**Table 9: Summary statistics of pH at the Residue Storage Area from 2013 to 2023**

Station	Year	Mean	Median	Min	Max
Well 1 Deep	2012	5.90	5.89	5.81	6.02
	2013	5.75	5.76	5.51	5.92
	2014	5.68	5.72	5.15	5.88
	2015	6.46	6.67	5.15	7.43
	2016	7.55	7.57	6.51	7.77
	2017	7.64	7.72	7.37	7.9
	2018	7.82	7.85	7.66	7.94
	2019	7.06	7.64	5.95	8.23
	2020	5.98	5.95	5.46	6.41
	2021	5.64	5.59	5.32	7.23
	2022	5.48	5.48	5.27	5.86
	2023	5.55	5.53	5.3	6.16
Well 2 Shallow	2012	5.39	5.32	4.97	5.85
	2013	5.64	5.72	4.89	5.98
	2014	5.70	5.74	4.65	6.05
	2015	5.75	5.73	4.52	6.12
	2016	6.11	6.11	5.61	6.68
	2017	6.00	5.95	5.42	7.13
	2018	6.18	6.13	5.73	6.87
	2019	6.16	6.17	5.36	6.9
	2020	6.45	6.57	5.6	6.96
	2021	5.72	5.59	5.02	6.92
	2022	5.47	5.47	4.81	5.75
2023	5.68	5.68	5.22	6.03	
Well 2 Deep	2013	8.13	8.16	6.98	8.35
	2014	8.08	8.12	7.38	8.2
	2015	8.27	8.25	7.24	8.57
	2016	8.48	8.55	6.65	8.75
	2017	8.32	8.41	6.4	8.56
	2018	8.24	8.18	6.61	8.74
	2019	8.45	8.53	6.66	9.21
	2020	8.22	7.83	6.67	9.35
	2021	7.05	6.98	5.56	7.89
	2022	6.57	6.44	5.7	7.35
2023	6.90	6.87	5.57	7.69	
Well 3 Deep	2012	5.66	5.62	5.33	5.96

	2013	5.75	5.73	5.08	6.03
	2014	5.69	5.73	5.06	6.05
	2015	5.67	5.63	5.11	6.03
	2016	5.93	5.95	5.73	6.23
	2017	5.99	5.98	5.8	6.25
	2018	5.89	5.9	5.55	6.16
	2019	6.07	6.02	4.49	6.48
	2020	5.94	5.87	5.6	6.28
	2021	5.84	5.84	5.24	6.06
	2022	5.9	5.9	5.7	6.0
	2023	5.98	5.98	5.74	6.11
Well 4 Deep	2012	7.77	7.76	7.71	7.98
	2013	7.03	6.77	6.43	8.08
	2014	7.25	7.14	6.71	7.85
	2015	7.97	8.11	6.31	8.27
	2016	8.04	8.06	7.54	8.45
	2017	8.29	8.27	8.03	8.68
	2018	8.46	8.37	7.97	8.88
	2019	8.08	8.4	6.1	9.05
	2020	8.06	8	7.53	8.6
	2021	7.55	7.62	6.91	8.06
	2022	7.15	7.07	6.81	7.82
2023	6.76	6.81	6.04	7.6	

**Table 10: Summary statistics of conductivity at the Residue Storage Area from 2013 to 2023**

Station	Year	Mean	Median	Min	Max
Well 1 Deep	2012	125.97	126	118	136
	2013	135.18	136	116	161
	2014	149.98	149	129	179
	2015	233.45	253	151	273
	2016	278.12	282	234	296
	2017	393.89	391	291	503
	2018	559.55	566	470	642
	2019	464.76	636	196	693
	2020	239.26	231	168	306
	2021	187.20	186	158	236
	2022	175.11	176	150	207
	2023	163.95	162.28	116.35	213.26

Appendix – Weather

Well 2 Shallow	2012	87.69	87	76	123
	2013	84.76	85	62	131
	2014	96.49	94	72	154
	2015	97.46	96	83	120
	2016	127.69	115	92	208
	2017	127.23	126	90	206
	2018	138.43	137	104	194
	2019	166.57	165	124	225
	2020	194.06	195	131	236
	2021	242.33	227	118	397
2022	332.96	335	162	481	
2023	266.70	266	170	372	
Well 2 Deep	2013	227.55	228	179	233
	2014	226.41	226	209	236
	2015	222.67	224	212	233
	2016	213.16	213	141	220
	2017	219.07	219	159	233
	2018	217.58	218	198	229
	2019	222.41	220	207	283
	2020	268.80	251	218	467
	2021	296.94	294	220	454
	2022	364	338	224	545
2023	295.38	260	214	549	
Well 3 Deep	2012	111.26	106	96	141
	2013	104.79	102	84	144
	2014	115.43	116	88	157
	2015	128.63	127	104	163
	2016	133.51	132	104	182
	2017	123.96	121	81	178
	2018	129.30	128	99	173
	2019	124.12	126	84	159
	2020	132.54	132	103	166
	2021	132.59	134	97	153
2022	129.84	129	110	146	
2023	127.30	128	118	134	
Well 4 Deep	2012	313.74	314	299	316
	2013	206.24	173	135	315
	2014	237.23	234	152	297
	2015	231.61	227	178	283

2016	263.85	268	226	283
2017	271.08	268	261	298
2018	265.54	269	245	284
2019	274.87	274	267	287
2020	266.99	276	195	288
2021	275.77	280	202	308
2022	241.9	236	196	304
2023	185.15	197	115	279

**Table 11: Summary statistics of Total Dissolved Solids at the Residue Storage Area from 2013 to 2023**

Station	Year	Mean	Median	Min	Max
Well 1 Deep	2012	0.08	0.082	0.077	0.088
	2013	0.09	0.088	0.075	0.105
	2014	0.10	0.097	0.084	0.116
	2015	0.15	0.165	0.098	0.178
	2016	0.18	0.183	0.152	0.192
	2017	0.26	0.254	0.189	0.327
	2018	0.36	0.368	0.305	0.417
	2019	0.30	0.413	0.127	0.451
	2020	0.16	0.15	0.109	0.199
	2021	0.12	0.12	0.1	0.15
2022	0.11	0.11	0.1	0.13	
2023	0.11	0.11	0.08	0.14	
Well 2 Shallow	2012	0.06	0.056	0.05	0.08
	2013	0.06	0.055	0.04	0.085
	2014	0.06	0.061	0.047	0.1
	2015	0.06	0.062	0.054	0.078
	2016	0.08	0.075	0.06	0.135
	2017	0.08	0.082	0.059	0.134
	2018	0.09	0.089	0.068	0.126
	2019	0.11	0.107	0.081	0.146
	2020	0.13	0.127	0.085	0.154
	2021	0.16	0.15	0.08	0.26
2022	0.22	0.22	0.11	0.310.24	
2023	0.17	0.17	0.11	0.24	
Well 2 Deep	2013	0.15	0.148	0.116	0.151
	2014	0.15	0.147	0.136	0.153
	2015	0.14	0.145	0.138	0.151
	2016	0.14	0.139	0.092	0.143



Appendix – Weather

	2017	0.14	0.142	0.104	0.151		2017	-224.67	-193.55	-441.4	84.4
	2018	0.14	0.142	0.13	0.149		2018	-57.77	-47.4	-201.6	189.2
	2019	0.14	0.143	0.134	0.184		2019	-14.25	-43.9	-375.3	416.1
	2020	0.17	0.163	0.142	0.304		2020	334.93	338.68	176.1	442.5
	2021	0.19	0.19	0.14	0.3		2021	368.11	356.88	116.05	443.93
	2022	0.24	0.22	0.15	0.35		2022	405.55	432.39	157.98	452.64
	2023	0.19	0.17	0.14	0.36		2023	360.47	378.29	59.37	470.28
Well 3 Deep	2012	0.07	0.069	0.063	0.092	Well 2 Shallow	2012	298.19	310	120.1	341.5
	2013	0.07	0.066	0.055	0.094		2013	208.00	309	-293.7	461.7
	2014	0.08	0.075	0.057	0.102		2014	155.62	225.5	-269	466.5
	2015	0.08	0.082	0.068	0.106		2015	252.57	325.35	-94.9	419.2
	2016	0.09	0.0855	0.068	0.118		2016	207.31	270.9	-284.9	453.5
	2017	0.08	0.079	0.053	0.116		2017	133.60	206.1	-340.8	419.7
	2018	0.08	0.083	0.064	0.112		2018	87.34	74.7	-357.6	415.8
	2019	0.08	0.082	0.055	0.103		2019	261.99	334.8	-58.6	439.3
	2020	0.09	0.086	0.067	0.11		2020	70.07	58.3	-298.2	419.3
	2021	0.09	0.09	0.06	0.1		2021	73.23	81.035	-264.16	352.08
	2022	0.08	0.08	0.07	0.1	2022	75.69	70.55	-258.71	412.54	
	2023	0.08	0.08	0.08	0.09	2023	95.86	104.76	-356.83	382.78	
Well 4 Deep	2012	0.20	0.204	0.194	0.206	Well 2 Deep	2013	-428.69	-467.7	-499.5	131.9
	2013	0.13	0.112	0.087	0.205		2014	-370.11	-411.1	-495.4	84.4
	2014	0.15	0.152	0.099	0.193		2015	-339.74	-392.1	-486.3	114
	2015	0.15	0.148	0.115	0.184		2016	-384.51	-424.8	-502.1	104.1
	2016	0.17	0.174	0.147	0.184		2017	-350.22	-403.5	-502.9	130.9
	2017	0.18	0.174	0.17	0.197		2018	-404.73	-461.3	-497.2	1.7
	2018	0.17	0.175	0.159	0.184		2019	-324.12	-394.5	-501	118.4
	2019	0.18	0.178	0.173	0.187		2020	-328.86	-367.5	-478.4	50.2
	2020	0.17	0.18	0.127	0.187		2021	-164.04	-152.06	-547.3	186.96
	2021	0.18	0.18	0.13	0.2		2022	-234.42	-280.68	-633.11	111.5
	2022	0.16	0.15	0.13	0.2	2023	-366.70	-379.88	-633.36	115.39	
	2023	0.12	0.13	0.07	0.18	Well 3 Deep	2012	260.80	253.7	210.3	306.5
							2013	364.17	375.9	119.7	442
							2014	388.15	422.1	147.2	448.4
							2015	401.78	440.2	143.2	462.9
							2016	411.14	430.4	179.2	462.7
							2017	364.46	415.6	-82.3	448.7
							2018	402.53	420.1	210.6	461.5
						2019	404.15	432.1	-112.1	587.1	
						2020	367.73	417.4	176.55	448.5	

**Table 12: Summary statistics of Oxidation Reduction Potential at the Residue Storage Area from 2013 to 2023**

Station	Year	Mean	Median	Min	Max
Well 1 Deep	2012	-329.15	-351.9	-361.5	-116.6
	2013	-271.97	-266.4	-375.2	1.7
	2014	-134.61	-250.6	-291	347.3
	2015	-180.86	-251	-481.6	336.2
	2016	-299.52	-333	-426.9	99.8

Appendix – Weather

Well 4 Deep	2021	378.08	390.18	231.02	446.21
	2022	392.08	390.4	212.57	453.35
	2023	369.88	377.33	211.45	448.1
	2012	-220.88	-337.6	-416.4	41.9
	2013	98.46	349.5	-439.7	404.7
	2014	44.72	63.95	-501	370.1
	2015	42.97	35.7	-597.8	378.1
	2016	-243.04	-287.95	-556.5	270.8
	2017	-289.34	-337.65	-500	244.4
	2018	-407.39	-448.5	-553.4	137.1
	2019	-263.38	-337.5	-625.1	341.4
	2020	-257.84	-402.5	-465.7	236.1
	2021	241.54	322.29	-146.76	487.44
2022	444.72	486.45	-6.53	550.4	
2023	335.80	474.93	-86.45	553.68	

**Table 13: Summary statistics of Water Elevation at the Residue Storage Area from 2013 to 2023**

Station	Year	Mean	Median	Min	Max
Well 1 Deep	2012	131.87	131.9028	131.3874	132.1029
	2013	131.98	132.02045	131.4862	132.3255
	2014	131.84	131.874	131.2755	132.2039
	2015	131.95	131.969	131.4605	132.3551
	2016	131.99	131.99605	131.7217	132.3484
	2017	131.94	131.9131	131.6611	132.3429
	2018	132.06	132.0801	131.6035	132.4215
	2019	131.83	131.8278	131.5443	132.1565
	2020	131.96	131.92065	131.6256	133.2077
	2021	132.06	132.02075	131.5125	132.991
	2022	131.99	131.9716	131.4236	132.7686
	2023	131.31	131.5441	127.9078	132.9078
	Well 2 Shallow	2012	113.93	113.9759	113.7681
2013		113.73	113.7411	113.4821	114.0309
2014		113.71	113.7275	113.2566	114.0288
2015		113.72	113.7245	113.4027	113.9952
2016		113.79	113.7951	113.4259	113.9767
2017		113.91	113.901	113.6009	114.1578
2018		113.88	113.8941	113.5997	114.0373
2019	112.86	112.8703	112.5978	113.0763	

Well 2 Deep	2020	112.89	112.9037	112.591	113.0219
	2021	113.99	114.196	112.7591	114.8644
	2022	114.2	114.1629	113.661	115.0802
	2023	114.3051	114.2904	113.6155	115.1724
	2013	113.63	113.6258	113.2945	114.0146
	2014	113.55	113.555	113.2056	113.9122
	2015	113.62	113.6285	113.2716	113.8631
	2016	113.68	113.69145	113.4072	113.9247
	2017	113.58	113.6053	113.2709	113.8921
	2018	113.56	113.5582	113.3718	113.8193
	2019	113.56	113.6087	113.1624	113.8587
	2020	113.60	113.5883	113.274	113.8842
	2021	113.53	113.5511	113.0675	114.1489
2022	113.28	113.2821	112.6758	114.0687	
2023	113.21	113.1770	112.6446	114.1597	
Well 3 Deep	2012	133.86	133.8236	133.3948	134.8782
	2013	133.31	133.3025	132.5933	134.6628
	2014	133.03	133.0324	131.7996	135.0034
	2015	133.21	133.08375	132.4325	134.7668
	2016	132.84	132.8195	132.1306	134.1729
	2017	132.69	132.6946	131.7471	134.1696
	2018	132.76	132.72105	132.1707	134.0623
	2019	132.54	132.46745	131.9998	133.7396
	2020	132.60	132.5027	131.9855	134.9102
	2021	133.16	133.0799	131.8971	135.3918
	2022	133.33	133.3953	131.511	136.5113
	2023	133.69	133.6958	132.3863	135.78
	Well 4 Deep	2012	137.94	137.942	137.7747
2013		138.03	138.07165	137.7004	138.3314
2014		138.09	138.1236	137.6385	138.4492
2015		138.34	138.442	137.585	138.8119
2016		138.56	138.5831	137.6365	139.0317
2017		138.63	138.6243	138.1569	139.1022
2018		138.72	138.7313	138.3324	139.0918
2019		138.46	138.44065	136.1444	138.8241
2020		138.48	138.4218	135.7233	139.8605
2021		138.91	138.86745	128.6848	139.9577
2022		138.1	138.03125	137.0165	138.9141
2023		138.03	137.974	136.0023	139.112

