

Newfoundland and Labrador Hurricane Season Outlook 2021

Submitted to:

Water Resources Management Division
Department of Environment and Climate Change
4th Floor, West Block, Confederation Building
PO Box 8700, St. John's, NL A1B 4J6

Submitted by:

Wood Environment & Infrastructure Solutions,
133 Crosbie Road
PO Box 13216
St. John's, NL A1B 4A5

June 1st, 2021

Wood Project #: ME2112703

Important Notice

This report was prepared exclusively for Water Resources Management Division Department of Environment and Climate Change by Wood Environment & Infrastructure Solutions. The quality of the information, conclusions and estimates contained herein is consistent with the level of effort involved in Wood services and based on: i) information available at the time of preparation, ii) data supplied by outside sources and iii) the assumptions, conditions and qualifications set forth in this report. This report is intended to be used by Water Resources Management Division Department Environment and Climate Change only, subject to the terms and conditions of its contract with Wood. Any other use of, or reliance on, this report by any third party is at that party's sole risk.

Table of Contents

1.0 Atlantic Hurricane Season Outlook 2021 4

2.0 Historical Newfoundland and Labrador Hurricane Season..... 10

3.0 Analog Years and Historical Tracks..... 12

4.0 Newfoundland and Labrador Hurricane Season Outlook 2021... 17

5.0 Closure..... 18

Table of Figures

Figure 1. Nino Regions..... 5

Figure 2. ENSO Probability Forecast (©IRI/CPC)..... 6

Figure 3. Global SST Anomaly on April 27th (above) and May 30th (below), 2021. The red boxes show the MDR region and the purple boxes show the Niño 3.4 region (©NOAA). 8

Figure 4. Tropical storm frequency by year for Newfoundland and Labrador between 1901 and 2020.10

Figure 5. Tropical storm frequency by month for Newfoundland and Labrador between 1901 and 2020.11

Figure 6. Newfoundland tropical cyclone probability using the 1886-2016 historical track (©FSU).12

Figure 7. Upper air (500 hPa) Geopotential Heights (m) composite anomaly (1981-2010 climatology) for June through November of the analogy years (2007, 2005, 1995) (©NOAA).....13

Figure 8. North Atlantic Hurricane Tracking Chart (2007) (©NHC).....14

Figure 9. North Atlantic Hurricane Tracking Chart (2005) (©NHC).....15

Figure 10. North Atlantic Hurricane Tracking Chart (1995) (©NHC).16

Figure 11. Newfoundland and Labrador Hurricane Season Risk Map.17

Table 1. Atlantic basin 2021 hurricane forecast by NOAA, CSU, and TSR.4



1.0 Atlantic Hurricane Season Outlook 2021

June 1st marks the beginning of the Atlantic hurricane season. Although the season stretches for the period between June 1st to November 30th, hurricanes and tropical storms can form prior to or beyond that period. Nevertheless, the season’s climatological activity peak occurs around September 10th each season.

There are numerous sources of weather data used in this hurricane outlook. The major sources are based on the prediction of the El Niño-Southern Oscillation (ENSO) condition, Sea Surface Temperatures (SSTs) model prediction, and weather patterns. The ENSO condition forecast has limited long-range skills. Models also have limited long-range skill in predicting SSTs, vertical wind shear, moisture availability, stability, and predicting weather patterns. The seasonal weather pattern prediction is an outlook to an overall season, but it does not convey the daily and weekly changes in the pattern. The short-term variability in the weather pattern plays a major role in the development, intensification, and track of tropical storms and hurricanes. Furthermore, the intensity and duration of the predicted tropical storms and hurricanes can occur under different combinations of climatic signals, which amplifies the uncertainty of such predictions.

Several agencies issue an Atlantic hurricane season outlook, including the National Oceanic and Atmospheric Administration (NOAA), Colorado State University (CSU), and the Tropical Storm Risk Center (TSR). The NOAA’s 2021 Atlantic Hurricane Seasonal Outlook predicts an above-normal season (60%), followed by a 30% chance of near-normal season and a 10% chance of a below-normal season. Colorado State University (CSU) predicts above-normal activity. It is worth noting that the NOAA uses a 30-year (1981-2010) based-climatology and the CSU has updated their 30-year climatology to include (1991-2020). Meanwhile, Tropical Storm Risk (TSR) predicts that the 2021 Atlantic hurricane season is expected to be 25-30% above the long-term (1950-2020) normal and 5-10% above the recent (2010-2020) ten-year normal. Table 1 below summarizes the prediction from each source.

Table 1. Atlantic basin 2021 hurricane forecast by NOAA, CSU, and TSR.

	NOAA	CSU	TSR
Named Storms	13-20	17	18
Hurricanes	6-10	8	9
Major Hurricanes	3-5	4	4

The El Niño-Southern Oscillation (ENSO) is a measure of the SST anomaly in different Niño regions in the equatorial Pacific, as seen in Figure 1. The neutral phase is determined when the Niño 3.4 region is within ± 0.5 °C. La Niña and El Niño are terms used when the Niño 3.4 region is in the negative and positive phases, respectively.






Figure 1. Niño Regions.

The La Niña phase is climatologically associated with weaker vertical wind shear, weaker trade winds, and increased atmospheric instability across the Atlantic basin, which in turn helps the overall organizational structure of any tropical storm. Meanwhile, the El Niño phase suppresses storm organization due to stronger wind shear, stronger trade winds, and greater atmospheric stability across the Atlantic basin. It is also worth noting that both ENSO phases have opposite effects on the Pacific and Atlantic basins.

Figure 2 shows the probabilistic ENSO forecast based on the Niño 3.4 SST anomaly. The figure shows that Neutral ENSO conditions are more favourable through part of this summer, but the uncertainty in determining the Niño 3.4 SST phase increases as we head into fall. The figure shows the probability of Neutral conditions decreases this fall with an increasing chance of La Niña conditions. The end of the hurricane season favours La Niña conditions, but there exists a chance of the neutral phase still being most dominant. This is an indication of model uncertainty.

Most of the ENSO models are favouring Neutral ENSO conditions for this summer with a wider uncertainty envelope for this fall. This trend is reflected in the above-normal Atlantic basin hurricane season activity discussed earlier.




Figure 2. ENSO Probability Forecast (©IRI/CPC).

Hurricane season predictions consider the various and complex interactions of intra-seasonal variables such as the ENSO and Atlantic Multidecadal Oscillation (AMO), Madden-Julian Oscillation (MJO), mid-level moisture, and Atlantic sea level pressure and upper air patterns. The MJO is an index that measures the variability (1 to 3 months) in the tropical atmosphere. The AMO is a model that measures the decadal SST variability in the Atlantic basin. The current positive phase of the AMO (above normal SST in the North Atlantic) is conducive to lower surface pressures and moist atmospheric mid-levels over the tropical Atlantic.

The latest SST observation (Figure 3) shows a slight warming trend in the tropical Atlantic, near the Main Development Region (MDR), 10-20 °N, and 60-20 °W (red boxes). The figure also shows slight cooling within the Atlantic tropics between April 27th and May 27th. Such relatively cooler temperatures are expected to be associated with drier mid-levels and higher surface pressure in the tropical Atlantic, which in turn decreases the formation of tropical cyclones, at least in the early part of the season. The slightly cooler anomaly in the tropical Atlantic correlates with a weak negative phase of AMO. There is some disagreement between the agencies on the AMO phase, given that the colder SSTs in the North Atlantic, near Newfoundland, are usually more associated with the negative phase of the AMO, as noted by CSU.

The increased cooling trend in the mid-Atlantic basin is of interest, especially in the main Canadian Bioregions. This could be attributed to the combination of the strength of the Labrador Current and the cooler air temperatures that affected the Atlantic in May. Concurrently, there is an increase of SST in the same region due to the Gulf Stream. Historically, the cooling trend in the Canadian Bioregions is likely to subside by July-August and be replaced with warmer SST. The current below-normal SST in the Canadian Bioregions is likely to negatively affect any tropical activity or storm intensifications in Atlantic Canada for the time being.

Figure 3 also shows the near-normal SST anomaly in the Niño 3.4 region, especially just north of the equator and can also be noticed in Figure 3 (purple boxes) between April 27th and May 27th. But it should be noted that this trend is not stable, rather perturbed at the current time due to the upwelling process in the Niño 1+2 (Figure 1).

Figure 3 also shows a cooling trend in the Gulf of Mexico. This trend is likely due to the weather patterns that affected that area during May that brought an ample amount of precipitation.




Figure 3. Global SST Anomaly on April 27th (above) and May 27th (below), 2021. The red boxes show the MDR region and the purple boxes show the Niño 3.4 region (©NOAA).

The latest models are consistently showing warmer SST anomaly in the MDR region and drier conditions across the southern African Sahel region throughout this fall. Perhaps the only exception is the few wet periods across the southern African Sahel Region in July and part of August. Although the warmer SST can fuel thunderstorm clusters, which could eventually lead to tropical storms, below-normal precipitation in the southern African Sahel Region can deter the development of troughs that ultimately aid in storm development off the coast of Africa.

The Wood Seasonal Outlook Team is anticipating an above-normal Atlantic Hurricane Season. The anticipated SST perturbation in the Niño 3.4 region is likely to bring a noticeable monthly variation to the tropical wave activity. The Team also predicts that the main threat of tropical storm development would be on the western Atlantic and the Gulf of Mexico. Such storms are likely to develop and be fueled by the weather patterns and anomalous warmer SSTs, respectively.

2.0 Historical Newfoundland and Labrador Hurricane Season

The archive records from the National Hurricane Centers and the Canadian Hurricane Center records show Newfoundland being impacted by a higher number of tropical storms than Labrador, as seen in Figure 4. The figure shows the frequency of tropical storms by year for Newfoundland and Labrador between 1901 and 2020.




Figure 4. Tropical storm frequency by year for Newfoundland and Labrador between 1901 and 2020.

The same records reveal the month of September as the peak month for tropical storm activity, followed by October (Figure 5).




Figure 5. Tropical storm frequency by month for Newfoundland and Labrador between 1901 and 2020.

Figure 6 shows the Florida State University (FSU) probabilistic tropical cyclones trekking over Newfoundland using 1886-2019 historical data. Storms that affect Newfoundland most regularly approach from the south-southwest and generally pass near or west of Bermuda before arriving on the Newfoundland Coast.




Figure 6. Newfoundland tropical cyclone probability using the 1886-2019 historical track (©FSU).

3.0 Analog Years and Historical Tracks

The Niño 3.4 region SST was in the neutral, negative phase (-0.5°C) throughout April 2021. There has been little evidence of upwelling in May and the region’s SST remains mostly unchanged and still in the neutral phase (-0.5°C). The upwelling process is likely to be intermittent at times, thus, producing a perturbed SST in that region and affecting the weather patterns in the Atlantic Basin.

Given the projected ENSO conditions, the best analog years are 2005, 2013, and 2017. Figure 7 shows the upper air (500 hPa) geopotential height (m) composite anomaly based on 1981-2010 climatology for the period from June and through to November for the combined analog years. The main features in this figure are the anomalously lower heights over NWT and Nunavut, higher heights over Atlantic Canada, and the lower heights over the Atlantic Basin tropics. The progressive weather pattern from west to east is expected to enable the lower heights over NWT and Nunavut to intrude into Atlantic Canada and thus paving the

way for any developed storm to track through Atlantic Canada.




Figure 7. Upper air (500 hPa) Geopotential Heights (m) composite anomaly (1981-2010 climatology) for June through November of the analogy years (2005, 2013, 2017) (©NOAA).

The hurricane track charts for the analog years are given in Figures 8 to 10. Generally, the analog years are depicting an above-normal hurricane season with numerous storms tracking into Atlantic Canada and affecting Newfoundland and Labrador.

The extratropical storms that had a direct impact on Newfoundland and Labrador were Ophelia (Sep 2005) and Nate (Oct 2017). No impact was recorded in 2013. The center of the remnant low pressure of post-tropical storm Andrea dissipated just south of Newfoundland on Sep 13, 2013.




Figure 8. North Atlantic Hurricane Tracking Chart (2005) (©NHC).




Figure 9. North Atlantic Hurricane Tracking Chart (2013) (©NHC).




Figure 10. North Atlantic Hurricane Tracking Chart (2017) (©NHC).

4.0 Newfoundland and Labrador Hurricane Season Outlook 2021

The Nino 3.4 region has experienced pockets of warmer SSTs in early March, eventually caused a fluctuation in the daily SSTs. These pockets were able to further weaken the La Nina setup earlier in the year. The three-month average (Feb-Mar-Apr) SSTs in the Nino 3.4 region indicated a weak La Niña that, although fluctuated from Mar-May, continued weakening further. The latest daily SSTs indicate a neutral phase of ENSO. Wood is anticipating this neutral phase to continue throughout this summer, but it is likely to remain in the negative-neutral phase and continue to oscillate. The projected oscillating SST is likely to provide preferable conditions for storm formation or leading to the weakening/strengthening of developing tropical storms.

The western North Atlantic SSTs also depicted an oscillating pattern from above-normal in March, to below-normal in part of April, above normal in early May, below-normal in mid-May, before bouncing to be above-normal in late May. This region is likely to continue exhibiting such SST oscillation through part of this summer.

Wood anticipates an above-normal Atlantic Hurricane season and a normal to slightly above-normal storm track across Atlantic Canada, specifically affecting Newfoundland and Labrador with extratropical storms. There is a moderate risk of a tropical storm and a low risk of a hurricane making landfall in Newfoundland. Figure 11 depicts the associated risk area. There is a risk of winds gusting up to 140 km/h and 100-130 mm of rainfall associated with any hurricane making landfall.




Figure 11. Newfoundland and Labrador Hurricane Season Risk Map.

5.0 Closure

We trust that this report meets your needs. Please do not hesitate to contact the author if you have any questions or comments regarding the hurricane season outlook.

Yours sincerely,

Wood Environment & Infrastructure Solutions



Robert Giglio

Forecast Lead/Meteorologist

Reviewed by:



Dr. Diar Hassan

Sr. Scientist, Product and Business development

J. Chris Innes

Discipline Lead, Met-Ocean Services