


SYMBOLS

- Geological boundary (defined, approximate, assumed, gradational) . . . . .
- Unconformity (defined, approximate, assumed) . . . . .
- Bedding, tops known (horizontal, inclined, overturned) . . . . .
- Bedding, tops unknown (inclined, vertical) . . . . .
- Igneous foliation: flow banding, compositional layering, autaxitic structure (inclined) . . . . .
- Facing direction on pillow lava (dip known, dip unknown) . . . . .
- Foliation of unknown relative and/or absolute age, only foliation recognized in outcrop: schistosity, cleavage (horizontal, inclined, vertical, dip unknown) . . . . .
- Gneissic layering (horizontal, inclined, vertical, dip unknown) . . . . .
- Composite regional foliation ( $S_m$ ) in pre-Helikian rocks: schistosity, staly cleavage (inclined, vertical) . . . . .
- $S_{m+1}$  foliation (inclined, vertical, dip unknown) . . . . .
- $S_{m+2}$  foliation (inclined) . . . . .
- Mineral rodding lineation, age unspecified . . . . .
- Axis of minor fold, age unspecified . . . . .
- Axis of minor fold of  $S_{m+1}$  . . . . .
- Shape of folded planar feature on flat outcrop surface or sense of symmetry of small folds looking along plunge direction . . . . .
- Structural trend observed on aerial photographs . . . . .
- Lineament observed on aerial photographs . . . . .
- Fault (defined, approximate, assumed) . . . . .
- Thrust (defined, approximate, assumed) . . . . .
- Anticline . . . . .
- Antiform . . . . .
- Syncline . . . . .
- Synform . . . . .
- Overturned anticline, antiform, syncline, synform . . . . .
- Esker (direction of flow known; unknown) . . . . .
- Locality where mineral age has been determined (in millions of years) . . . . .
- Whole-rock age derived from samples collected at various localities in unit (in millions of years) . . . . .
- Mineral occurrence . . . . .
- Drumlinoid ridge . . . . .

- MINERAL OCCURRENCE ABBREVIATIONS**
- |                             |                             |
|-----------------------------|-----------------------------|
| py . . . . . pyrite         | mag . . . . . magnetite     |
| ch . . . . . chalcocopyrite | Fe . . . . . iron formation |
| sp . . . . . bornite        | hem . . . . . hematite      |
| cc . . . . . chalcocite     | fl . . . . . fluorite       |
| gn . . . . . galena         | mo . . . . . molybdenite    |
| sp . . . . . sphalerite     | U . . . . . uranium         |
| pp . . . . . pyrrhotite     | Rb . . . . . rubidium       |
| asb . . . . . asbestos      | G . . . . . gossan zone     |
- ISOTOPIC AGES**
- For whole-rock and mineral ages shown on the map the abbreviations preceding the ages refer to the following methods and minerals:
- |                                                        |
|--------------------------------------------------------|
| Ri . . . . . Rubidium-strontium whole rock isochron    |
| Rr . . . . . Rubidium-strontium whole rock errorchron  |
| KD . . . . . Potassium-argon mineral age on biotite    |
| Kh . . . . . Potassium-argon mineral age on hornblende |
| Km . . . . . Potassium-argon mineral age on muscovite  |
| Pl . . . . . Lead-lead isochron age                    |
| Pip . . . . . Lead-lead mineral age on pitchblende     |



1. MAIN PROVINCE (ARCHEAN)
2. MAKKOVIK SUBPROVINCE (EARLY PROTEROZOIC)
3. GRENVILLE PROVINCE (LATE PROTEROZOIC)

Lines indicate trend of major aeromagnetic elevations and depressions

Geology compiled by B. Ryan, 1979. Sources given in Appendix I of the accompanying report.

Geological cartography by Mineral Development Division, Department of Mines and Energy, Government of Newfoundland and Labrador, St. John's.

Copies of this map may be obtained from the Publications and Information Section, Mineral Development Division, Department of Mines and Energy, P.O. Box 4750, St. John's, Newfoundland A1C 5T7.

Base map assembled and produced by Maritime Resource Management Services Ltd., Antigonish, Nova Scotia. The report has been reviewed to verify intervals by hand from National Topographic Series 1:50,000 sheets with contours in feet. Theoretical north-south orientation of the map is not precise, but are close approximations based on the form of the original contours and elevations given on the 1:25,000 sheets.

None of the names of lakes and rivers shown on this map are of historical usage and are not shown on 1:25,000 maps for this area.

Magnetic declination is approximately 23° westerly at the center of sheet.


Many of the mineral occurrences shown on this map are taken from exploration company maps, and some may not be accurately located at this scale.

All bedrock units exhibit varying degrees of metamorphism and deformation. See accompanying report for description.

This report was financed under the Canada-Newfoundland Mineral Development Subsidy Agreement (1977-1981) by contributions from the Government of Newfoundland and Labrador (15 percent) and from the Department of Regional Economic Expansion (85 percent) and Energy, Mines, and Resources (85 percent) of the Government of Canada.

Legend and symbols are common to Sheet 1 and Sheet 2. Some map units and symbols may not appear on this map.

For an explanation of the use of the foliation designation  $S_m$ ,  $S_{m+1}$ , etc. see Appendix III of accompanying report.



LEGEND

GRENVIAN	<p>40 Flysch series glacial till, Boulder Falls, and sandy outwash. Extensive areas of swamp along Kabinak River.</p> <p><b>SEDIMENTARY, IGNEOUS, AND METAMORPHIC ROCKS OF THE GRENVILLE PROVINCE FORELAND ZONE</b></p> <p><b>GABRO, DIORITE, DIABASE</b></p> <p>39 Medium to fine grained mafic and intermediate intrusive rocks of unknown absolute age.</p> <p><b>SEAL LAKE GROUP</b></p> <p>38 SALMON LAKE FORMATION: Red shales, basaltic flows, and diabase sills.</p> <p>37 WHISKY LAKE FORMATION: Red and grey shales, silt, and argillite; minor and quartzite.</p> <p>36 WUCHUK LAKE FORMATION: Shale, grey to red quartzite, shales, chert, and muscovite; 36a, calcareous and gabbro units.</p> <p>35 BESSE LAKE FORMATION: 35a, white, grey and pink quartzite, arkose, and conglomerate; 35b, amygdaloidal and vesicular basalt.</p>
	<p><b>MICHAEL GABRO</b></p> <p>34 Medium to fine grained alkalic gabbro; actinolite-plagioclase-biotite schist (metagabbro).</p> <p><b>NIPISHISH LAKE INTRUSIVE SUITE</b></p> <p>33 CROOKED RIVER GRANITE: Pink to grey, undeformed to grossly muscovite and biotite bearing aplite and fine to medium grained biotite granite.</p> <p>32 DITEN LAKE-WALKER LAKE GRANITE: 32a, pink to red quartzite, gneiss, and muscovite; 32b, coarse grained, massive to strongly foliated; biotite ± hornblende granite, granodiorite and monzonite; 32c, green to grey medium grained coarse.</p> <p><b>BRUCE RIVER GROUP</b></p> <p>31a Red to pink, quartz-feldspar porphyry.</p> <p>31b Grey to black, plagioclase porphyry. May be intrusive in part.</p> <p><b>SYLVIA LAKE FORMATION:</b> 29a, brown to green, fine brecciated arkosite; 29b, dark green to grey, massive to slightly porphyritic, andesite and basalt; 29c, massive to green, plagioclase porphyritic, andesite and porphyritic; 29d, 29e, amygdaloidal sequence of massive to porphyritic, grey and green siltstone, arkosite, and trachyandesite flows, agglomerate, breccia and bedded tuff (29f), alternating with massive to porphyritic, red, purple, grey, and black siltstone and shale gneissic, breccia, breccia, agglomerate, and bedded tuff (29g); 29i, volcanoclastic mudstone, sandstone and conglomerate (interbedded with above units). Sequence is strongly schistose and recrystallized in the Minnivik Lake - Sperry Lake area.</p> <p>28a BROWN LAKE FORMATION: 28a, red, polymictic conglomerate and minor white oligoclase conglomerate; 28b, pink, red, buff, and green volcanoclastic sandstone, cherty clay silt, minor conglomerate, and siltstone.</p> <p>28b HEGARTY LAKE FORMATION: 27a, massive to well bedded, buff to maroon, arkosic sandstone, minor polymictic conglomerate, and mudstone; 27b, red and grey, tabular to oolitic, polymictic conglomerate; 27c, massive and porphyritic mafic flows, silt and siltstone.</p> <p><b>SOUTHERN KAIPOK VALLEY INTRUSIVE SUITE</b> (relative and absolute ages not firmly established)</p> <p>26 Diorite.</p> <p>25 JUNIOR LAKE GRANODIORITE: Undeformed to weakly foliated, medium grained, hornblende granodiorite to quartz diorite.</p> <p>24 Magmatic, medium to coarse grained, undeformed to weakly foliated, biotite granodiorite.</p> <p>23 Pink, undeformed to weakly foliated, medium grained, potassic granite.</p> <p>22a Grey, medium to fine grained, foliated quartz monzonite.</p> <p>22b Quartz-feldspar-hornblende-biotite gneiss and migmatite.</p>
PALEOHELIKIAN	<p><b>SEDIMENTARY, IGNEOUS, AND METAMORPHIC ROCKS OF THE MAIN PROVINCE AND THE MAKKOVIK SUBPROVINCE FORELAND ZONE</b></p> <p><b>SYNKINETIC AND LATE KINEMATIC GRANITES</b></p> <p>18a ISLAND HARBOUR GRANITE: Pink and grey, medium grained, equigranular to porphyritic, granodiorite. Border zone (20a), contains numerous garnet enclaves and is mylonitic.</p> <p>18b Foliated, medium grained granite, granodiorite and quartz monzonite.</p> <p><b>UNDIVIDED SUPRACRUSTAL ROCKS</b> (lithologically similar to Moran Lake Group equivalents and lower AHB Group)</p> <p>17 Metasedimentary rocks varying in composition from biotite and felsic tuffs to laminated amphibolite.</p> <p>16a Metasedimentary rocks varying in composition from slate, phyllite, and gneiss to garnetiferous muscovite-biotite schist.</p> <p><b>ALLER GROUP</b></p> <p><b>UPPER SEQUENCE</b></p> <p>15a Grey to pink, porphyritic and nonporphyritic phyllite (in part intrusives), igneous, and air-fall tuff, and volcanic breccia.</p> <p>15b Pink, grey and green, laminated, volcanoclastic siltstone and sandstone.</p> <p><b>LOWER SEQUENCE</b> (in tectonic contact with refoliated Archean gneiss)</p> <p>14a, 14b, arkosites, intermediate to mafic metasedimentary flows, silt, bedded tuff, and breccia; 14c, laminated garnetiferous mafic tuff.</p> <p>13a Grey to black, siltstone, mudstone, and slate, locally containing amphibole porphyroblasts, garnetiferous biotite-muscovite schist.</p> <p>13b Bedded white quartzite; minor actinolite and biotite schist.</p>
	<p><b>MORAN LAKE GROUP</b> (in depositional contact with Archean granite and gneiss)</p> <p>12a JOE POND FORMATION: Massive and pillowed basalt, pillow breccia and bedded tuff; minor dolostone and chert.</p> <p>12b WARREN CREEK FORMATION: 12b, grey to black mudstone, slate, siltstone, and sandstone with minor limestone, dolostone and chert; 12b, grey sandstone; 12b, brown dolostone.</p> <p>12c Schistose mafic metasedimentary rocks of diverse origin; bedded tuffs present west of Island Point. Monzonite locally present east of Barlow Lake and north of Island Point.</p> <p>12d Black, granitic, and chlorite-bearing phyllite and slate; minor chert, dolostone, and sandstone.</p> <p><b>MORAN LAKE GROUP EQUIVALENTS</b> (in tectonic contact with refoliated Archean gneiss and granite)</p> <p>11a Schistose mafic metasedimentary rocks of diverse origin; bedded tuffs present west of Island Point. Monzonite locally present east of Barlow Lake and north of Island Point.</p> <p>11b Black, granitic, and chlorite-bearing phyllite and slate; minor chert, dolostone, and sandstone.</p>
ARCHEAN	<p><b>KANAIKOTOK VALLEY COMPLEX</b></p> <p><b>GABRO AND DIABASE</b></p> <p>9a Coarse to fine grained mafic dikes, generally undeformed but locally schistose. May include post-Archean dikes.</p> <p><b>KANAIKOTOK INTRUSIVE SUITE</b></p> <p>7a Massive to weakly foliated to grossly, medium grained, grey to pink, granite, granodiorite, and tonalite.</p> <p><b>ULTRAMAFIC ROCKS</b></p> <p>8 Serpentinized peridotite and talc-carbonate rocks.</p> <p><b>FLORENCE LAKE GROUP</b></p> <p>5 LISSE LAKE FORMATION: Felsic lavali tuff, lapilliferous, minor chert and marble.</p> <p>4 ADLAKOK FORMATION: Isomictic to felsic tuff, volcanic breccia, porphyritic silt, minor marble.</p> <p>3 SCHIST LAKES FORMATION: Massive and pillowed mafic flows and silt, intermediate and felsic tuffs and silt, minor marble.</p> <p><b>MAGGO GNEISS AND WEEKES AMPHIBOLITE</b></p> <p>2a Layered, quartzofeldspathic gneiss with numerous amphibole inclusions; cut by pegmatite and actinolite dikes, and locally highly migmatized.</p> <p>2b Layered amphibolite, locally garnetiferous.</p>
	<p><b>KAIPOK VALLEY COMPLEX</b></p> <p><b>BALLET POND SCHISTS</b></p> <p>10a Sh, quartzofeldspathic and mafic schist secondarily derived from 1', 2', 2a, 2c, and 2d; 10b, interbedded, sh, argillaceous and partially refoliated bedded granite; 10c, schistose leucogranite.</p> <p><b>GABRO AND DIABASE</b></p> <p>6a Coarse to fine grained mafic dikes, locally as swarms. Undeformed to slightly schistose, and calcareous to siliceous in 7', 2a, but commonly recrystallized parallel to superimposed foliation in 2a.</p> <p><b>KANAIKOTOK INTRUSIVE SUITE EQUIVALENTS</b></p> <p>7b Massive to weakly foliated, medium grained, grey tonalite and granodiorite.</p> <p><b>FLORENCE LAKE GROUP EQUIVALENTS</b></p> <p>4 Deformed, felsic lavali tuff, volcanic breccia, and siliceous volcanoclastic sediments; minor chert and marble.</p> <p>3 Deformed pillow lava.</p> <p><b>MAGGO GNEISS AND WEEKES AMPHIBOLITE EQUIVALENTS</b></p> <p>2a 2a, layered, quartzofeldspathic gneiss and amphibolite; 2b, partially retrogressed and refoliated layered gneiss and amphibolite, commonly mylonitic; 2c, grey refoliated migmatite.</p> <p>2b Layered amphibolite, locally garnetiferous.</p>

GEOLOGY OF THE CENTRAL MINERAL BELT  
(CENTRAL PART - SHEET 2)



Published 1982