PROVISIONAL MAP
12-I/6 & 12-I/5

1:50,000

CANADA

EDITION 2

CARTE PROVISOIRE
12-I/6 & 12-I/5



BELLBURNS

ST BARBE SOUTH DISTRICT

NEWFOUNDLAND

Scale 1:50,000 Échelle

Map 85-63

LEGEND

ORDOVICIAN

MIDDLE ORDOVICIAN

$M^{O}GT$ - Goose Tickle Formation or equivalent: dark gray shales and green gray sandstones.

TABLE HEAD GROUP (MIDDLE ORDOVICIAN)

$M^{O}BC$ - Black Cove Formation: black shale.

$M^{O}TC$ - Table Cove Formation: interbedded fine, thin bedded limestones and gray shales; slump beds.

$M^{O}TP$ - Table Point Formation: undivided, TP_{-1} - cyclic limestones and dolomitic limestones on dolostones; TP_{-2} - thickly bedded, light gray to white, dominantly fenestral limestones; TP_{-3} - well bedded, dark gray, fossiliferous limestones with characteristic dolomitic-argillaceous seams; TP_{-d} dolomitized equivalents.

ST. GEORGE GROUP (LOWER-MIDDLE ORDOVICIAN)

$l\text{-}M^{O}A$ - Aguathuna Formation: light gray to gray dolostones with interbedded green-gray and gray shales; locally bodies of matrix breccia ($l^{O}MBr$).

$l^{O}C$ - Catoche Formation: well bedded, gray, fossiliferous and bioturbated limestones with characteristic dolomite mottling, cryptalgal mounds locally; $l^{O}cd$ - diagenetic and epigenetic dolostone replacing Catoche limestones.

$l^{O}BH$ - Boat Harbour Formation: interbedded, dark to light gray, dolomitic limestones and buff dolostones, some cryptalgal limestones, dolomitization common throughout, locally preserved matrix breccia bodies ($l^{O}MBr$).

$l^{O}WB$ - Watts Bight Formation: dark gray to black, fine to medium crystalline dolostone characterized by light gray to cream color mottling, and large cryptalgal mounds.

CAMBRIAN

PORT AU PORT GROUP (MIDDLE-UPPER CAMBRIAN)

$\mathrm{\in}^{U}PJ$ - Petit Jardin Formation: yellow, buff to white weathering, light gray to white, microcrystalline to finely crystalline dolostones, stromatolite mounds common, locally oolitic and stromatolitic limestones in middle of the formation, cherty dolostones towards top, red, green and gray shale interbeds in basal strata.

$M^{\in}MP$ - March Point Formation: dark gray, thinly bedded dolostones, minor dark gray shale.

LABRADOR GROUP (LOWER CAMBRIAN

$l^{\in}HB$ - Hawke Bay Formation: white to pink, clean, quartz sandstones, some argillaceous sandstone units.

$l^{\in}F$ - Forteau Formation: undivided black, oolitic and bioclastic limestone and gray shale; Fsh - basal shale member; Fls - upper limestone member.

$l^{\in}B$ - Bradore Formation: red, arkosic sandstones and pebbly sandstones and gray and pinkish gray, micaceous sandstones and siltstones.

PRECAMBRIAN

PE gr - Pink to red, micaceous, fine to coarse crystalline granite, locally rafts of quartz mica schist.

SYMBOLS

- Unconformity
- Geologic boundary
- Fault; approximate, assumed, dip, downthrow side
- Active mine
- Mineral showing, sphalerite
- No visible mineralization, zinc revealed by zinc reagent liquid
- Diamond drill hole
- Strata and dip of bedding, tops known, tops unknown, inverted horizontal
- Cleavage
- Fold axes
- Outcrop
- Fossil

map 85-63

121(m)