

Hole ID: GB98-01 Location: Stacked at Springdale Company: Nfld Limited
 Core Size: BQ Logged by: T. Rice Date logged: Nov. 27/98

Page: 1 of 1

From	To	Description	Samples	Notes
0.0m	4.5m	Overburden		
4.5m	160.0m	<u>Mafic Volcanic</u> Variable light to dark green-grey fine-grained basaltic flow. Upper 6.0 m may represent crystal tuff. Locally, core is aphyric and has a vitreous luster. Fabric (flow banding) occasionally visible at 70 to 45 degrees to core long axis. Thin (1-3mm) Carbonate veining sporadically orientated throughout interval. Mineralization is sparse throughout interval (trace to 1% pyrite). Locally, up to 2-3% pyrite in interval 118.0m-138.0m. Pyrite appears in thin continuous banded stringers (1-2 mm). Highest concentration of pyritic stringer in section 120.6 - 121.5 (6-8%). Lower contact is sharp and at 40-50 degrees to core axis; ambiguous to which unit is older.	118.4 - 119.8, 120.6 - 121.5, 123.4 - 124.2, 124.2 - 125.3, 125.3 - 126.6, 130.3 - 131.3, 136.1 - 137.1	Moderate mechanically broken core Samples taken previously (No sample #'s recorded)
160.0m	169.2m	<u>Felsic Volcanic</u> Light to dark grey, fine to medium grained, plagioclase porphyritic low-member rhyolite. Porphyroblasts are rounded and range in size from 1-3 mm. Upper 6.0 m of interval is silica saturated and carbonate altered. It is light pink in colour. Local trace bebbly pyrite. EOH		Silica alteration below contact related to structural incompetency along the contact (no gouge noted to indicate a fault, however).