

[illegible]

[illegible]

Copper Hill Resources Inc.											Hole # PH07-03		Page Number 3	
Drilling Company		Collar Elevation	Bearing of Hole from True North	Total Metreage	Dip of hole at collar	Location of Hole in Relation to a fixed point on the claim		Mapsheet	Claim #					
Date Started	Date Completed	Date Logged	Logged By:					Location						
Exploration Co., owner or optionee		Date Submitted	Other:					Property Name						
Metreage From	To	Rock Type	Description	Planar Feature Angle	Core Specimen Metreage (tags)	Sample Number	Sample metreage From	To	Actual Sample Length	Assays				
										Cu (ppm)	Pb (ppm)	Zn (ppm)	Au (g/t)	Ag (g/t)
		Felsic Volcanic (cont)	51.58-51.63 & 53.38-53.51 Two zones with sharp but irregular contacts, lower including fragments of felsic, upper with bands parallel to edge. Appears more intermediate, bit epidote-sericite altered, trace Sp. Intrusive??											
			52.20-52.70 more sericite-epidote rich zone with a few quartz pods. See offset of bands & veins @ 15-10°TCA											
			53.86 break in laminations, probable minor slump/penecontemporaneous deformation @45-50°TCA, hosts Po.		1.00	23240	59.97	60.97	1.00	9	6	809	5	0.2
			61.87-63.65 Mafic dike. Contacts sharp @ 200°TCA (upper) & 35°TCA(lower)											
			Common hair cracks @10-20°TCA hosts minor Py+/-Sp. Right from top of felsics see trace Sp with Po-Py associated, also with green chlorite.											
			Local pygmatic folding of siliceous veins/layers. Jan thinks just compression in soft sediment.											
69.81	120.63	Dacitic tuff & crystal tuff	Banded quartz-sericite-biotite volcanic. Quartz-rich layers more brown/buff in colour. Sericite flaky & very light green. Contains variable crystals of grey quartz or white feldspar.<<1% fine disseminated Py & Po, most on S ₀ , some on low angle cracks. Trace Sp noted near top of unit only.Local siliceous bands may be lapilli or disrupted layers. Overall texture is coarser/flakier than above felsic. (over)	S ₀ 85°										
					1.00	23241	72.24	73.24	1.00	1	2	76	5	0.2
					1.00	23242	84.43	85.43	1.00	3	2	69	5	0.2
							75.55	75.65	petro					
							87.39	87.45	petro					
							104.64	104.71	petro					

[illegible]

[illegible]

[illegible]

[illegible]

Copper Hill Resources Inc.															
										Hole # PH07-03	Page Number 8				
Drilling Company		Collar Elevation	Bearing of Hole from True North	Total Metreage	Dip of hole at collar		Location of Hole in Relation to a fixed point on the claim		Mapsheet	Claim #					
Date Started	Date Completed	Date Logged	Logged By:						Location						
Exploration Co., owner or optionee		Date Submitted	Other:						Property Name						
Metreage From To	Rock Type	Description	Planar Feature Angle	Core Specimen Metreage (tags)	Sample Number	Sample metreage From To		Actual Sample Length	Assays						
									Cu (ppm)	Pb (ppm)	Zn (ppm)	Au (g/t)	Ag (g/t)		
	Felsic Volcanic (cont)	211.45-213.17 grey to brownish, banded felsic volcanic of quartz sericite mainly with grey or white quartz crystals. Has quartz stringers with Po-Cp-Sp-Gn & bands of brownish coarse flakey matted chlorite or biotite with net-like Po-Sp+/-Cp. Thin lamnations/streaks of Sp-Po-Gn. Brown chlorite bands are 1-2cm @90°TCA & irregular mottled zones throughout		1.00	26751	STD CDN-FMC-2									
				0.72	26752	211.45	212.45	1.00	224	462	2800	22	10.3		
				0.84	26753	212.45	213.17	0.72	36	77	378	5	0.7		
				0.84	26754	213.17	214.01	0.84	8	5	205	5	0.2		
				1.04	26755	214.01	215.05	1.00	1	2	113	5	0.2		
				1.00	26756	215.05	216.05	1.00	2	2	248	5	0.2		
		213.17-214.01 grey-brown, fine silty, felsic volcanic. More sedimentary look, no crystals. Composed of quartz-biotite/chlorite-Po 1% disseminated Gt crystals of <=1mm. 5% quartz-chlorite-Sp-Cp-Po stringers/pods @ 90°TCA Minor brassy Po on hair cracks at 0-10°TCA		1.00	26757	216.05	217.05	1.00	3	2	175	5	0.2		
				1.01	26758	217.05	218.06	1.00	2	9	167	5	0.2		
				1.00	26759	218.06	219.06	1.00	4	2	145	5	0.2		
				1.00	26760	219.06	220.06	1.00	4	2	226	5	0.2		
				1.14	26761	220.06	221.20	1.14	6	50	236	5	0.2		
		214.01-221.20 Light grey, quartz phyric, felsic crystal tuff of silica biotite-chlorite-Po. Top 1-2m bit darker, gradationally lightening downhole. Thin sometimes wavy streaks of biotite-Po-chlorite on S. Has variable <= 1-2/3% disseminated Gt crystals to 1.5mm, +/- Po/Cp. Local bit bleached zones with hair cracks have biotite to chlorite. 5-10% quartz-Gt-chlorite-Po/Cp stringers(rare epidote) with coarse Gt. Quartz white and massive or grey coarse.	S ₀ , 90°	1.00	26762	221.20	222.30	1.00	210	139	969	5	2.3		
				1.00	26763	222.30	223.20	1.00	1215	4600	19200	101	25.3		
				0.98	26764	223.20	224.18	1.00	69	84	1542	5	0.7		
				1.00	26765	224.18	225.18	1.00	49	15	1310	5	0.7		
				1.00	26766	225.18	226.18	1.00	101	5	1304	5	0.5		
				0.99	26767	226.18	227.17	1.00	60	37	597	5	0.5		
				1.00	26768	227.17	228.17	1.00	52	28	401	5	0.4		
				1.00	26769	228.17	229.17	1.00	13	2	57	5	0.2		
				0.91	26770	229.17	230.02	0.91	12	3	36	5	0.2		
					26770a	Blank of dike									

Copper Hill Resources Inc.

Hole #										Page Number						
PH07-03										9						
Drilling Company		Collar Elevation		Bearing of Hole from True North		Total Metreage		Dip of hole at collar		Location of Hole in Relation to a fixed point on the claim		Mapsheet		Claim #		
Date Started	Date Completed	Date Logged		Logged By:								Location				
Exploration Co., owner or optionee		Date Submitted		Other:								Property Name				
Metreage From To		Rock Type		Description		Planar Feature Angle	Core Specimen Metreage (tags)	Sample Number	Sample metreage From To		Actual Sample Length	Assays				
												Cu (ppm)	Pb (ppm)	Zn (ppm)	Au (g/t)	Ag (g/t)
				221.20-230.06 white & grey, quartz crystal rhyolitic tuff. Felsic material, minor sericite+/-biotite/chlorite+sulphides. Banding contorted and folded. Very siliceous rock. Has coarse stringers/pods of quartz-chlorite-Sp-Cp?-Po+/-Gn mainly in top 30cm. Other veins and fracture fill stringers of coarse Po(brassy). Zones of matrix Sp-G Cp-Po enclosing felsic grains from 222.21-223.00, but sulphides all through as streaks. disseminated and bands. Sulphide associated with green/black chlorite and/or sericite is less common.												
230.02 236.84		Mafic Dike		Dark grey, strong magnetic, high plagioclase porphyritic mafic dike with granite inclusion. Sharp contacts, lower @ 40°TCA												
236.84 239.73		Felsic Volcanic		Similar to above but greyer. Disseminated Po+/-Cp. Sp in thin wispy lams & few coarser bands ~85-90°TCA. Rock is felsic matrix with sericite, chlorite. Green chlorite & brassy Po or Py on fractures. Sp darker. Better banding of felsic and sericite-chlorite. Quartz crystals.												
239.73 241.61		Mafic Dike		Dark grey, strong magnetic, high plagioclase porphyritic mafic dike												
241.61 242.62		Felsic Volcanic		Green/grey, quartz phyric felsic tuff. Minor folding of laminations.		S ₀ @ 80°										
EOH				Disseminated Po & blebs creamy Sp?												