


MAP 2000-33
WEST MICMAC LAKE
NEWFOUNDLAND
Scale 1 : 50 000



LANDFORM CLASSIFICATION

Each outlined area is assigned a classification consisting of up to three genetic categories and modifiers that designate the types of deposits within each area. Each category within a classification is listed in order of dominance and is separated from the other categories by a slash (/). Typically, the areas are divided so that three landform or deposit types are identified within a given area. The classification system is also used to denote the approximate percentage of landforms occurring within an outlined area, but those which comprise less than 5 percent of the area are not included in the classification. Four variations of the landform system are as follows:

- Where three different landforms are included in a single map unit they are each separated by a single slash (/) and their relative percentages are (00-85), (15-35), and (0-15).
- Where two landforms are included in a single map unit, a double slash (//) or single slash (/) is used to separate them, and their relative percentages are (00-85) and (15-35) for double slash, or (00-85) and (15-40) for a single slash.
- A hyphen between two landform types indicates that they are approximately equal in area. For example, T-Rc indicates that till veneer and rock concealed by vegetation are equal in area.
- A composite symbol is used to show combinations of the above cases. For example, T1C1D1 indicates that about 60-85 percent of the area is covered by fluvial sediment, 15-40 percent by glaciofluvial sediments, and is underlain by till.

The stratum data reported on this map have been referenced from the Newfoundland Stratum Database (Taylor et al., 1994)

LANDFORM CLASSIFICATION: GENETIC

Symbol	Depositional Environment	Origin and Characteristics of Materials
F	Fluvial	Alluvium consisting of silt and clay to boulder gravel; forms terraces and plains associated with modern stream channels, their floodplains and deltas; usually less than 1 m thick, deposited by fluvial action at or below maximum flood levels.
C	Colluvial	Colluvium consists of coarse-grained bedrock derived materials, but may include sand, silt or clay; accumulates on the lower parts, or at the base of steep rock faces; transported by gravity.
E	Aeolian	Medium to fine grained sand and silt, well sorted, poorly compacted; commonly occur as dunes up to 10 m high; transported and deposited by wind.
G	Glaciofluvial	Fine grained sand to coarse grained cobble gravel occur as plains, ridges (eskers), hummocks, terraces and deltas; generally greater than 1 m thick, deposited as outwash in an ice-contact position or proglacially.
L	Lacustrine	Silt, clay, gravel and sand occur as plains and channels; silt and clay deposited in freshwater lakes from lake-shore, sand and silt by lake-floor currents, gravel and sand by shoreline wave action.
M	Marine	Clay, silt, gravel and sand; sand is present in some places, generally moderately to well sorted and commonly stratified, but may be massive; occurs as beach ridges, deltas, terraces and bars deposited in a marine environment; gravel and sand by shoreline wave action; may include shells, clay and silt deposited from suspension and turbidity currents; gravel is generally a w/washed lag.
T	Till	Includes all types of till composed of dominantly unsorted, transported and subsequently deposited by till from glacier ice with no significant sorting by water.
O	Bog	Poorly drained accumulations of peat, peat moss and other organic matter; developed in areas of poor drainage.
R	Rock	Bedrock.

LANDFORM CLASSIFICATION: MORPHOLOGY

Symbol	Morphology	Description
a	apron	A relatively gentle slope at the foot of a steeper slope, commonly used to describe colluvium at the base of a rock escarpment; consists of materials derived from the usually steeper upper slope.
b	blanket	Any deposit greater than 1.5 m thick, minor irregularities of the underlying unit are masked but the major topographic form is still evident.
c	concealed by vegetation	Vegetation matted developed on either colluvium surfaces or a thin layer of angular frost-shattered and frost-heaved rock fragments overlying bedrock; includes areas of shallow (less than 1 m), discontinuous overburden.
d	drummoor	Elongate ridges between 1.5 and 20 m high, 20 and 300 m wide, and 200 to 5000 m long; ridges have a rounded and porous in the up-ice direction and gently curving sides that taper in the down-ice direction; exhibit a convex longitudinal profile, commonly with a steeper slope in the up-ice direction; consist of stratigraphically formed deposits shaped in a streamlined form parallel to the direction of glacial flow; commonly consists of till, although some may contain stratified drift; may have a rock core.
e	eroded and dissected	Series of closely spaced gullies or deeply incised channels; can have a dendritic pattern or may be a single straight or arcuate channel; gullies and channels may contain undrifted streams.
f	fan	A gently sloping accumulation of debris deposited by a stream issuing from a valley onto a levelled, less slope at the mouth of the valley from which the stream issues; the fan shape results from the deposition of material as the stream swings back and forth across the levelled; fluvial fans are usually derived from eroded glacial and glaciofluvial deposits; glaciofluvial fans (lobes) are deposited in standing water rather than a terrestrial environment; colluvial fans are derived from bedrock and are usually steeper (i.e., cone shaped).
h	hummock	An apparently random assemblage of knobs, mounds, ridges and depressions without any pronounced parallelism, sigmoidal form or orientation; formed by glacial melting during ice stagnation and disintegration; includes subglacial, englacial, supraglacial and stratified materials.
k	kettle	A basin or bowl-shaped closed depression or hollow in glacial drift; results from the melting of a buried or partly buried detached block or lens of glacier ice; commonly occurs in association with hummocks.
l	linedated	Elongate spine-shaped ridges between 8 and 60 m high, 70 and 300 m wide and up to 4000 m long; ridges are commonly straight sided, taper at one or both ends, and have a flat longitudinal profile; consist of subglacially formed deposits shaped in a streamlined form parallel to the direction of glacial flow; commonly consist of silt, although some may contain stratified drift; may have a rock core.
p	plain	A comparatively flat, level, or slightly undulating tract of land; materials are either till, glaciofluvial, lacustrine, marine, lacustrine or organic sediments; bedrock features are commonly masked by the overlying sediments.
r	ridge	Narrow, elongated and commonly steep-sided feature that rises above the surrounding terrain; materials are either till, glaciofluvial, fluvial or lacustrine sediments; generally formed by fluvial and glaciofluvial erosion and marine wave action.
t	terrace	Long, narrow, level or gently inclined step-like surface, bounded along one edge by a steeper descending slope or scarp and along the other by a steeper ascending slope or scarp; materials are either till, glaciofluvial, fluvial or lacustrine sediments; generally formed by fluvial and glaciofluvial erosion and marine wave action.
v	veneer	Any deposit less than 1.5 m thick; morphology of the underlying unit is evident.
w	weathered	A thin layer, generally less than 1 m thick, of frost-heaved and frost-shattered bedrock fragments.
x	complex	Commonly used to indicate numerous esker ridges that are closely spaced; can be used when any genetic category exhibits numerous surface expressions in a small area, and in which no single element can be defined.

LANDFORMS AND SURFICIAL GEOLOGY OF THE WEST MICMAC LAKE MAP SHEET (NTS 13K/09)

MAP 2000-33

LANDFORM CLASSIFICATION

MORPHOLOGY (F)	GENETIC									
	Fluvial (C)	Colluvial (E)	Aeolian (G)	Glaciofluvial (L)	Lacustrine (M)	Marine (T)	Glacial (O)	Organic (R)	Rock	
apron (a)	Ca									
blanket (b)	Fb	Cb								Rb
concealed by vegetation (c)		Cc								
drummoor (d)				Td						
eroded and dissected (e)	Fa	Ca	Ea	Ga	La	Ma	Ta			Re
fan (f)	Ff	Cf		Tf						
hummock (h)			Ec	Gc						hc
kettle (k)				Gk						kc
linedated (l)				El						lc
plain (p)	Fp			Gp	Lp	Mp	Tp			Rp
ridge (r)	Fr			Gr	Lr	Mr	Tr			Rr
terrace (t)	Ft			Gt	Lt	Mt	Tt			Rt
veneer (v)	Fv	Cv	Ev	Gv	Lv	Mv	Tv			Rv
weathered (w)				Gw	Lw	Mw	Tw			Rw
complex (x)	Fx	Cx	Ex	Gx	Lx	Mx	Tx			Rx
unclassified	F	C	E	G	L	M	T	O		R

SYMBOLS

- Geological boundary (assumed)
- Scarp face at edge of fluvial terrace
- Esker (flow direction known or assumed, unknown)
- Meltwater channel (small, large)
- Crevasse of major moraine ridge
- Trend of ribbed or minor moraine ridges
- Beach ridges
- Crevasse III ridge
- Sand dunes
- Drummoor
- Crag-and-tail hill
- Futing
- Roché Moutonnée
- Sitation (direction known, unknown)
- Kettle hole (small, large)
- Sinkhole (small, large)
- Observation site

Elevation in feet above mean sea level. Contour interval 50 feet.
NOTE: All symbols and classifications may not occur on this map.


Geology by M.J. Bateman, Geological Survey, Department of Mines and Energy, Government of Newfoundland and Labrador.
Digital Cartography by T.J. Seals and L.C. Ryan, map editing by D.M. Taylor, Geological Survey, Department of Mines and Energy, Geological Survey Branch, 174 pages, Open File NF LD 2195 (version 3).
Copies of this map may be obtained from the Geoscience Publications and Information Section, Geological Survey, Department of Mines and Energy, P.O. Box 8700, St. John's, Newfoundland, Canada, A1B 4X6.
E-mail: pub@zippog.gov.ns.ca
http://www.geoscan.gov.ns.ca

OPEN FILE 13K/09/267
This map supersedes Open File 13K/09/175

PUBLISHED 2000

Reference
Taylor, D.M., St. Onof, L. and Veitch, S.V., 1994. Newfoundland stratum data base. Newfoundland Department of Mines and Energy, Geological Survey Branch, 174 pages, Open File NF LD 2195 (version 3).

Recommended citation:
Bateman, M.J., 2000. Landforms and Surficial Geology of the West Micmac Lake Map Sheet (NTS 13K/09), Labrador. Department of Mines and Energy, Geological Survey, Map 2000-33, Open File 13K/09/267



INDEX MAP