

CONTINUING STRATIGRAPHIC AND TRILOBITE STUDIES OF THE WATTS BIGHT FORMATION (ST. GEORGE GROUP), PORT AU PORT PENINSULA, WESTERN NEWFOUNDLAND

W.D. Boyce, L.M.E. McCobb¹ and I. Knight
Regional Geology Section

¹Department of Geology, National Museum of Wales, Cathays Park, Cardiff, CF10 3NP UK

ABSTRACT

Thick-bedded, stylonodular lime mudstone and wackestone, encasing thin sheets and lenses of grainstone and rudstone, and locally impressive, but isolated thrombolitic boundstone mound complexes midway along the southern coast of the Port au Port Peninsula, host Skullrockian trilobites and other macrofossils. The subtidal shelf facies of the Tremadocian Watts Bight Formation (St. George Group), deposited during transgression through early high-stand, likely were deposited during the Stonehenge transgression defined in the central Appalachians.

*Trilobites from the lower to middle Watts Bight Formation, 1.5 km west of Ship Cove and at Pigeon Head, are dominated by the genera *Millardicurus* and *Sympysurina*. *Sympysurina* sp. cf. *S. convexa*, is recorded for the first time, as is an indeterminate genus, increasing the known diversity of the trilobite fauna from the formation, and strengthening correlation with the Tribes Hill Formation of New York State amongst other co-eval shelf sequences along the eastern and southern margin of Laurentia.*

INTRODUCTION

The trilobite genus *Sympysurina* is long established as an important faunal presence in Furongian to Lower Ordovician rocks of the allochthonous successions of the Cow Head Group (Kindle and Whittington, 1958; Whittington, 1968; Fortey and Skevington, 1980; Kindle, 1982; Fortey *et al.*, 1982; Fortey, 1983; James and Stevens, 1986; Karim, 2008) and the Cooks Brook Formation (Boyce *et al.*, 1992). It is also well represented in co-eval autochthonous platform carbonate shelf sequences throughout the Caledonian–Appalachian–Ouachitan orogenic belt of eastern and southern Laurentia, from Greenland (Poulsen, 1927, 1937; Cowie and Adams, 1957; McCobb and Owens, 2008; McCobb *et al.*, 2011, *in revision*), through Scotland (Ingham *et al.*, 1985), to northwestern Vermont (Shaw, 1951), the Champlain Valley and Mohawk Valley of New York–Vermont (Cleland, 1900, 1903; Fisher, 1954; Landing *et al.*, 2003; Westrop *et al.*, 1993), New Jersey (Weller, 1903; Westrop *et al.*, 1993), Pennsylvania (Raymond, 1910; Butts and Moore, 1936), Maryland (Sando, 1957), West Virginia (Woodward, 1951), Virginia (Orndorf *et al.*, 1988; Taylor *et al.*, 1992), and Oklahoma (Stitt, 1971, 1977, 1983) in the United States. *Sympysurina* and other trilobites, along with brachiopods,

cephalopods, corals, echinoderms and gastropods, were lately discovered in western Newfoundland in the autochthonous Watts Bight Formation, St. George Group along the south coast of the Port au Port Peninsula near Ship Cove (Boyce *et al.*, 2011; McCobb *et al.*, 2011) and in 2012 at Pigeon Head a little farther to the west (Figure 1 and Plate 1). Collectively, the Port au Port Peninsula fossils fill a significant gap in the distribution of Skullrockian faunas in the Ordovician shelf along the Appalachian Orogen.

LITHOSTRATIGRAPHY

The Watts Bight Formation on the Port au Port Peninsula belongs to the older of two long-lived, third-order sequences of Tremadocian and Floian age, separated by the Boat Harbour Disconformity (Knight and James, 1987; Section 2 of Ji and Barnes, 1994; Knight *et al.*, 2007, 2008). Knight *et al.* (2008), supporting the work of Ji and Barnes (1993, 1994), argued that the Tremadocian rocks consist of two sequences lying on either side of an older disconformity within the lower part of the Boat Harbour Formation (see Figure 2). A long-lived sequence including the upper Berry Head Formation, the Watts Bight Formation and the lower member of the Boat Harbour Formation, form a deepening




Figure 1. Geology of the Port au Port Peninsula and location of sections studied in the Watts Bight Formation.



Plate 1. The Pigeon Head composite section with Lower Cove in the background. Arrows indicate the locations of measured sections.

to shallowing Skullrockian succession that is terminated at the older disconformity. Stairisan peritidal carbonate parasequences of the Middle Boat Harbour Formation occur above

the lower disconformity and terminate at the Boat Harbour Disconformity (Knight *et al.*, 2008).

The Watts Bight Formation has now been studied in four sections along the south coast of the Peninsula (Knight *et al.*, 2008; Boyce *et al.*, 2011; *see Figure 1*). In its reference section at Isthmus Bay, it comprises three parts (Knight *et al.*, 2008), a lower boundstone-dominated interval, which includes the celebrated Green Head mound complex (Pratt and James, 1982), a middle interval of burrowed limestone that grades upward into partially dolomitized lime grainstone and an upper interval, once more dominated by a thrombolite-stromatolite mound complex. A markedly incised erosion surface overlain by conglomerate marks the top of the upper interval to separate the Watts Bight Formation from the overlying lower Boat Harbour Formation. The Watts Bight Formation spans the deeper-water (DW) *Cordylodus lindstromi* to *Cordylodus angulatus* Lineage Zones (Ji and Barnes, 1994, figures 18, 19 and 20); the lower member




Figure 2. Stratigraphy of the St. George Group, Port au Port Peninsula from Knight et al. (2008) showing stratigraphic intervals discussed in the text.




Figure 3. Detailed graphic logs of sections through the incomplete Watts Bight Formation at Pigeon Head, showing trilobite localities and conodont localities of Ji and Barnes (1994).

of the Boat Harbour Formation is dated as *Rossodus manitouensis* Zone.

The Watts Bight succession along the mid-southern shore of the Port au Port Peninsula at Ship Cove, Pigeon Head and Lower Cove (*this report* and Boyce *et al.*, 2011) is less structured. Thrombolitic mounds are only present, locally, close to the base of the formation. At Ship Cove and Pigeon Head, stratified muddy limestones abut against, and onlap the mounds, suggesting isolated mound complexes of various sizes. At Lower Cove, mounds are reduced to a few thin beds. Instead, bedded muddy limestone dominates the sections in this area.

At Pigeon Head, the Watts Bight Formation is incomplete and difficult to trace along the cliff and foreshore due to access and several, often subtle, faults. The base of the Watts Bight Formation in the east is covered at sea level, whereas the top of the formation at the Head is cut off by a fault that juxtaposes the middle Watts Bight Formation against rocks of the lower Boat Harbour Formation.

The more easterly of the two sections begins close to the base of the formation (Figure 3). There, massive, clotted and digitate thrombolitic microbial mounds are enclosed within bedded muddy limestone at the base of a cliff. Large cephalopods and gastropods, many partly silicified, and cavities filled by fibrous calcite epitomize the mounds that also host trilobites, articulate brachiopods and silicified gastropod opercula. Macrofossils are also common in limestone flanking the mounds. Well-bedded limestone in the overlying succession in this cliff and at the foreshore to weakly cliffed section at Pigeon Head itself consists of muddy to grainy limestone with some interbeds of burrowed dolostone capped, at Pigeon Head, by a largely dolomitized unit of grainstone that is truncated by the fault mentioned above.

The stratified limestone ranges from decimetre-thick beds of stylonodular lime mudstone to stylolitic to unevenly stratified, variously bioturbated, dolomitic lime wackestone–mudstone with discontinuous lenses of intraclastic and bioclastic grainstone, to beds of intraclastic–bioclastic grainstone, pebbly grainstone and rudstone intercalated, on a fine scale, with burrowed wackestone. Razor sharp, planar erosion surfaces in these grainy beds are likely hardgrounds. The grainy beds locally enclose small isolated microbial mounds. Dolostone beds that are commonly burrow-mottled are interspersed in the interval. The muddy to grainy rock types are repetitively arranged in coarsening-upward parasequences. Trilobites are scattered predominantly in the grainy beds suggesting that they are allochthonous and likely reworked from adjoining muddy sediment into the dominantly intraclastic grainstone during storms. These skeletal beds can be traced for hundreds of metres along strike, and include, in one case, abundant *Sympysurina* pygidia scat-

tered in a fine-grained matrix that hosts millimetre eocrinoid ossicles.

Overlying the 29 m of bedded muddy to grainy limestone at Pigeon Head, is a prominent unit of grainstone, 4 m thick. This grainstone, which is predominantly dolomitized, likely correlates with a similar unit overlying burrowed carbonate at the Isthmus Bay section (the middle interval of Knight *et al.*, 2008). The grainstone is dominantly intraclastic, consisting of well-rounded, very coarse to granular carbonate sand grains, some small pebbles and a scattered bioclast. Horizontal thin stratification marks the base of the unit. It also locally encloses domal microbial mounds as well as large, non-rounded mounds consisting of radial-digitate fan-like structures that project irregularly into the grainstone. Crossbedded to massive grainstone enclosing small mounds overlie the basal bed; scour surfaces are also present.

The Watts Bight succession likely deepened through the lower interval, and commenced shallowing in the middle to culminate at the incision surface as seen at Isthmus Bay. At Ship Cove, the section spans the base through the middle interval of the formation whereas at Pigeon Head except for the mounds from the very top of the lower interval, the succession dominantly belongs to the middle limestone interval.

TRILOBITE BIOSTRATIGRAPHY OF THE WATTS BIGHT FORMATION

A varied trilobite fauna including *Bellefontia*, “*Hystricurus*”, *Millardicurus* and *Sympysurina* has been collected from the Watts Bight Formation on the Port au Port Peninsula at Isthmus Bay, Ship Cove and Pigeon Head. The trilobites occur together with the articulate brachiopod *Finkelnburgia* sp. undet., various unidentified ellesmeroceratid cephalopods, gastropods that include *Ecculiomphalus* sp. undet., *Lytospira*, *Ophileta* sp. cf. *O. supraplana*, *Rhombella* sp. undet. as well as unidentified forms. Very small eocrinoid ossicles are scattered to locally abundant in the grainstones. The trilobites recovered from the formation at Ship Cove and Pigeon Head occur predominantly in rocks equivalent to the middle interval of the formation and the very top of the lower mound interval.

ISTHMUS BAY

A small pygidium of “*Hystricurus*” *ellipticus* (Cleland, 1900) (Plate 2A–D) was recovered from a single horizon approximately 20 m above the base of the Watts Bight Formation at Isthmus Bay, 3 m above the base of the Green Head Mound Complex (*see* Boyce *et al.*, 2011, page 220, Figure 4). A cranidium of *Millardicurus* sp. undet. (Plate 3A) was also discovered during recent re-examination of




Plate 2. “*Hystricurus*” *ellipticus* (Cleland, 1900)?, pygidia. A – D) 2010F001, Isthmus Bay section (NFM F-803); see also Boyce et al. (2011; Plate 19B). Length is about 3mm. A) Dorsal view. B) Posterior view. C) Lateral view. D) Oblique posterior view. E-H). PH-10A, Pigeon Head. Pygidium (NFM F-1001). E) Dorsal view. F) Posterior view. G) Lateral view. H) Oblique posterior view.




Plate 3. *A*) *Millardicurus* sp. undet. from 2010F001, Isthmus Bay section. Latex replica of cranidium (NFM F-994). Dorsal view. *B – F*) *Millardicurus* sp. cf. *M. armatus* (Poulsen, 1937) from 2012F02 = 2010F025, Ship Cove section. *B*) Cranidium, dorsal view (NFM F-995). *C*) *Librigena*, dorsal view (NFM F-996). *D – F*) Pygidium (NFM F-997). *D*) Dorsal view. *E*) Posterior view. *F*) Lateral view.

material from this locality. The trilobites are from the same bed as Z2-3B of Ji and Barnes (1994), which marks the bases of the shallow-water (SW) *Polycostatus falsioneotensis*–*Rossodus tenuis* Assemblage Zone and the deeper-water (DW) *Cordylodus angulatus* Lineage Zone.

SHIP COVE SECTION

An abundant, trilobite fauna occurs in association with brachiopods, cephalopods and gastropods in the Watts Bright Formation, 1.5 km west of Ship Cove (Boyce *et al.*, 2011; Figure 3). Recovered between 11.5 and 33 m above the base of the formation, it comprises the following:

- Bellefontia gyracantha* (Raymond, 1910)?
- "Hystricurus" ellipticus* (Cleland, 1900)?
- Millardicurus* sp. cf. *M. armatus* (Poulsen, 1937)
- Sympysurina myopia* Westrop in Landing *et al.*, 2003

Although no new species were discovered in 2012, better material of *Millardicurus* sp. cf. *M. armatus* (Poulsen, 1937) was obtained from locality 2010F025 (Boyce *et al.*, 2011, Figure 5), including a cranium (Plate 3B), librigena (Plate 3C), as well as the only pygidium (Plate 3D–F); a librigena of *Sympysurina myopia* Westrop in Landing *et al.*, 2003 was also found at locality 2010F031 (Plate 4). Both the librigena and pygidium of *Millardicurus* sp. cf. *M. armatus* are more similar to equivalent sclerites of *M. armatus* than to any other known species of *Millardicurus*, supporting our earlier identification of this species in the formation (Boyce *et al.*, 2011).




Plate 4. *Sympysurina myopia* Westrop in Landing *et al.* (2003) from 2010F031, Ship Cove section. Librigena (NFM F-998), dorsal view.

PIGEON HEAD SECTION

Two sections at, and just east of, Pigeon Head (Figure 3



Plate 5. Grainstone beds at Pigeon Head that host fossil horizons PH-10A = 2012F008 and PH-10C = 2012F010 (arrows).

and Plates 1, 5; Appendix) that lies 1.95 km west of the Ship Cove section, have yielded new fossil horizons. The Pigeon Head section overlaps with the lower part of Section 6 of Ji and Barnes (1994, page 78). The two sections collectively yielded the following trilobites:

- Gen. et sp. undet. (Plate 6)
- "Hystricurus" ellipticus* (Cleland, 1900)? (Plates 2E–F and 7)
- Millardicurus* sp. undet.
- Sympysurina* sp. cf. *S. convexa* (Cleland, 1900) (Plates 8–10)

Sympysurina sp. cf. *S. convexa* (Cleland, 1900) – previously unknown in the Watts Bright Formation – ranges throughout the section at Pigeon Head. At least one of its horizons (PH-14) is a prominent bedding plane replete with scattered pygidia. The bedding plane can be traced over 100 m along strike. A single, unidentified, fragmentary cranium (Plate 6) from horizon PH-10B in the upper part of the section is also new to the formation. It comprises the occipital ring and posterior glabella with parts of the interocular fixigenae, and has a prosopon of fine pustules/tubercles suggesting it may be a hystricurine.

CONODONTS

Ji and Barnes (1994) marked their sample sites in the field with blue painted numbers; of these, Z6-2, Z6-3 and Z6-6 remain visible at the Pigeon Head section. These samples, integrated into the logged section, show that the trilobite fauna overlaps with a collection of conodonts that Ji and Barnes (1994) placed in their shallow-water (SW) *Polycostatus falsioneotensis*–*Rossodus tenuis* Assemblage Zone, which is equivalent to the deeper-water (DW) *Cordylodus angulatus* Lineage Zone of Ji and Barnes (1994). The abun-





Plate 6. Trilobite gen. et sp. undet. from PH-10B, Pigeon Head. Cranidial axis (gabella and occipital ring) with prosopon of fine pustules / tubercles (NFM F-999). A) Dorsal view. B) Posterior view. C) Lateral view. D) Oblique anterior view. E) Closeup of latex replica, showing the prosopon of fine tubercles / pustules.




Plate 7. “*Hystricurus*” *ellipticus* (*Cleland*, 1900)? from PH-14, Pigeon Head. Cranidium (NFM F-1000). A) Dorsal view. B) Anterior view. C) Lateral view. D) Oblique anterior view.

dance of *Polycostatus falsioneotensis*¹ in the bedded limestone of the Pigeon Head section mimics that seen in the middle burrowed carbonate and grainstone interval at Isthmus Bay.

DISCUSSION

The presence in the Watts Eight Formation of *Bellefontia gyracantha* (Raymond, 1910)?, “*Hystricurus*” *ellipticus* (*Cleland*, 1900)? and *Sympysurina myopia* Westrop in Landing *et al.* (2003), support a correlation with the Tribes Hill Formation of New York State, USA (Fisher, 1954) that allows a link with the McKenzie Hill Formation of Oklahoma (see Boyce *et al.*, 2011); *Millardicurus* sp. cf. *M. armata*

(Poulsen, 1937) provides a correlation with the Antiklinalbugt Formation of North-East Greenland (Boyce *et al.*, 2011; McCobb *et al.*, 2011).

Sympysurina sp. cf. *S. convexa* (*Cleland*, 1900), which ranges from PH-02 to PH-14 in the Pigeon Head composite section, strengthens previous correlations, as the type material of *S. convexa* occurs in the Tribes Hill Formation (Fisher, 1954; Westrop *et al.*, 1993). *Sympysurina convexa* also occurs in the Kittatinny Formation of New Jersey where, according to Westrop *et al.* (1993, page 1625), it was originally described as *Illaenurus columbiana* by Weller (1903, pages 133-134; Plate V, figures 1-4); the species has also been reported in West Virginia by Woodward (1951, page

¹ Landing in Landing *et al.* (1996, page 676) regards *Polycostatus falsioneotensis* Ji and Barnes, 1994 as a junior synonym of *Semiacontiodus iowensis* (Furnish, 1938).




Plate 8. *Symphysurina* sp. cf. *S. convexa* (Cleland, 1900). A) PH-14, outcrop, Pigeon Head. Pygidium, dorsal view. B – F). PH-11, Pigeon Head. B – E) Pygidium (NFM F-1002). B) Dorsal view. C) Posterior view. D) Lateral view. E) Oblique posterior view. F) Pygidium (NFM F-1003), silicon cast of external mold, dorsal view.




Plate 9. *Symphysurina* sp. cf. *S. convexa* (Cleland, 1900) from PH-05, Pigeon Head. A) Pygidium (NFM F-1004), clay cast of external mold, dorsal view. B) Pygidium (NFM F-1005), clay cast of external mold, dorsal view.





Plate 10. *Symphysurina* sp. cf. *S. convexa* (Cleland, 1900) from PH-02, Pigeon Head. A) Dorsal view. B) Lateral view. C) Oblique posterior view.

67) from the “Chepultepec–Stonehenge limestone”. Butts and Moore (1936, page 23) report “one pygidium of a trilobite resembling *Tsinania columbiana* (Weller)” from the Stonehenge Formation of the Bellefonte Quadrangle, Pennsylvania. The unidentified fragmentary cranium collected from Pigeon Head (see Plate 6) together with *S.* sp. cf. *S. convexa* indicates greater diversity of the Watts Bight trilobite fauna than suggested by prior studies.

CONCLUSIONS

1. New fossil horizons in the lower Watts Bight Formation on Port au Port Peninsula has yielded two previously unrecorded species, including an indeterminate genus and *Sympysurina* sp. cf. *S. convexa*.
2. *Sympysurina* sp. cf. *S. convexa*, *S. myopia*, *B. gyracantha*? and “*H.*” *ellipticus*?, strengthen the correlation of the lower half of the Watts Bight Formation with the Tribes Hill Formation of New York State, USA.
3. The *Millardicurus* sp. cf. *M. armatus* from the Watts Bight Formation compares well to *M. armatus* from the Antiklinalbugt Formation in North-East Greenland (McCobb *et al.*, *in revision*). This reinforces a likely correlation between the two formations, which is also indicated by the presence of *Sympysurina* species.

ACKNOWLEDGMENTS

Dr. McCobb travelled to western Newfoundland with financial support from the National Museum of Wales, and with the help of a grant from the Geological Association of Canada.

Ruth Gingrich is gratefully acknowledged for finding the first trilobites at Pigeon Head and for important trilobite and brachiopod fossil finds there and at Ship Cove. Helen Gillespie facilitated the use of the photographic setup at the Department of Earth Sciences (Memorial University). The specimens illustrated in this report are housed in the Natural History Annex of The Rooms Provincial Museum (NFM), St. John’s; Adrienne Noftall, Alexanne Oake and Sarah Furlotte assisted in the curation of the material under the supervision of Monica Squires, and Randy Batten (Natural History Collections Manager) provided specimen numbers. Dr. Alana Hinckley reviewed the manuscript. Tony Paltanavage is thanked for his drafting skills.

REFERENCES

- Boyce, W.D., Botsford, J.W. and Ash, J.S.
1992: Preliminary trilobite biostratigraphy of the Cooks Brook formation (Northern Head group), Humber Arm

Allochthon, Bay of Islands, western Newfoundland. In Current Research. Government of Newfoundland and Labrador, Department of Mines and Energy, Geological Survey Branch, Report 92-1, pages 55-68.

Boyce, W.D., McCobb, L.M.E. and Knight, I.
2011: Stratigraphic studies of the Watts Bight Formation (St. George Group), western Newfoundland. In Current Research. Government of Newfoundland and Labrador, Department of Natural Resources, Geological Survey, Report 11-1, pages 215-240.

Butts, C. and Moore, E.S.
1936: Geology and Mineral Resources of the Bellefonte Quadrangle, Pennsylvania. United States Geological Survey, Bulletin 855, 111 pages.

Cleland, H.F.
1900: The Calciferous of the Mohawk Valley. Bulletins of American Paleontology, Volume 13, 26 pages.

1903: Further notes on the Calciferous (Beekmantown) Formation of the Mohawk Valley, with descriptions of new species. Bulletins of American Paleontology, Volume 18, 25 pages.

Cowie, J.W. and Adams, P.J.
1957: The geology of the Cambro-Ordovician rocks of central East Greenland. Part 1. Stratigraphy and Structure. Meddelelser om Grønland, Volume 153, Number 1, 193 pages.

Fisher, D.W.
1954: Lower Ordovician (Canadian) stratigraphy of the Mohawk Valley, New York. Geological Society of America Bulletin, Volume 65, pages 71-96.

Fortey, R.A.
1983: Cambrian-Ordovician trilobites from the boundary beds in western Newfoundland and their phylogenetic significance. In Special Papers in Palaeontology, Number 30, pages 23-27.

Fortey, R.A. and Skevington, D.
1980: Correlation of Cambrian-Ordovician boundary between Europe and North America: new data from western Newfoundland. Canadian Journal of Earth Sciences, Volume 17, pages 382-388.

Fortey, R.A., Landing, E. and Skevington, D.
1982: Cambrian-Ordovician boundary sections in the Cow Head Group, western Newfoundland. In The Cambrian-Ordovician Boundary: Sections, Fossil Distributions, and Correlations. Edited by M.G. Bassett and

- W.T. Dean. National Museum of Wales, Geological Series Number 3, Cardiff, pages 95-129.
- Ingham J.K., Curry G.B. and Williams, A.
1985: Early Ordovician Dounans Limestone fauna, Highland Border Complex, Scotland. *Transactions of the Royal Society of Edinburgh*, Volume 76, Number 4, pages 481-513
- James, N.P. and Stevens, R.K.
1986: Stratigraphy and correlation of the Cambro-Ordovician Cow Head Group, western Newfoundland. *Geological Survey of Canada, Bulletin* 366, 143 pages.
- Ji, Z. and Barnes, C.R.
1993: A major conodont extinction event during the Early Ordovician within the Midcontinent Realm. *Palaeogeography, Palaeoclimatology, Palaeoecology*, Volume 104, pages 37-47.
1994: Lower Ordovician conodonts of the St. George Group, Port au Port Peninsula, western Newfoundland, Canada. *Palaeontographica Canadiana*, Number 11, 149 pages.
- Karim, T.S.
2008: Olenid-dominated trilobite fauna from the Shallow Bay Formation (Cow Head Group), Cambrian-Ordovician boundary interval, western Newfoundland. *Canadian Journal of Earth Sciences*, Volume 45, pages 407-425.
- Kindle, C.H.
1982: The C.H. Kindle collection: Middle Cambrian to Lower Ordovician trilobites from the Cow Head Group, western Newfoundland. *In Current Research, Part C. Geological Survey of Canada, Paper 82-1C*, pages 1-17.
- Kindle, C.H. and Whittington, H.B.
1958: Stratigraphy of the Cow Head region, western Newfoundland. *Geological Society of America Bulletin*, Volume 60, pages 315-342.
- Knight, I. and James, N.P.
1987: The stratigraphy of the Lower Ordovician St. George Group, western Newfoundland: the interaction between eustasy and tectonics. *Canadian Journal of Earth Sciences*, Volume 24, pages 1927-1951.
- Knight, I., Azmy, K., Boyce, W.D. and Lavoie, D.
2008: Tremadocian carbonate rocks of the lower St. George Group, Port au Port Peninsula, western Newfoundland: lithostratigraphic setting of diagenetic, isotopic and geochemistry studies. *In Current Research, Government of Newfoundland Labrador, Department of Natural Resources, Mines Branch, Report 08-1*, pages 115-149.
- Knight, I., Azmy, K., Greene, M.G. and Lavoie, D.
2007: Lithostratigraphic setting of diagenetic, isotopic, and geochemistry studies of Ibexian and Whiterockian carbonate rocks of the St. George and Table Head groups, western Newfoundland. *In Current Research, Government of Newfoundland Labrador, Department of Natural Resources, Geological Survey, Report 07-1*, pages 55-84.
- Landing, E., Westrop, S.R. and Knox, L.A.
1996: Conodonts, stratigraphy, and relative sea-level changes of the Tribes Hill Formation (Lower Ordovician, east-central New York). *Journal of Paleontology*, Volume 70, pages 656-680.
- Landing, E., Westrop, S.R. and Van Aller Hernick, L.
2003: Uppermost Cambrian Lower Ordovician faunas and Laurentian platform sequence stratigraphy, eastern New York and Vermont. *Journal of Paleontology*, Volume 77, pages 78-98.
- McCobb, L.M.E. and Owens, R.M.
2008: Ordovician (Ibex-Whiterock) trilobites from central east Greenland. *In Advances in Trilobite Research. Edited by I. Rábano, R. Gozalo and D. García-Bellido. Cuadernos del Museo Geominero*, 9. Instituto Geológico y Minero de España, Madrid, pages 253-258.
- McCobb, L.M.E., Boyce, W.D. and Knight, I.
2011: Correlation of Lower Ordovician (Ibexian) faunas in north-eastern Greenland and western Newfoundland - new trilobite and biostratigraphic data. *In Ordovician of the World. Edited by J. C. Gutiérrez-Marco, I. Rábano, I. and D. García-Bellido. Cuadernos del Museo Geominero*, 14. Instituto Geológico y Minero de España, Madrid, pages 339-346.
- McCobb, L.M.E., Boyce, W.D., Knight, I. and Stouge, S.
In revision: Early Ordovician (Skullrockian) Trilobites of the Antiklinalbugt Formation, North-east Greenland, and their biostratigraphic significance.
- Orndorff, R.C., Taylor, J.F. and Traut, R.W.
1988: Uppermost Cambrian and lowest Ordovician conodont and trilobite biostratigraphy in northwestern Virginia. *Virginia Minerals*, Volume 34, Number 2, pages 13-20.
- Poulsen, C.
1927: The Cambrian, Ozarkian and Canadian faunas of

- northwest Greenland. *Meddelelser om Grønland*, Volume 70, Number 2, pages 233-343.
- 1937: On the Lower Ordovician faunas of east Greenland. *Meddelelser om Grønland*, Volume 119, Number 3, pages 1-72.
- Pratt, B.R. and James, N.P.
1982: Cryptalgal-metazoan bioherms of early Ordovician age in the St. George Group, western Newfoundland. *Sedimentology*, Volume 29, pages 543-569.
- Raymond, P.E.
1910: Notes on Ordovician trilobites; II, Asaphidae from the Beekmantown. *Annals of the Carnegie Museum*, Volume 7, pages 35-44.
- Sando, W.J.
1957: Beekmantown Group (Lower Ordovician) of Maryland. *Geological Society of America, Memoir* 68, 161 pages.
- Shaw, A.B.
1951: The paleontology of northwestern Vermont. I. New Late Cambrian trilobites. *Journal of Paleontology*, Volume 25, pages 97-114.
- Stitt, J.H.
1971: Late Cambrian and earliest Ordovician trilobites, Timbered Hills and lower Arbuckle Groups, western Arbuckle Mountains, Murray County, Oklahoma. *Oklahoma Geological Survey, Bulletin* 110, 83 pages.
1977: Late Cambrian and earliest Ordovician trilobites, Wichita Mountains area, Oklahoma. *Oklahoma Geological Survey, Bulletin* 124, 79 pages.
- 1983: Trilobites, biostratigraphy, and lithostratigraphy of the McKenzie Hill Limestone (Lower Ordovician), Wichita and Arbuckle Mountains, Oklahoma. *Oklahoma Geological Survey, Bulletin* 134, 54 pages.
- Taylor, J.F., Repetski, J.E. and Orndorff, R.C.
1992: The Stonehenge transgression: A rapid submergence of the central Appalachian platform in the Early Ordovician. In *Global Perspectives on Ordovician Geology*. Edited by B.D. Webby and J.R. Laurie. Proceedings of the the Sixth International Symposium on the Ordovician System, Sydney, Australia. A.A. Balkema, Rotterdam, pages 409-418.
- Weller, S.
1903: Report on Palaeontology. Volume III. The Palaeozoic Faunas. *Geological Survey of New Jersey*. The John L. Murphy Publishing Company, Trenton, New Jersey, 462 pages.
- Westrop, S.R., Knox, L.A. and Landing, E.
1993: Lower Ordovician (Ibexian) trilobites from the Tribes Hill Formation, central Mohawk Valley, New York State. *Canadian Journal of Earth Sciences*, Volume 30, pages 1618-1633.
- Whittington, H.B.
1968: Zonation and correlation of Canadian and early Mohawkian series. In *Studies of Appalachian Geology: Northern and Maritime*. Edited by E-an Zen, W.S. White, J.B. Hadley and J.B. Thompson, Jr. Wiley-Interscience, New York, pages 49-60.
- Woodward, H.P.
1951: Ordovician System of West Virginia. *West Virginia Geological Survey, Volume XXI*, 627 pages.

APPENDIX — Macrofossil horizons in the Watts Bight Formation (St. George Group), Pigeon Head composite section, Port au Port Peninsula, NTS 12B/10 (Stephenville) and NTS 12B/11 (Mainland), UTM Zone 21, NAD27

The fossil horizons are listed in descending order and prefixed by “PH-”. “2010F”, “2012F”, “K-2011-” and “K-2012-” refer to the field collection numbers. “(+)” and “(-)” signify fossil casts and molds, respectively.

PH-17 = 2012F007

Brachiopoda—Articulata

Gen. et sp. undet. – sulcate

PH-16 = 2012F006

Arthropoda—Trilobita

Sympysurina? sp. undet. – questionable pygidium

Echinodermata—Crinoidea

Gen. et sp(p). undet. – debris.

PH-15 = 2012F004

Brachiopoda—Articulata

Gen. et sp. undet. (-)

PH-14 = K-2011-F14 = K-2011-F15 = K-2012-003A-5 = 2012F001 = 2012F003 = 2012F005 = 2012F012 = 2012F013

Arthropoda—Trilobita

Hystricurus “ellipticus” (Cleland, 1900)? – see Plate 7*Sympysurina* sp. cf. *S. convexa* (Cleland, 1900) – flattened and weathered cranidia (+), librigenae (+), thoracic segments (+), pygidia (+) – see Plate 8A

Brachiopoda—Articulata

Gen. et sp. undet.

Echinodermata—Crinoidea

Gen. et sp(p). undet. – abundant ossicles

Mollusca—Gastropoda

Gen. et sp. undet.

PH-13 = K-2011-F13

Undetermined skeletal material

PH-12 = K-2011-F12 = K-2011-F12A = 2012F002 = 2012F011

Arthropoda—Trilobita

Sympysurina sp. cf. *S. convexa* (Cleland, 1900) – cranidium (+), internal mold of a pygidium (-), incomplete pygidium (+)

Ichnofauna

Thalassinoides sp(p). undet.

Mollusca—Gastropoda

Gastropod gen. et sp. undet. – robust, planispiral

Z6-6 of Ji and Barnes (1994, page 92)**PH-11 = K-2011-F04 = 2012F016**

Arthropoda—Trilobita

Sympysurina sp. cf. *S. convexa* (Cleland, 1900) – 1.9 cm long pygidium (+, -) – see Plates 8B-F

Echinodermata—Crinoidea

Gen. et sp(p). undet. – ossicles

Mollusca—Cephalopoda

Gen. et sp. undet. – cross-section of straight form

Mollusca—Gastropoda

Gen. et spp. undet. – cross-sections of large, robust, low spiraled forms

PH-10C = 2012F010 (see Plate 5)

Mollusca—Gastropoda

Gen. et sp. undet.

PH-10B = 2012F009

Arthropoda—Trilobita

Gen. et sp. undet. – glabella (+,-) – see Plate 6

Millardicurus sp. undet. – fragmentary cranidium (+)

PH-10A = 2012F008 (see Plate 5)

Arthropoda—Trilobita

“*Hystricurus*” *ellipticus* (Cleland, 1900)? – pygidium (+) – see Plates 2E-H

PH-09 = 2012F026

Brachiopoda—Articulata?

Gen. et sp. undet.

Mollusca—Cephalopoda?

Gen. et sp. undet. – possible straight form

Mollusca—Gastropoda

Gen. et spp. undet. – specimen clusters were photographed, but not collected

PH-08 = 2012F017

Mollusca—Cephalopoda

Gen. et spp. undet. – straight forms (cross-sections)

PH-07 = 2012F025

Mollusca—Gastropoda

Rhombella sp. undet. – photographed, but not collected

Z6-3 of Ji and Barnes (1994, page 92)

Z6-2 of Ji and Barnes (1994, page 92)

PH-06 = K-2011-F11

Arthropoda—Trilobita

Gen. et sp. undet. – fragments

PH-05 = 2012F018B

Arthropoda—Trilobita

Sympysurina sp. cf. *S. convexa* (Cleland, 1900) – pygidia (+) – see Plate 9

PH-04 = 2012F018A

Arthropoda—Trilobita

Sympysurina sp. cf. *S. convexa* (Cleland, 1900)

PH-03 = K-2011-03-01 = 2012F021 = 2012F024

Arthropoda—Trilobita

Millardicurus sp. undet. – cranidium (-), librigenae (+, -)

Sympysurina sp. cf. *S. convexa* (Cleland, 1900)? — external mold of cranidium

Mollusca—Gastropoda

Gastropod gen. et sp(p). undet. – photographed, not collected

PH-02 = 2012F020

Arthropoda-Trilobita

Sympysurina sp. cf. *S. convexa* (Cleland, 1900) – cranidium (+) – see Plate 10

Brachiopoda-Articulata

Finkelnburgia? sp. undet.

PH-01 = 2012F019

Brachiopoda-Articulata

Gen. et sp. undet.

Echinodermata-Crinoidea

Gen. et sp(p). undet. – debris, including ossicles

Mollusca-Cephalopoda

Gen. et sp(p). undet. – cross-sections

Mollusca-Gastropoda

Gen. et sp(p). undet. – cross-sections of large, robust, low spiraled forms