

MAP 2017-01
FOGO
NEWFOUNDLAND AND LABRADOR



LEGEND

Each outlined area is assigned a classification consisting of up to three generic categories and modifiers that designate the types of deposits within each area. Each category within a classification is listed in order of dominance and is separated from the other categories by a slash (e.g., TuR). Generally, the areas are divided so that up to three landforms or deposit types are identified within a given area. The classification system is also used to derive the approximate percentage of landforms occurring within an outlined area, but those that form less than 5 percent of the area are not included in the classification. Four variations of the landform system are as follows:


- Where three different landforms are included in a single map unit they are each separated by a single slash (/) and their relative percentages are (85-100), (15-30), and (5-15).
- Where two landforms are included in a single map unit, a double slash (//) is used to separate them, and their relative percentages are (85-95)/(5-15) for double slash, or (80-85)/(15-40) for single slash.
- A hyphen between two landform types indicates that they are approximately equal in area. For example, TuRc indicates that till veneer and rock concealed by vegetation or a thin regolith are equal in area.
- A composite symbol is used to show combinations of the above cases. For example, TcR indicates that about 60-85 percent of the area is covered by fluvial sediment, 15-40 percent by glaciofluvial sediments, and is underlain by R.

GENETIC CLASSIFICATION

Symbol	Depositional Environments	Origin and Characteristics of Materials
O	Organic	Poorly drained accumulations of peat, peat moss and other organic matter; developed in areas of poor drainage
F	Fluvial	Alluvium consisting of silt and clay to bouldery gravel; forms terraces and plains associated with modern stream channels; fine floodplains and deltas; usually less than 1 m thick; deposited by fluvial action at or below maximum flood levels
C	Colluvial	Coarse-grained bedrock derived materials; may include sand, silt or clay; accumulates on the lower parts, or at the base of steep rock faces; transported by gravity
E	Aeolian	Medium to fine-grained sand and silt; well sorted; poorly compacted; commonly forms as dunes up to 10 m high; transported and deposited by wind
G	Glaciofluvial	Fine-grained sand to coarse-grained silt/clay; forms plains, ridges (saddles), hummocks, terraces and deltas; generally greater than 1 m thick; deposited as outwash in an ice-contact or proglacial position
L	Lacustrine	Silt, clay, gravel and sand; forms as plains and terraces; silt and clay are deposited in freshwater lakes from suspension; sand and silt by lake-floor currents; gravel and sand by shoreline wave action
M	Marine	Clay, silt, gravel and diatomite; sand is present in some places; generally moderately to well sorted and commonly stratified; may be massive; forms beach ridges, deltas, terraces and bars deposited in a marine environment; gravel and sand are formed by shoreline wave action; may include shells, clay and silt deposited from suspension and turbidity currents; gravel is generally a well-sorted lag
Tv	Glacial	Includes all types of till composed of diamicton; transported and subsequently deposited by or from glacier ice with no significant sorting by water; these include relatively fine (Tv) or coarse (Tc, Td, Tl, Tt) till with little to no surface expression; features produced by wetted flowing ice (Tl, Tl, Tl) or sediment deposited through ice disintegration (Tl, Tl)
R	Rock	Bedrock, either exposed (R) or concealed by vegetation (Rc)

MORPHOLOGY

Symbol	Morphology	Description
a	apron	A relatively gentle slope at the foot of a steeper slope, commonly used to describe collium at the base of a rock escarpment; consists of materials derived from the usually steeper upper slope
b	blanket	Any deposit greater than 1.5 m thick; minor irregularities of the underlying unit are masked but the major topographic form is still evident
c	concealed by vegetation	Vegetation that develops on either colluvial surfaces or a thin layer of angular frost-shattered and frost-heaved rock fragments overlying bedrock; includes areas of shallow (less than 1 m), discontinuous overwash
d	drumstick	Elongate ridges between 1.5 and 20 m high, 20 and 300 m wide, and 200 to 5000 m long; ridges have a rounded end pointing in the up-ice direction and gently curving sides that taper in the down-ice direction; exhibit a commonly longitudinal profile; commonly with a steep slope in the up-ice direction; consist of subglacially formed deposits shaped in a streamlined form parallel to the direction of glacial flow; commonly consist of till, although some may contain stratified drift; may have a rock core
e	eroded and dissected	A series of closely spaced gullies or deeply incised channels; can have a dendritic pattern or may be a single straight or arcuate channel; gullies and channels may contain underfit streams
f	fan	A gently sloping accumulation of debris deposited by a stream issuing from a valley onto a lowland; has its apex at the mouth of the valley from which the stream issues; the fan shape results from the deposition of material as the stream swings back and forth across the lowland; fluvial fans are usually derived from eroded glacial and glaciofluvial deposits; glaciofluvial fans (alluvial fans) are deposited in standing water rather than in a terrestrial environment; colluvial fans are derived from bedrock and are usually steep (i.e., cone shaped)
h	hummock	An apparently random assemblage of knolls, mounds, ridges and depressions without any pronounced parallelism, significant form or orientation; formed by glacial melting during ice stagnation and disintegration; includes subglacial, englacial, supraglacial and stratified materials
k	kettle	A basin or bowl-shaped closed depression or hollow in glacial drift; results from the melting of a buried or partly buried detached block or lens of glacier ice; commonly occurs in association with hummocks
l	lineated	Elongate spindle-shaped ridges between 6 and 60 m high, 75 and 300 m wide and up to 4000 m long; ridges are commonly straight sided, taper at one or both ends, and have a flat longitudinal profile; consist of longitudinally formed glacial drift (a streamlined form parallel to the direction of flow; commonly consist of till, although some may contain stratified drift; may have a rock core; includes slope lineated bogs (L))
p	plain	A comparatively flat level, or slightly undulating tract of land; materials are either till, glaciofluvial, alluvial, marine, lacustrine or organic sediments; bedrock features are commonly masked by the overlying sediments
r	ridge	Narrow, elongated and commonly steep-sided feature that rises above the surrounding terrain; materials are either till, glaciofluvial, fluvial, marine, lacustrine, or organic sediments; includes string bogs (R)
t	terrace	Long, narrow level or gently inclined step-like surface, bounded along one edge by a steeper ascending slope or scarp and along the other by a steeper ascending slope or scarp; materials are either till, glaciofluvial, fluvial or lacustrine sediments; generally formed by fluvial and glaciofluvial erosion or marine wave action
v	veneer	Any deposit less than 1.5 m thick; morphology of the underlying unit is evident
w	weathered	A thin layer, generally less than 1 m thick, of frost-heaved and frost-shattered bedrock fragments
x	complex	Commonly used to indicate numerous either ridges that are closely spaced; can be used where any genetic category exhibits numerous surface expressions in a small area, and in which no single element can be defined at this scale



SURFICIAL GEOLOGY FOGO MAP AREA (NTS 2E/09)

MAP 2017-01

SYMBOLS

Geological boundary	Drumlin (direction known, unknown)
Scarp face at edge of terrace	Crag-and-tail hill
Cirque	Till ramp
Esker (flow direction known or assumed, unknown)	Fluting
Meltwater channel (small, large)	Ridgeline (small, large)
Crestline of major moraine ridge	Thrust of ribbed or minor moraine ridges (numbers indicate relative age)
Beach ridges	Kettle hole (small, large)
Sand dunes	Sea-hole (small, large)
Avalanche track	Observation site
	Delta
	Recolonization age

Note: All symbols and classifications may not occur on this map.

Geology by K.R. Sheppard.
GIS/digital cartography by T.J. Sears.
Digital elevation data supplied by the Shuttle Radar Topography Mission (SRTM), a partnership between NASA and the National Geospatial Intelligence Agency (NGA). Flown aboard the NASA Space Shuttle Endeavour (11 - 22 February 2000). Additional information available from: <http://seamless.usgs.gov/products/srtm/>.
The age dates and glacial stations, where included, on this map have been obtained from Taylor (2001a, b).
Elevation in feet above mean sea level. Contour interval 50 feet.
Copies of this map may be obtained from the Geoscience Publications and Information Section, Geological Survey, Department of Natural Resources, Government of Newfoundland and Labrador, P.O. Box 8700, St. John's, NL, Canada, A1B 4X8.
This map is subject to review and revision. Comments to the author concerning errors or omissions are invited.
Based on maps published by Surveys and Mapping Branch, Department of Natural Resources, Ottawa, Canada.
OPEN FILE G2E/09/1820.
This map supersedes Map 2011-21, Open File G2E/09/1699.


PUBLISHED 2017.
Department: <http://www.nr.gov.nl.ca/nr/>
Geological Survey: <http://www.nr.gov.nl.ca/nr/mines/geoscience/>
E-mail: info@geogov.nl.ca

References:
Taylor, D.M. (Compiler).
2001a. Carlton 14 data list for Newfoundland and Labrador. Government of Newfoundland and Labrador, Department of Mines and Energy, Geological Survey. <http://rps.geosurv.gov.nl.ca/>.
2001b. Newfoundland and Labrador Station Database. Government of Newfoundland and Labrador, Department of Mines and Energy, Geological Survey. <http://rps.geosurv.gov.nl.ca/>.

Recommended citation:
Sheppard, K.R.
2017. Surficial geology of the Fogo map area (NTS 2E/09). Government of Newfoundland and Labrador, Department of Natural Resources, Geological Survey, Map 2017-01. Open File G2E/09/1820.

Note:
Open File reports and maps issued by the Geological Survey Division of the Newfoundland and Labrador Department of Natural Resources are made available for public use without being formally edited or peer reviewed. They are based upon preliminary data and evaluation. The purchaser agrees not to provide a digital reproduction or copy of this product to a third party. Derivative products should acknowledge the source of the data.

Disclaimer:
The Geological Survey, a division of the Newfoundland and Labrador Department of Natural Resources (the "authors and publishers"), retains the sole right to the original data and information found in any product produced. The authors and publishers assume no legal liability or responsibility for any alterations, changes or misrepresentations made by third parties with respect to these products or the original data. Furthermore, the Geological Survey assumes no liability with respect to digital reproductions or copies of original products or for derivative products made by third parties. Please consult with the Geological Survey to ensure originality and correctness of data and/or products.



Digital Elevation Model (DEM) from Shuttle Radar Topography Mission (SRTM) of the Fogo map area (shaded from the northeast). The image varies surface information not readily illustrated on the surficial map. Terrain variability is evident, with areas of bedrock highlighted as rougher textured areas (consistent with surficial map), and areas of thicker till shown as smoother textured areas.